A HANDBOOK

OF

CRYPTOGAMIC BOTANY

BY

ALFRED W. BENNETT, M.A., B.Sc., F.L.S.

LECTURER ON BOTANY AT ST THOMAS'S HOSPITAL

AND

GEORGE MURRAY, F.L.S.

SENIOR ASSISTANT, DEPARTMENT OF BOTANY, BRITISH MUSEUM
AND EXAMINER IN BOTANY, GLASGOW UNIVERSITY

WITH 378 ILLUSTRATIONS

LONDON
LONGMANS, GREEN, AND CO.
AND NEW YORK: 15 EAST 16th STREET
1889

All rights reserved

PREFACE.

In presenting to the botanical public this 'Handbook of Cryptogamic Botany,' the result of the labour of several years, the authors are deeply sensible of its inevitable defects. In traversing so wide a field, it is impossible that a single worker, or even two, can be practically acquainted with more than a comparatively small portion of it. It is necessary, therefore, to consult a literature, the extent of which, even for a single year, is appalling, and in which it is often difficult to distinguish between trustworthy and untrustworthy observations. The attempt has, notwithstanding, been made by the authors to acquaint themselves with the contents of every important publication of recent years bearing on Cryptogamic Botany, and issued in English, French, German, Italian, or Latin. It is beyond hope but that inaccuracies have crept in, or that observations which should have been noted have escaped attention. We shall be grateful to workers and writers who will inform us of any such inaccuracies or omissions, and especially to those who will kindly supply us, with a view to future editions, with copies of papers containing records of new and original observations or theories. Those relating to Vascular Cryptogams, Muscineæ, Algæ, and Schizophyceæ should be directed to Mr. Bennett; those relating to Fungi, Mycetozoa, and Schizomycetes to Mr. Murray; these being the portions of the work actually written respectively by each of us, although we hold ourselves severally responsible for the whole contents of the volume.

So rapidly are facts accumulating, and new views of affinity being promulgated, that it may be necessary to change one's opinion on some points even in the interval between the printing of the earlier and later sheets of a volume like this; and this must be held to account for any

slight discrepancies that may be apparent between the general scheme of classification contained in the Introduction, and the details as carried out in the work itself.

We have also to acknowledge the permission given by the publishers of the following works for electros to be taken from the illustrations contained in them, viz. :—De Bary, 'Comp. Morph. und Biol. der Pilze, Mycetozoen, und Bacterien,' and 'Vorlesungen über die Bacterien'; Sachs, 'Lehrbuch der Botanik'; Goebel, 'Grundzüge der Systematik'; Luerssen, 'Die Kryptogamen'; Schenk, 'Handbuch der Botanik'; Zopf, 'Die Spaltpilze': Hauck, 'Die Meeresalgen'; Reinke, 'Lehrbuch der Botanik'; Thomé, 'Lehrbuch der Botanik'; Le Maout et Decaisne, 'Traité Général de Botanique'; Solms-Laubach, 'Einleitung in die Palæophytologie.'

Of the remaining illustrations, many have been taken from nature; others have been copied from the illustrations of previous works, especially from Cooke's 'British Freshwater Algæ'; and for others we have to thank the courtesy of the Councils of the Royal and Linnean Societies, and the publishers of the 'Annals of Botany.'

In those branches of Cryptogamic Botany which have not been the immediate object of our own researches, we have freely consulted experts in these several departments, and have received from all the greatest kindness and most valuable assistance. In particular we wish to express our obligations in this respect to Mr. W. Carruthers, Pres. L.S., F.R.S.; Mr. J. G. Baker, F.R.S.; Professor F. O. Bower, F.L.S.; Dr. R. Braithwaite, F.L.S.; Mr. E. M. Holmes, F.L.S.; and Mr. G. C. Karop, F.R.M.S.

December, 1888.

ALFRED W. BENNETT,
6 Park Village East, London, N.W.

GEORGE MURRAY,
British Museum (Natural History),
Cromwell Road, London, S.W.

CONTENTS.

	PAGE
INTRODUCTION	1
FIRST SUBDIVISION: VASCULAR CRYPTOGAMS	10
HETEROSPOROUS VASCULAR CRYPTOGAMS	21
CLASS I. RHIZOCARPEÆ	21
,, II. SELAGINELLACEÆ	38
ISOSPOROUS VASCULAR CRYPTOGAMS	53
CLASS III. LYCOPODIACEÆ	53
,, IV. FILICES	64
,, V. OPHIOGLOSSACEÆ	95
,, VI. EQUISETACEÆ	100
FOSSIL VASCULAR CRYPTOGAMS	114
SECOND SUBDIVISION: MUSCINEÆ	132
CLASS VII. MUSCI	136
,, VIII. HEPATICÆ	156
FOSSIL MUSCINEÆ	172
THIRD SUBDIVISION: CHARACEÆ	173
CLASS IX. CHARACEÆ	173
FOSSIL CHARACEÆ	183
FOURTH SUBDIVISION: ALGÆ	184
CLASS X. FLORIDEÆ	191
,, XI. CONFERVOIDEÆ HETEROGAMÆ	219
,, XII. FUCACEÆ	228
,, XIII. PHÆOSPOREÆ	237
,, XIV. CONJUGATÆ	258
,, XV. CONFERVOIDEÆ ISOGAMÆ	272
,, XVI. MULTINUCLEATÆ	280
,, XVII. CŒNOBIEÆ	291
FOSSIL ALGÆ	303

PAGE

FIFTH SUBDIVISION: FUNGI 305

GROUP I. PHYCOMYCETES 323

CLASS XVIII. OOMYCETES 323

,, XIX. ZYGOMYCETES 335

GROUP II. SPOROCARPEÆ 353

CLASS XX. ASCOMYCETES 353

,, XXI. UREDINEÆ 383

,, XXII. BASIDIOMYCETES 388

SIXTH SUBDIVISION: MYCETOZOA 401

CLASS XXIII. MYXOMYCETES 401

,, XXIV. ACRASIEÆ 405

SEVENTH SUBDIVISION: PROTOPHYTA 407

GROUP I. SCHIZOPHYCEÆ 408

CLASS XXV. PROTOCOCCOIDEÆ 409

,, XXVI. DIATOMACEÆ 419

,, XXVII. CYANOPHYCEÆ 426

GROUP II. SCHIZOMYCETES 449

CLASS XXVIII. SCHIZOMYCETES 449

INDEX 457

Errata

Page 11, line 22, *for* oophore *read* oophyte
,, 11, ,, 22, *for* sporophoie *read* sporophyte
,, 122, description of fig. 94, for *Haulea* read *Hawlea*
,, 186, line 10, *for* CŒNOBIÆ *read* CŒNOBIEÆ
,, 187, ,, 12, for *Mesocarpeæ*, *Desmidieæ* read *Mesocarpaceæ*, *Desmidiaceæ*
,, 187, lines 19, 24, *for* Desmidieæ *read* Desmidiaceæ
,, 187, line 26, *for* Mesocarpeæ *read* Mesocarpaceæ
,, 190, ,, 4, *for* Desmidieæ *read* Desmidiaceæ
,, 208, last line, *for* HYPNÆACEÆ *read* HYPNEACEÆ
,, 209, line 1, *for* Hypnæa *read* Hypnea
,, 209, ,, 4, *for* Rytiphlæa *read* Rytiphlœa
,, 213, description of fig. 191, line 1, for *corymbosa* read *corymbifera*
,, 250, ,, ,, ,, 223, ,, 2, *for* propagules *read* sphaceles
,, 280, line 6 from bottom, *for* Dasycladæceæ *read* Dasycladaceæ
,, 296, last line, for *polyhedra* read *polyhedræ*
,, 311, description of fig. 271, line 1, *for* Lib. *read* de By.
,, 313, ,, ,, ,, 275, ,, 1, for *Portulaceæ* read *Portulacæ*
,, 319, line 3, omit full stop after Johow
,, 326, description of fig. 287, line 3, also last line of page, for *Portulaceæ* read *Portulacæ*
,, 335, line 2, insert comma after Pythium
,, 343, ,, 15 from bottom, *for* Tremellini *read* Tremellineæ
,, 381, ,, 13, *for* Barenetski *read* Baranetzki

HANDBOOK

TO

CRYPTOGAMIC BOTANY.

INTRODUCTION.

No general handbook to Cryptogamic Botany has appeared in the English language since the Rev. M. J. Berkeley's in 1857. Since then this department of botanical science has gone through little less than a revolution. Not only has the number of known forms increased enormously, but additions of great importance have been made to our knowledge of structure by the use of the microscope, and to the genetic connection of different forms by the careful following out of the life-history of particular species. The present work is an attempt to bring within the reach of botanists, and of the public generally who are interested in the study of nature, an acquaintance with the present state of our knowledge in this branch of science. It is not intended to replace in any way the numerous excellent handbooks or monographs which exist of special families or groups. Its scope is quite different. Neglecting the minor differences by which genera, or in many cases even orders, are distinguished from one another, the aim of the authors has been to bring before the reader the main facts of structure, of development, and of life-history, which mark the larger groups, contrasting them with one another, and referring only to the broader lines of demarcation within those groups. It is hoped that the work will be found useful to the beginner as well as to the more advanced student.

One great difficulty in our work has been to observe a due proportion in the space allotted to the different groups; and this has been increased by the necessity for a very different mode of treatment in the higher and the lower forms. Of the Vascular Cryptogams—more nearly allied in many respects to Phanerogams than to the lower Cryptogams—our knowledge is, with some exceptions, as minute and exhaustive as that

16

of Flowering Plants ; and it is improbable that any living forms remain to be discovered differing in any material point of structure from those already known. Here, therefore, we are able to discuss systems of classification which claim something like finality ; and the difficulty of the compiler of a handbook is the enormous amount and the minute detail of the material to his hand, from which he has to cull those portions which seem suitable for his object. In order not to extend this portion of the work beyond due limits, it has been necessary frequently to practise rigid compression—beyond, probably, what many of our readers specially interested in these groups would have desired. The same remarks apply, to a large extent, to the Muscineæ. But in the Thallophytes, and especially in the lower Algæ and Chlorophyllous Protophyta, the case is very different. From the extremely minute size of many of these, and the much smaller extent to which they have been studied, new forms are constantly being discovered, and important additions are yearly being made to our knowledge of their life-history and of their structure. It is highly probable that among these groups, as well as in some of the orders of Fungi, forms will yet be discovered which cannot be assigned to any type at present known, gaps in the life-history of many species will yet be filled up, and organisms hitherto placed in widely separated families will ultimately be found to be phases in one cycle of development. We have therefore, in this branch of our subject, brought before our readers every fact of importance known to us which is vouched for by observers in whom we have confidence ; and the classification here submitted is a purely tentative one. In the Algæ, the Fungi, and the Protophytes, we do not attempt an exhaustive enumeration of orders or families which shall include every known organism, but describe in detail only those types which are of greater importance, and of which our knowledge is more complete.

Something must be said on the classification adopted. In the Vascular Cryptogams and in the Muscineæ this proceeds on generally recognised lines, in which there is not much room for difference of opinion. But a very different treatment seemed necessary of the Thallophytes, and of the relationship to one another of the Algæ and Fungi, and of the different orders within each of these groups. Here the systems proposed are almost as numerous as the original investigators, and it has been necessary to choose that which appeared to the authors to bring together those organisms which are most nearly related to one another. Whether these two familiar terms represent a natural bifurcation in the classification of the lower organisms, is a question which has been very variously answered by different observers and theorisers. About fifteen years ago a system of classification of the Thallophytes was pro-

posed, on authority entitled to the highest respect,[1] which altogether abolished the bifurcation into Algæ and Fungi. On this system the sole character made use of in their primary classification was the mode of reproduction. First came the Protophyta, in which no sexual mode of reproduction is known, followed by three primary classes (in ascending order)—the Zygosporeæ, Oosporeæ, and Carposporeæ—distinguished solely by the degree of complexity of the sexual process. Each of these four classes was then divided into a series containing chlorophyll and a series not containing chlorophyll, the former including the organisms hitherto known as Algæ, the latter those hitherto known as Fungi.

In support of this view it was urged, with great plausibility, that, reproduction being the most important event in the life-history of a plant, the mode in which this is brought about must become fixed in each group by heredity ; while such a subordinate character as the presence or absence of chlorophyll is seen, in the higher plants, to be entirely without importance in determining affinity. But a little consideration will show that it is unsafe to apply the same rule to more highly and to less highly organised forms. In the higher forms of life the mode of sexual reproduction becomes, in its main features, absolutely fixed ; and throughout the vast range of Angiosperms—as in the higher animals—there is entire uniformity in this respect in all important points ; while in external morphology, and in the mode in which they obtain their livelihood, there is the greatest diversity, even within a narrow circle of affinity. In the animal kingdom we may point, as an illustration of this law, to the existence of such a family as the Cetacea among Mammalia; among flowering plants we have only to consider such phenomena as the occurrence of parasitism, insectivorous habits, or the suppression of chlorophyll, in individual genera dispersed through a large number of natural orders. Even in subsidiary characters connected with the process of reproduction there is not the uniformity that might have been expected. While such an apparently subordinate point as the number of cotyledons in the embyro is so constant as to give its name to primary divisions of Phanerogams, a character which might have been supposed to be much more important (but which, it is instructive to observe, is connected with the mode in which the germinating embryo receives its nutriment)—viz. the presence or absence of endosperm—is not always constant, even within narrow limits. The first necessity of a nascent organism is to live ; and hence it is not surprising to find that in the lower forms of life the one character which remains most constant within wide circles of affinity is the mode of life. In the course of development

[1] See Sachs's *Text-book of Botany*, 2nd English edition, p. 244.

of the higher forms nature may be said to have tried a variety of experiments in the mode of reproduction ; on the whole there is a continual advance, but still by no means infrequent fallings back to simpler modes ; and unless this law of *retrogression* is taken into account, any system of classification must be *pro tanto* imperfect and misleading.

If these considerations have any weight, it is not surprising that, although the system of classification of Thallophytes above alluded to has been adopted by a few authorities in this country and on the Continent, it has not met with general acceptance. The adoption of its leading principle, that '*in each class* Fungi have diverged as ramifications from various types of Algæ,'[1] is seen to lead to such startling results as the collocation in the same class of Spirogyra and Mucor, of Volvox and Peronospora, of Callithamnion and Agaricus. It may, on the contrary, be safely asserted that several of the most important groups among Fungi (take, for example, the Uredineæ and the Basidiomycetes) display no traces of genetic affinity with any known class of Algæ ; and if, on the other hand, we have forms like Saprolegnia and Chytridium among Fungi, or Leptothrix and Beggiatoa among Protophyta, which betray strong indications of a degraded affinity with groups of Algæ, this by no means contradicts the general law that Fungi as a class form an altogether independent series.

Retrogression may take the form of the suppression of either the vegetative or the reproductive organs ; and wherever you have one of these sets of organs displaying strong development, while the other set of organs is very feeble or altogether wanting, you have *primâ facie* evidence of retrogression. Of this examples will be given in the sequel.

While, therefore, we adopt the Protophyta as a primary class, with the general limits proposed by Sachs, we have no hesitation in reverting to the time-honoured division of the higher Thallophytes into the two great groups of Algæ and Fungi.

The classification of Fungi adopted is that of de Bary, consisting of a main series (called the series of the Ascomycetes), composed as follows : (1) Peronosporeæ (with Ancylisteæ and Monoblepharis), (2) Saprolegnieæ, (3) Mucorini or Zygomycetes, (4) Entomophthoreæ, (5) Ascomycetes, (6) Uredineæ ; and of divergent groups as follows : (7) Chytridineæ, (8) Protomyces and Ustilagineæ, (9) Doubtful Ascomycetes (Saccharomyces, &c.), (10) Basidiomycetes.

The groups 1–4 are Phycomycetes, and 7 and 8 of the second series go with them ; while 9 stands in relation to 5, and 10 to 6; and they are

[1] Sachs's *Text-book*, p. 244, foot-note.

so considered together in the linear series in which they come in the book. The Phycomycetes approach the Algæ (Chlorophyceæ) very nearly; and the other groups of Fungi bear a relation to the Phycomycetes which seems to negative any supposition of their independent connection with algal forms.

One other point had to be decided, whether to commence at the bottom or at the top of the series. Had our purpose been to construct theoretically a genealogical tree for the lower forms of vegetable life, the former course must necessarily have been pursued, and in the laboratory there is no doubt much to be said in favour of proceeding from the simple to the more complicated types. But to the general student, 'from the known to the unknown' is a very sound principle. And, among flowerless plants, not only are the higher types far the best known to the ordinary observer, but they are also those about the life-history of which we have the greatest certainty of knowledge. We have been confirmed in our belief of the correctness of this decision by observing that in the last edition of Huxley and Martin's 'Elementary Biology' these authors have (in the zoological section) abandoned the ascending for the descending order.

The question of terminology is one of the greatest stumbling-blocks to the student of cryptogamy. Not only are new terms being constantly introduced, many of them quite needlessly or from an erroneous idea of structure; but some that are in continual everyday use are employed in different senses by different writers of repute. The first requisite in a terminology, after accuracy, is simplicity; and to this end we have, wherever possible, used anglicised instead of Latin and Greek forms. Many of the terms which we employ throughout this volume—such as *sporange*, *archegone*, *antherid*, *cœnobe*, *sclerote*, *epiderm*, &c.—will probably be accepted at once; and it seems strange that the awkward and uncouth foreign forms of these words should have held their ground so long. With others there will no doubt be greater hesitation; but we hope to see all, or nearly all, of the anglicised forms we have used gradually introduced into all English works on cryptogamic botany, and the same principle possibly extended in other cases where we have not ventured to apply it.

A striking instance of the uncertainty which still surrounds cryptogamic terminology is afforded by the various senses in which different writers use the everyday term 'spore.' Le Maout and Decaisne and Asa Gray speak of spores as 'the analogues of seeds;' Berkeley describes the unfertilised oospheres of Fucus as spores; Vines includes under the term all reproductive cells produced either asexually or sexually; while Sachs defines a spore as a reproductive cell produced

either directly or indirectly by an act of fertilisation, reserving the term 'gonidium' for those which are produced without any previous act of impregnation. It is obvious that one practical defect of this last suggestion is that it may necessitate a perpetual change of terminology as our knowledge advances. Every fresh extension of the domain of sexual fecundation—and it is probable that many such will take place—will involve the removal of a fresh series of reproductive cells from the category of gonidia to that of spores, even though they may not be the immediate result of an act of fertilisation. Again, if the spores of ferns and mosses are the indirect result of impregnation, it is difficult to say why the term should not ultimately include all reproductive bodies whatever, except the spores of the 'apogamous ferns' with which Farlow and de Bary have recently made us acquainted, and of other similar abnormal productions, which are certainly not the result of impregnation, direct or indirect.

It seems a sounder principle—and is certainly more convenient to the student—to base a system of terminology on facts which can be confirmed by actual observation, rather than on unproved hypotheses. We propose, therefore, as the basis of our terminology, to restore the term *spore* to what has been in the main hitherto its ordinary signification, and to restrict its use to *any cell produced by ordinary processes of vegetation, and not directly by a union of sexual elements, which becomes detached for the purpose of direct vegetative propagation.* The spore may be the result of ordinary cell-division or of free-cell-formation. In certain cases (*zoospore*) its first stage is that of a naked primordial mass of protoplasm. In rare instances it is multicellular, breaking up into a number of cells (*polyspore*, composed of *merispores*, or breaking up into *sporids*).

The simple term *spore* will, for the sake of convenience, be retained in Muscineæ and Vascular Cryptogams; but in the Thallophytes it will generally be used in the form of one of those compounds to which it so readily lends itself, expressive of the special character of the organ in the class in question. Thus, in the Protophyta we have *chlamydospores*; in the Myxomycetes, *sporangiospores*; in the Saprolegnieæ and many Algæ, *zoospores*; in the Uredineæ, *teleutospores*, *æcidiospores*, *uredospores*, and *sporids*; in the Basidiomycetes, *basidiospores*; in the Ascomycetes (including Lichenes), *ascospores*, *polyspores*, and *merispores*; in the Diatomaceæ, *auxospores*; in the Œdogoniaceæ, *androspores*; in the Florideæ, *tetraspores*; and others belonging to special groups. The cell in which the spores are formed will, in almost all cases, be called a *sporange*; and this term will be compounded in the same way as spore.

In describing the heterosporous Vascular Cryptogams it is usual to speak of the spores which give rise to the female prothallium and those which give birth to antherozoids as 'macrospores' and 'microspores' respectively. The first of these terms is doubly objectionable: firstly, etymologically, the proper meaning of μακρός being not 'large,' but 'long;' and secondly, from the close similarity in sound of the two terms, an inconvenience, especially in oral instruction, which every teacher must have experienced. Seeing that the correct and far preferable terms *megaspore* and *microspore* are used by Berkeley, Areschoug, Carpenter, and others, it is difficult to understand how 'macrospore' can ever have got into general use; and these terms, together with *megasporange* and *microsporange*, will be used in the following pages. For similar reasons *megazoospore* is always used instead of 'macrozoospore.'

The *male* organs of fecundation are so uniform in their structure throughout Cryptogams that very little complication has found its way into their terminology. The cell or more complicated structure in which the male element is formed is uniformly known among Cormophytes as well as Thallophytes as an *antherid*; the fecundating bodies are almost invariably naked masses of protoplasm, provided with vibratile cilia, endowed with apparently spontaneous motion, and bearing the appropriate name of *antherozoids* or 'spermatozoids.' The former of these is perferable for two reasons; from its etymological connection with antherid, and because the use of terms compounded from 'sperm' should, for reasons to be detailed presently, be avoided for male organs. In only two important groups, Florideæ and Ascomycetes, are the fecundating bodies destitute of vibratile cilia and of spontaneous motion: in the former case they are still usually termed 'antherozoids;' in the latter 'spermatia,' and their receptacles 'spermogonia.' In order to mark the difference in structure from true antherozoids, it is proposed to designate these motionless bodies in both cases *pollinoids*; the term 'spermogone' is altogether unnecessary, the organ being a true antherid.

A satisfactory terminology of the *female* reproductive organs presents greater difficulties, from the much greater variety of structure, and the larger number of terms already in use. The limits we have placed to the use of the term 'spore' and its compounds require the abandonment of 'oospore' for the fertilised ovum or oosphere in its encysted state (enclosed in a cell-wall), anterior to its segmentation into the embryo; and this is the most important change involved in the terminology of the present volume.

In devising a term which shall include all those bodies which are the immediate result of impregnation, it was necessary to take two points specially into account. Firstly, the term must be capable of

defence on etymological grounds ; and secondly, it must, like 'spore,' be suited for ready combination. After much consideration we have decided on adopting the syllable *sperm*. No doubt the objection will present itself that the Greek σπέρμα, like the Latin 'semen,' while originally meaning the ultimate product of fertilisation, came afterwards to signify the male factor in impregnation ; and hence, in zoology, terms derived from these roots are used for the male fertilising bodies. But the objection applies to a much smaller extent to phyto-terminology, and the use in the proposed sense of the syllable 'sperm' is justified by the universal employment in phanerogamic botany of such terms as 'gymnosperm,' 'angiosperm,' 'endosperm,' and 'perisperm.' Of cryptogamic terms, where the syllable is used in the reverse sense, 'sperm-cell,' for antherozoid or pollen-grain, has never come into general use in this country ; 'spermatozoid' is easily replaced by 'antherozoid ;' 'spermogonium' is simply a peculiar form of antherid, and 'spermatium' has already been referred to. Accepting this term as the least open to objection of any that could be proposed, it will be found to supply the basis of a symmetrical system of terminology, which will go far to redeem the confusion that at present meets the student at the outset of his researches. For the unfertilised female protoplasmic mass the term *oosphere* is already in general use ; and, though not all that could be desired, it is proposed to retain it. The entire female organ before fertilisation, whether unicellular or multicellular, is designated by a set of terms ending in *gone*, such as *archegone* and *carpogone*, again following existing analogy.

The term *reproduction* itself is often far too vaguely employed by botanical writers. We propose to limit its use, in accordance with its etymology, to the production of a new individual, that is, to a process of impregnation ; all cases of non-sexual multiplication being described as *propagation*.

The object of the writer of a handbook is to gather up and to collate material already existing, winnowing, to the best of his judgment, the wheat from the chaff. Except, therefore, where original observations may have been made by the compiler himself, it will contain nothing new. In compiling from the writings of the original observers it was thought best, as far as possible, to use their own words, and this will account for the frequent close resemblance in the following pages of the descriptions contained to those in such works as de Bary's 'Comparative Anatomy of the Phanerogams and Ferns,' the 'Comparative Morphology and Biology of the Fungi, Mycetozoa, and Bacteria,' by the same writer, the scheme of which has been mainly adopted in outline, and Goebel's 'Outlines of Classification and Special Morphology.' Admirable, on

the whole, as are the translations of these works by which the Clarendon Press has enriched English scientific literature, it is the original work rather than the translation that we have in all cases followed. We wish here to express the great obligation under which we lie to these writers, and to acknowledge the extent to which we have borrowed from them. In the chapter on Fossil Vascular Cryptogams we have, to a considerable extent, followed Graf zu Solms-Laubach's excellent 'Einleitung in die Palæophytologie,' though with some modifications. To the description of each group or family we have appended a bibliography of the researches on which that description is founded ; these having again been consulted, wherever possible, in the original language.

FIRST SUBDIVISION.

VASCULAR CRYPTOGAMS.

The Vascular Cryptogams include all the highest forms of cryptogamic life, and constitute a well-marked group of plants intermediate between the Gymnosperms, or lowest division of Flowering Plants, and the lower or Cellular division of Flowerless Plants. From the former they differ mainly in the mode in which fertilisation is effected ; from the higher forms of the latter in the much greater differentiation of tissues. The term 'vascular' Cryptogams is, however, strictly speaking, correct only to a limited degree. Although the arborescent and fruticose species display as well-marked a differentiation of their tissues as Flowering Plants, into epidermal tissue, 'vascular' bundles, and fundamental tissue, and the bundles consist of distinct xylem and phloem (without any intermediate cambium, as in Gymnosperms), it is only rarely, as in that group of Flowering Plants, that the xylem is composed of vessels in the true sense of the term.

The Vascular Cryptogams and the highest families of Cellular Cryptogams are distinguished from Flowering Plants by an obvious Alternation of Generations between Sexual Generation or *Oophyte*, and Non-sexual Generation or *Sporophyte*. The former is a small and purely cellular structure, usually of very temporary duration, the purpose of which is to bear the sexual organs of reproduction, male *antherids* and female *archegones*, the structure of which is uniform in all essential characters throughout the class, and which are borne on a cellular expansion, the *prothallium*. This prothallium may be either monœcious or diœcious—that is, the male and female organs may be borne on the same or on different prothallia. The act of fertilisation consists in the impregnation of an *oosphere*, a naked mass of protoplasm contained within the *central cell* of the archegone, by one or more *antherozoids*, minute masses of protoplasm endowed with spontaneous motion by means of vibratile cilia, which escape from the cells of the antherid and penetrate to the central cell of the archegone. The immediate result of the impregnation of the oosphere is that it invests itself

with a cell-wall of cellulose, and thus becomes an *oosperm*, which develops into the *embryo*, and finally into the sporophyte, which is often of great size and extended length of life. It is this that is commonly known in popular language as the Fern, Club-moss, &c. On it are produced, without any process of fertilisation, the *spores*, which are always single cells, protected by two or three coats, and giving rise on their part, when they germinate, to the oophyte. This is the cycle of development in the Isosporous families of Vascular Cryptogams. In the Heterosporous families there are produced (always on the same plant) two different kinds of spore, differing very greatly in size—larger *megaspores* and smaller *microspores*. The former produce female prothallia, that is, those which bear archegones only; the latter male prothallia, bearing antherids only, or even antherozoids, without the intervention of a male prothallium or antherid. The spores are always endogenous structures, produced by free-cell-formation within a spore-case or *sporange*, which again, in the heterosporous families, is a *megasporange* or a *microsporange*, according as it contains megaspores or microspores. The sporanges are often collected into groups known as *sori*, and these may again be enclosed within special chambers, *sporocarps* or *conceptacles*. Various kinds of vegetative propagation, similar to those of Flowering Plants, also occur on both oophyte and sporophyte; and in certain exceptional cases either the oophore or the sporophore may be entirely suppressed, constituting the phenomena of *apogamy* and *apospory* respectively. These will be described especially under Ferns, where they are of the most common occurrence.

It may be useful in this place to compare the structure of the organs of reproduction and the phenomena of impregnation in Vascular Cryptogams with those of Phanerogams, and to endeavour to trace the requisite homologies,[1] although these cannot be fully understood without some knowledge of the details of structure described under the separate families. It has been usual to compare the pollen-grain of Phanerogams with the antherozoid of Vascular Cryptogams; but this is not strictly accurate. The essence of the act of fecundation consists in the coalescence of the protoplasmic contents of an active (male) and of a passive (female) cell. The protoplasmic endoblast or cell-contents in each case must, therefore, be homologous; that is, the antherozoid with the protoplasmic contents of the pollen-grain which, in Flowering Plants, is brought into contact with the embryonic vesicle by means of the pollen-

[1] Hofmeister was the first to point out these homologies in his *Vergleich. Unters. über Keimung u.s.w. der höheren Kryptogamen* (see the Ray Society's translation, *On the Germination, &c., of the Higher Cryptogamia*, p. 438). They have since been traced out more completely by Hanstein, Celakovsky, and others.

tube ; or probably the more correct homology is that of the nucleus of the antherozoid with the 'generative' nucleus of the pollen-grain. The most important difference between Cryptogams and Phanerogams lies in the mode in which this contact between the male and female elements is brought about. In Flowering Plants it takes place by the penetration into the embryo-sac, through the micropyle of the ovule, of the extension of the inner coat of the pollen-grain known as the pollen-tube, excited into activity by the viscid secretion of the stigma. The pollen-grain has therefore in the first place to be conveyed from the anther to the stigma ; and the various parts of the flower of Flowering Plants are all more or less concerned, directly or indirectly, with arrangements for facilitating the conveyance of pollen. In Flowerless Plants, on the contrary, contact between the male and female elements is effected by the protoplasmic contents escaping from the male cells and coming directly into contact with the oosphere in consequence of an independent power of motion imparted to these naked masses of protoplasm (antherozoids) by the vibratile cilia with which they are provided. This impregnation always takes place in water or moisture, and no external agency is needed to bring it about. Hence the absence from all Flowerless Plants of any conspicuous compound organ analogous to the flower of Phanerogams. The true homology of the pollen-grain of Phanerogams appears to be with the microspore of the heterosporous Vascular Cryptogams, notwithstanding the fact that the contents of the microspore break up into a number of antherozoids, each capable of impregnating an oosphere ; while the pollen-grain, as a rule, emits only a single pollen-tube. Here, as in other phenomena, we are guided to the true homology by comparing the highest Cryptogams with the lowest Phanerogams, the Gymnosperms, which form a connecting link between them and Angiosperms. In all Gymnosperms, as in some Angiosperms, the pollen-grain is divided into several cells, only one of which (very much larger than all the rest) emits a short pollen-tube. The large fertile cell in the pollen-grain of Gymnosperms corresponds to the larger fertile portion of the microspore of Selaginellaceæ, to the entire contents of the microspore of Marsileaceæ, to the terminal cell of the germinating filament in Salviniaceæ, and to the antherid of the isosporous Vascular Cryptogams. In Angiosperms the sterile cells of the pollen-grain of Gymnosperms appear to be sometimes entirely suppressed, and the pollen-grain becomes unicellular. The contents of the pollen-grain of Angiosperms, together with the intine or inner coating of the grain, are therefore homologous with the antherid of Cryptogams. Seeing that the motile antherozoids have to be conveyed to the oosphere through the medium of water, it is convenient for both antherid and archegone to be freely

exposed to the action of moisture at the time of maturity. Hence, in all the isosporous Vascular Cryptogams, by far the largest portion of the product of germination of the hermaphrodite spore is the cellular tissue or prothallium (reduced to a comparatively small size in the Lycopodiaceæ and Ophioglossaceæ), on which are borne both the antherids and the archegones. In the heterosporous Salviniaceæ the male prothallium is reduced to a simple unseptated germinating filament; and in Marsileaceæ it altogether disappears. In Selaginellaceæ it takes the form of the small sterile cells at one extremity of the microspore; in Gymnosperms, of the sterile cells of the pollen-grain; in Angiosperms it is almost entirely suppressed. The contents of the pollen-grain corre-



FIG. 1.—I., male catkin of *Zamia* (Cycadeæ). II. III., antheriferous scale and pollen-sacs.

FIG. 2.—Peltate scale of *Equisetum*, with sporanges *sg*. (After Sachs.)

sponding to the antherozoids, the pollen-grain itself becomes homologous with the microspore, the pollen-sac or anther-cell to the microsporange. Even in external appearance the pollen-sacs of Coniferæ and Cycadeæ bear a striking resemblance to the sporanges of some Vascular Cryptogams. The modes of formation of the pollen-grains within the pollen-sac and of spores within the sporange, from an original archespore, are identical in their main features.

Turning now to the female organs of reproduction, we must trace the homology of these back from the product of the union of the two elements, which in all the higher plants, whether flowering or flowerless, may be termed the oosperm, developing later into the embryo. In

Flowering Plants this is the product of the action of the contents of the pollen-grain on the protoplasmic embryonic or germinal vesicles ; in the higher Flowerless Plants, of the action of the antherozoid on the oosphere, or protoplasmic contents of the central cell of the archegone. The naked masses of protoplasm known as the germinal vesicles (or perhaps rather that one only which is ultimately fecundated) are therefore unquestionably homologous with the naked mass of protoplasm known as the oosphere. In Vascular Cryptogams the central cell which contains the oosphere is a portion of an archegone which is borne on a prothallium resulting from the germination of a spore or of a megaspore, as the case may be. To understand the homologies with the higher Phanerogams we must again have recourse to the intermediate Gymnosperms. In



FIG. 3.—Fertilisation of *Abies* (Coniferæ). *p*, pollen-grains ; *ps*, pollen-tube ; *e*, embryo-sac ; *c*, secondary embryo-sac or corpusculum. (× 60.)



FIG. 4.—Female prothallium of *Marsilea*, with archegone *a* and oosphere *o*. (After Hanstein.)

this class the female elements are not directly developed in the interior of the embryo-sac, but within certain chambers produced within the embryo-sac, the secondary embryo-sacs or 'corpuscula, the homologues of the central cell of the archegone. In connection with them there have been detected other structures comparable to the neck and the canal-cells of the archegone. The object of the peculiar structure of the archegone and the deliquescence of the canal-cells being to facilitate the passage of the motile antherozoids to the oosphere, these are no longer wanted when impregnation is effected by means of a pollen-tube. The archegone (except the central cell) is therefore reduced to a rudimentary condition in Gymnosperms, and disappears altogether in Angiosperms, where the embryonic vesicles are produced directly within

the embryo-sac, the homologue of the megaspore. The nucellus of the ovule must then be regarded as corresponding to the megasporange ; but it is difficult to carry the homology further. In Filices and Lycopodiaceæ an hermaphrodite prothallium usually exposes both kinds of sexual organ to the action of moisture ; in Equisetaceæ we find a normal differentiation into male and female prothallia, but produced from one kind of spore only ; in the heterosporous families the differentiation is carried back to the spores and sporanges, and the female prothallium is altogether a subordinate product, and never has any separate existence apart from the megaspore, within which it is more or less concealed, only that part which bears the archegones being exposed. In Angiosperms it has been suggested that we have a rudiment of the female prothallium surviving in the peculiar 'antipodal cells' found within the embryo-sac in certain natural orders ; but the homology is doubtful. In the Selaginellaceæ we find also the rudiments of two other structures which characterise the ovule of Flowering Plants. The sterile tissue which occupies the lower part of the megaspore in this order is probably the first appearance of the endosperm (albumen) which is found in the seed of a large number of Flowering Plants ; the purpose in both cases being the same, to provide the embryo with nutritive material during the early stages of its growth. In all Phanerogams the young embryo is borne on a longer or shorter pedicel of cellular structure, the suspensor or proembryo, which, again, we find for the first time in the same order of Cryptogams.

In all the isosporous Vascular Cryptogams the sexual generation or oophyte has an independent existence distinct from the spore which produced it ; while in the heterosporous families the prothallium recedes more and more into the background, existing only within the megaspore ; and at the same time the male organs or antherids become more rudimentary in structure. In Gymnosperms the female prothallium and the antherids (as distinct from the antherozoids) have become much reduced, and in Angiosperms have completely disappeared. While, therefore, in Characeæ alternation of generations disappears by the suppression of the non-sexual generation which bears the spores, in Phanerogams the same result is brought about by an exactly opposite process, the suppression of the sexual generation or prothallium with its antherids and archegones, and the coalescence of male and female elements takes place within the non-sexually produced embryo-sac of the ovule, which corresponds to the megaspore.

The life-history and general structure of the various organs in Vascular Cryptogams may now be described more in detail.

The immediate product of the germination of the spore or megaspore

is always a *prothallium*, which is usually a green plate of tissue lying flat on the soil, very commonly lobed or reniform in shape, sometimes microscopic, but often quite visible to the naked eye, from $\frac{1}{4}$ to $\frac{1}{2}$ an inch in diameter, and constituting, with its organs of reproduction, the oophore, *oophyte*, or sexual generation. The prothallium usually disappears as soon as the non-sexual generation has firmly rooted itself in the soil; but in Gymnogramme (Filices) it attains a larger size, and continues its existence for a considerable time, producing a succession of reproductive organs. The oophyte never becomes differentiated into stem and leaves, as in the Muscineæ, nor does it contain any vascular tissue. It usually consists of only a single layer of cells filled with chlorophyll, except a marginal cushion, where there are several layers. Here it puts out into the soil numerous colourless elongated cells, the organs of attachment, or *rhizoids*,[1] among which, or scattered over the whole of the under surface and margin, are the archegones and antherids. True vegetative budding takes place but rarely on the prothallium; but it sometimes, in certain Filices, exhibits apogamy, the sporophore springing directly from it, without the intervention of the sexual organs. In some Lycopodiaceæ and in the Ophioglossaceæ the prothallium is subterranean, destitute of chlorophyll, and cylindrical or tuberous. In the Hymenophyllaceæ (Filices) it is often filiform. In the isosporous families the prothallium is most commonly monœcious, less often dicœcious. In the heterosporous families it is far less fully developed. That which arises from a megaspore is very small, formed within the spore, and at no period maintains an independent existence; while that developed within a microspore is still more rudimentary.

The *archegones* of Vascular Cryptogams are produced on the prothallium, usually on the under side of the cushion. Each archegone consists of a swollen basal portion or *venter*, and a *neck*, usually composed of four longitudinal rows of cells; the venter is buried in the tissue of the prothallium, the neck alone projecting above it. The archegone originates from a superficial cell of the prothallium, which divides by a tangential wall into an inner and an outer cell; the latter then develops, by further divisions, into the four rows of cells constituting the neck, which is, therefore, always comparatively short. The inner cell puts out a protuberance between the neck-cells, which is first of all separated as the *neck-canal-cell*, and below it a small portion is again separated from the lower larger cell as the *ventral canal-cell*; the

[1] These organs are frequently termed root-hairs; but it is better to confine this term to the epidermal appendages (trichomes) of the roots of Phanerogams and of the sporophyte generation of Vascular Cryptogams, between which and true rhizoids there is a functional rather than a morphological homology.

lowermost portion of the lowest cell remains of a comparatively large size, and is the *central cell* containing the *oosphere*. When mature, the two canal-cells deliquesce into mucilage, which swells up considerably, drives apart the four apical lid-cells or *stigmatic cells* of the neck, and is ejected ; an open canal being thus formed to allow the access of the antherozoids to the oosphere, which always takes place in moisture, the ejected mucilage assisting also in this process.

The *antherids* appear as roundish papillæ on the margin, or dispersed over the under surface of the prothallium ; in some cases they are imbedded in its tissue. Each antherid consists of a comparatively small number of cells ; when mature the cell-walls are ruptured under water, and from each escapes a swarming *antherozoid*. The antherozoids are spirally coiled threads of protoplasm, the body of which is formed from the nucleus of the mother-cell, with a number of fine *vibratile cilia* on the anterior coils. There is generally attached to each antherozoid, as it escapes from its mother-cell, a vesicle of protoplasm containing starch-grains, formed out of the cytoplasm of the mother-cell, which, adhering to one of its posterior coils, is dragged along with it during its swarming, but becomes detached before its entry into the neck of the archegone. In the heterosporous families the antherid is of very simple structure, and is either produced directly within the microspore, or after the preliminary formation of a few cells, which must be regarded as a rudimentary prothallium.

The form and size of the non-sexual generation or *sporophyte* vary within very wide limits, from the filmy, moss-like Hymenophyllaceæ (Filices) to the arborescent tree-ferns (Filices), and must be described more in detail under the various families. It arises in the archegone, from the *oosperm* or fertilised oosphere. The first effect of impregnation is that the oosphere invests itself with a cell-wall of cellulose, thus becoming the oosperm, which then divides into a small number of undifferentiated cells, in which condition it is known as the *embryo*. In the earliest subsequent divisions of the embryo may be recognised the rudiments of the first *root*, of the first leaf or *cotyledon*, and of the apex of the *stem* ; while at the same time a lateral outgrowth termed the *foot* is formed at the bottom of the venter, and draws from the prothallium the first nourishment for the young plant. The venter at first grows vigorously, enveloping the embryo, until this latter finally protrudes free, leaving the foot still attached to it for some time as a nutritive organ. The primary root soon disappears, and in some Hymenophyllaceæ, and in Salvinia and Psilotum, is not followed by others ; but in the great majority of cases other true roots succeed in acropetal succession, and the prothallium then disappears. The cotyledon always

remains small ; the first stem bends upwards, and other leaves of a more complicated structure appear on it.

The *roots* of Vascular Cryptogams usually arise in acropetal succession on the stem (in some ferns on the leaf-stalk), and branch either monopodially or dichotomously. There is never one preponderating root continuous in a downward direction with the main stem corresponding to the tap-root in Flowering Plants. They are distinguished from the roots of Flowering Plants by the lateral roots springing, not from the procambium, but from the innermost cortical layer of the mother-root. They are abundantly covered with *root-hairs*, trichomic formations by means of which the nutritive materials are absorbed from the soil. They possess a true *root-cap*. Salvinia (Rhizocarpeæ) and Psilotum (Lycopodiaceæ) are altogether rootless, as also are a few Hymenophyllaceæ (Filices), the function of roots being performed by underground branches of the stem.

The degree of development of the *stem* varies within very wide limits. In the tree-ferns it is of erect habit, and attains great height and considerable thickness. In many herbaceous ferns the internodes of the erect stem are altogether suppressed, while the underground portion forms an elongated rhizome. In the existing Lycopodiaceæ, and in some Selaginellaceæ the very elongated creeping stem is mainly above ground ; in the rootless forms, like Psilotum, branches of the stem bending down into the soil perform the function of roots. In some paludose species belonging to the Rhizocarpeæ the stem is almost entirely suppressed, and in Salvinia the whole plant floats on the surface of the water. The mode of branching is either monopodial or dichotomous ; the leaves do not usually produce buds and branches in their axils, as in Flowering Plants.

In all the larger species the stem displays a distinct differentiation of tissues into the three systems, fundamental, epidermal, and 'vascular' or fascicular. The so-called *vascular bundles* are *closed*, like those of Monocotyledons—that is, they contain no formative cambium ; and they are usually but not always *concentric*, the phloem portion surrounding the xylem portion in the form of a *phloem-sheath*. Each bundle, or a group of bundles, is again very frequently surrounded by a single layer of strongly sclerenchymatous cells belonging to the fundamental tissue, the *vascular bundle-sheath*, where it encloses a single, or *plerome-sheath*, while it surrounds a group of bundles. The prevalent, though not the exclusive form of thickening in the xylem is that of *scalariform tracheides* ; true vessels formed from the coalescence of cells are rare ; in the phloem *sieve-tubes* are of common occurrence. Potonié holds that there is no sharp differentiation between the xylem and phloem portions

of the 'vascular' bundles of Vascular Cryptogams, and prefers for them the terms *hadrome* and *leptome* respectively. According to Van Tieghem, the secondary tissues, like those of Flowering Plants, proceed normally from two concentric generating layers—an external one in the cortex, forming bark outwardly and secondary cortex inwardly, and an inner one in the central 'vascular' cylinder intercalated in the liber and in the xylem of the primary 'vascular' bundles, producing secondary liber outwardly and secondary xylem inwardly. The epiderm is in most cases abundantly provided with trichomic appendages of various kinds.

The size and form of the leaves are extremely various. In Lycopodium, Selaginella, and some other genera, they are very small, unsegmented, and lanceolate, not unlike those of mosses, and form a dense imbricated clothing to the stem ; in Psilotum they are altogether rudimentary : in the Equisetaceæ they are reduced to divisions or teeth of a membranous sheath ; in Isoëtes (Selaginellaceæ), Pilularia (Rhizocarpeæ), and Phylloglossum (Lycopodiaceæ), they are long, narrow, and awl-shaped. In some ferns the barren and fertile leaves differ from one another in appearance, and especially in the degree of division of the lamina. In Salvinia they are of two kinds, one floating on the surface of the water and entire, the other submerged, very finely divided, and performing the function of a root ; in Azolla (Rhizocarpeæ) they are floating and bilobed. In some genera of Filices and their allies the leaves are quite entire ; in the Hymenophyllaceæ they are very delicate, consisting of only a single layer of cells, and in the smaller species closely resemble those of the foliose Hepaticæ ; while in most ferns they are of considerable (in the tree-ferns of gigantic) size, with well-marked petiole, rachis, and lamina, and distinguished by the great extent to which the lamina is divided. In most cases (except the Hymenophyllaceæ) they are abundantly provided with *stomates*. The tissue beneath the epiderm consists of a parenchymatous mesophyll containing abundance of chlorophyll, the portion of which adjacent to the upper epiderm is frequently developed as palisade-parenchyme. This mesophyll is permeated by 'vascular' bundles or veins, which branch off from the cauline bundles, and are distinguished, in the majority of ferns, by their dichotomous mode of branching, in contrast to the reticulate anastomosing in Dicotyledons, and the parallel arrangement in most Monocotyledons. Among Gymnosperms a similar arrangement is presented by Salisburia and Stangeria. The floral metamorphosis of the leaves of Flowering Plants does not occur in Vascular Cryptogams, nor their special agglomeration round the organs of reproduction as in mosses.

The mature *sporange*, theca, or spore-case, is usually a roundish

capsule borne on a stalk, of small size and simple structure. Its morphological value varies greatly, and will be referred to more particularly under the separate families. In the majority of Filices the sporanges are trichomic structures, and are collected into groups or *sori*, which are always located in connection with a 'vascular' bundle on the under side or margin of the leaf. In the Marattiaceæ they spring from a hypodermal mass of tissue. In the Ophioglossaceæ a segment of a leaf is transformed into sporanges. In Selaginella and Lycopodium they arise from the growing point of the stem above the axil of a leaf. In Psilotum they are sunk in the extremity of branchlets of a peculiar form. In the Salviniaceæ they are enclosed in receptacles or *sporocarps*, which are themselves modifications of divisions of the leaf. The mode of formation of the *spores* closely corresponds to that of the pollen-grains in Flowering Plants. The spore-forming tissue can always be traced back to a single cell or a row or layer of cells, the *archespore*, which may be distinguished at a very early period from the remaining cellular tissue by the nature of its contents. From this proceeds the *sporogenous tissue*, which afterwards becomes the mother-cells of the spores by perfectly regular divisions, the details of which differ in the different families. This is surrounded by one or more layers of cells, the *tapetal cells* or *tapete*, and the whole is enclosed in the wall of the sporange, itself composed of one or more layers of cells. In the heterosporous families the distinction between *megaspores* and *microspores* is manifested only at a comparatively late period in their development. In the isosporous families the spores are always strictly unicellular, very commonly elliptical or reniform in shape. The coat always consists of two separable layers—an outer cuticularised *exospore*, often elevated into warts or other prominences ; and an inner *endospore*, composed of cellulose, which bursts through the exospore on germination, producing the *germ-filament*, which develops by cell-division into the prothallium. In the megaspores of the heterosporous families these are further protected on the outside by a third separable, greatly hardened layer, the *epispore*. The mode in which the spores escape from the sporange differs in the different families.

A purely vegetative mode of propagation by means of gemmæ or bulbils borne on the sporophyte occurs especially in Filices and Equisetaceæ ; on the oophyte vegetative propagation is less common.

The classification of Vascular Cryptogams is attended with considerable difficulty. None of the systems as yet produced have much claim to be regarded as natural ; and, until some doubtful points are cleared up connected with fossil forms which may be links between existing families, the primary distinction into *Heterosporous* and *Isosporous*

Vascular Cryptogams is so convenient, that we have decided to adhere to it, without dogmatising as to its permanent retention. In the boundaries of the families, again, there is equal room for diversity of opinion. Whether to retain the Psiloteæ under Lycopodiaceæ, and the Isoëteæ under Selaginellaceæ, and whether to regard the true Filices, the Marattiaceæ, and the Ophioglossaceæ as constituting one, two, or three classes, are points on which there is much to be said in favour of classifications different from that which we have decided to adopt.

LITERATURE.

Thuret—(Zoospores and Antherids) Ann. Sc. Nat., vol. xiv., 1850, p. 214; and vol. xvi., p. 5.
Hofmeister—Germination, Development, and Fructification of the Higher Cryptogams, Ray Soc., 1862.
Schacht—Die Spermatozoiden, Braunschweig, 1864.
Dippel—(Fibrovasc. Bundles) Ber. Deutsch. Naturf. u. Aerzte, 1865.
Nägeli u. Leitgeb—(Root) Nägeli's Beitr. z. wiss. Bot., 1867.
Kny—(Prothallium) Sitzber. Ges. Naturf. Freunde, Berlin, 1868.
Millardet—(Prothallium) 1869.
Russow—Vergleich. Unters., Petersburg, 1872.
Janczewski—(Archegone) Bot. Zeit., 1872, p. 418.
Goebel—(Sporangia) Bot. Zeit., 1880 and 1881.
Sadebeck—Die Gefässkryptogamen, 1880.
Prantl—Morphologie der Gefässkrypt., 1881.
Van Tieghem—Bull. Soc. Bot. France, 1883, p. 169.
Potonié—(Vasc. Bundles) Jahrb. Bot. Gart. Berlin, 1883.
Leitgeb—(Spores) Ber. Deutsch. Bot. Gesell., 1883, p. 246.
Celakovsky—Pringsheim's Jahrb. f. wiss. Bot., xiv., 1884, p. 291.
Bower—(Leaf) Proc. Roy. Soc., xxxvii., 1884, p. 61.
Rabenhorst—Crypt. Flora Deutschland, Vasc. Crypt., by Luerssen, 1884-88.
De Bary—Comparative Anatomy of Phanerogams and Ferns, 1884.
Klein—(Dehiscence of Sporange) Bull. Soc. Bot. France, 1884, p. 292.
Leclerc du Sablon—(Spores) Ann. Sc. Nat., vol. ii., 1885, p. 5.
Campbell—(Antherozoids) Ber. Deutsch. Bot. Gesell., 1887, p. 120.
Baker—Handbook to Fern Allies, 1887.

HETEROSPOROUS VASCULAR CRYPTOGAMS.

Class I.—Rhizocarpeæ.

The Rhizocarpeæ or Hydropterideæ constitute a class composed of only a small number of genera, none of which includes more than a few species. They grow submerged in or floating on water, and derive their name from the circumstance that the non-sexual organs of propagation are produced in the radicular region, or near the base of the leaves.

The spores are of two kinds, one of which is many hundred times larger than the other. The larger spores, or *megaspores*, produced in *megasporanges*, are female ; the smaller spores, or *microspores*, produced

FIG. 5.—*Pilularia globulifera* L., with fructification, natural size. (After Luerssen.)

FIG. 6.—*Marsilea quadrifolia* L., with fructification, natural size, and fructification enlarged. (After Luerssen.)

in *microsporanges*, are male. The megaspore is not completely spherical, but has a distinct apical protuberance, which at the period of maturity is enveloped in a thick firm layer termed the *epispore*, formed by the

hardening of mucilage derived from the disorganisation and deliquescence of a portion of the contents of the sporange. The female prothallium is formed within the apical papilla of the megaspore, and is exposed by the bursting of the enveloping epispore. It never completely frees itself from the megaspore, and is usually altogether destitute of chlorophyll. It bears one or more *archegones*, differing from one another in smaller points of structure in the different genera. The microspores do not give birth to a male prothallium, nor even to antherids, in the sense in which the terms are employed elsewhere in



FIG. 7.—*Salvinia natans* L. *A* and *B* natural size, the latter with two aërial leaves and submerged fertile leaves ; *C*, two sporocarps, slightly magnified and diagrammatic, one containing a few megasporanges, the other a large number of microsporanges ; *D*, section of empty sporocarp, slightly magnified. (After Luerssen.)

Vascular Cryptogams ; the contents divide more or less directly into the parent-cells of the *antherozoids*, which, accompanied by peculiar vesicles attached to them, reach and impregnate the oosphere contained in the central cell of the archegone.

The external form of the sporophyte or non-sexual generation varies widely in the different genera. The growth of both stem and root is always the result of successive divisions from a single *apical cell*. The stem is extremely abbreviated in Salvinia (Schreb.) and Azolla (Lam.) ; procumbent and creeping in Marsilea (L.) and Pilularia (L.). It is traversed

by closed concentric 'vascular' bundles, each surrounded by its bundle-sheath, and the branching is always monopodial. The roots—except in Salvinia, which is rootless—are fibrous, are furnished with a root-cap, and branch monopodially. The leaves also vary greatly in form. In Pilularia they are erect, cylindrical, and setiform; in Marsilea (L.) the lamina consists of several distinct leaflets at the extremity of a more or less elongated petiole. In both these genera the vernation is circinate. In Azolla the leaves are deeply bifid. Salvinia is remarkably heterophyllous. While the majority of the leaves retain an ordinary leaf-like habit, others develop into coriaceous scutiform structures, while others again divide into a number of capillary segments, which perform the function of roots, and at the same time bear the non-sexual propagating organs.



FIG. 8. *Salvinia natans* L. Young plant still attached to the prothallium *pr*, and megaspore *sp*. *b*, scutiform leaf; *I* and *II*, first and second leaves; *L'* and *L''*, later aërial, and *w*, submerged leaf of the first whorl. (After Pringsheim, ×20.)

The fructification is of a more complicated structure than in other classes of Vascular Cryptogams. In Salvinia it springs from the lower teeth of the submerged leaves; in Azolla from the pendent section of the deeply bipartite leaf, or rather of one particular leaf; in Pilularia it stands beside and beneath the leaves; in Marsilea (L.) on the under side of the petiole, or of the base of the leaf itself. The Rhizocarpeæ are always monœcious, the two kinds of sporange being produced on the same individual, and usually in close proximity. The structure and degree of complexity of the fructification differ in the different genera. The sporanges are always associated together in groups. Each of these groups, known as a *sorus* or *sporocarp*, is a closed capsule-like chamber, which is of epidermal or trichomic origin, its wall or *indusium*, often considerably hardened, being an extension of the epiderm. Each sporocarp is regarded by Celakovsky as the homologue of the integumented ovule of Flowering Plants. In the Salviniaceæ the sporocarp is unilocular; in the Marsileaceæ it is plurilocular, and the wall indurated into a hard shell. Each sporocarp may contain sporanges of one kind only or of both kinds; in the former case male and female sporocarps are often associated together in

groups. The megasporanges are often considerably larger than the microsporanges.

In the early stages of their development no difference is exhibited between the megasporanges and microsporanges. In both cases the sporange originates in a papilla placed on the *placenta*, which divides first into an upper and a lower cell, the latter developing, by repeated transverse septation, into the pedicel, the former into the body of the sporange, and dividing ultimately into a large central tetrahedral *archespore*, surrounded by a layer which almost immediately breaks up into two layers of *tapetal cells* or mantle-layers. The archespore further divides into sixteen spore-mother-cells, and each of these into four special spore-cells arranged tetrahedrally. A difference is now manifested according as the sporange is to develop into a mega- or a microsporange. In the latter case each of the sixty-four cells develops into a microspore, while the tapetal cells become disorganised, and changed into the frothy mucilage which subsequently hardens and encloses the spores. In the former case only one of the sixty-four cells develops into a megaspore, growing rapidly at the expense of the others, and ultimately filling up the cavity of the sporange. The remaining sixty-three spore-cells, as well as the tapetal cells, become disorganised, and deliquesce into a frothy mucilage which envelops the ripe megaspore, ultimately hardening into the *epispore*, which splits to allow the emergence of the prothallium. In Azolla the mucilage of the microsporanges forms the peculiar *massulæ* which will be described later.

A more detailed description requires the division of the Rhizocarpeæ into the two orders *Salviniaceæ* (Salvinia and Azolla) and *Marsileaceæ* (Marsilea and Pilularia), which are, perhaps, not in reality very nearly related to one another.

Order 1.—Salviniaceæ.

The female prothallium of Salvinia is formed within the apical papilla of the megaspore. The protoplasm in this papilla appears to separate from that of the rest of the spore, and then breaks up by free-cell-formation into several portions, which remain for a time unclothed with cellulose; subsequently they secrete cell-walls, and form a tissue, which breaks through the cell-wall of the papilla, and forces its way through the epispore, which splits into a three-lobed body. The prothallium, when it first emerges from the epispore, has a somewhat triangular form, with an elevated ridge along its median line, and two wing-like appendages, subsequently forced apart by the growth of the embryo, which hangs down on each side of the spore. It contains a considerable amount of

chlorophyll, but never loses its connection with the megaspore, even after the commencement of the germination of the sporophyte. The first archegone makes its appearance on the elevated dorsal ridge of the prothallium, and subsequently two others are formed, one on each side of the first. If one of these is fertilised no more archegones are produced, and the prothallium ceases growing. But if no impregnation has taken place the prothallium continues to grow, and produces from one to three additional rows, each consisting of from three to seven archegones. In Azolla the prothallium has the form of a slightly convex disc, consisting, in its central part, of several layers of cells, at the margin of only one. A single archegone is first formed, near the centre of the prothallium. If this



FIG. 9.—*Salvinia natans*. Longitudinal section through megaspore and prothallium. *a*, wall of sporange; *b*, epispore, formed of hardened mucilage; *c*, coat of spore; *d*, diaphragm separating prothallium from spore-cavity; *pr*, prothallium; *m*, neck of archegone; *I*, *II*, first two leaves of embryo; *s*, scutiform leaf or cotyledon. (After Pringsheim, × 70.)



FIG. 10.—Archegone of *Salvinia natans* in three stages. *a*, *b*, *c*, divisions in the neck-cells; *d*, neck-canal-cell; *e*, oosphere; *h*, neck-cells. (After Pringsheim, × 150.)

is fertilised, no others are formed; if not, it is followed by a few others.

The *archegone* of Salvinia is a nearly globular cavity, its venter being completely buried in the tissue of the prothallium. The *central cell* is at first somewhat elliptical, its axis lying obliquely to the surface of the prothallium. Its apex is at first covered by four cells belonging to the epiderm, arranged in the form of a cross. Each of these four *neck-cells* divides by transverse septa into a row of three cells, the four rows thus forming a short neck. The large central cell now elongates upwards and forces its way between the lowermost four cells of the neck, and its

conical point becomes cut off by a septum, forming the *neck-canal-cell*. Below this a second very small portion of the central cell is again cut off to form the *ventral canal-cell*, so that the canal now consists of two cells. These two cells become transformed into mucilage, which escapes by forcing apart the four apical or *stigmatic cells*, leaving an open canal In the meantime the protoplasm of the large basal portion of the central cell has become transformed by contraction into the *oosphere*. The archegone is now ready for impregnation, the antherozoids reaching the oosphere through a funnel-shaped depression in the epispore and the open canal. After fertilisation the canal again closes up by the expansion of the stigmatic cells. The archegone of Azolla resembles that of Salvinia in all essential points.

The male prothallium of Salvinia is reduced to a mere rudiment. The microspores lie imbedded in a mass of hardened, granular, frothy mucilage, formed by the disorganisation of the tapetal cells. They do not escape from this mucilage, but the endospore of each develops into a tubular filament which pierces through the mucilage and the wall of the sporange. The extremity of this filament which projects outside the sporange is curved, and becomes cut off by a septum. The lower and larger of the two cells thus formed is regarded as a rudimentary prothallium; the terminal cell, which again divides into two, as a rudimentary antherid. The protoplasm of each of the two antheridial cells divides into four, and each of these eight masses of protoplasm escapes as an *antherozoid*. Each antherozoid is a corkscrew-like coil of protoplasm, bearing vibratile cilia of great length at its broader extremity. To the same extremity is attached a vesicle, composed of a portion of the protoplasm of the antheridial cells which was not used up in the formation of the antherozoids, and which does not leave the antherozoid during the period of its 'swarming.'



FIG. 11.—*Salvinia natans*. *A*, microsporange, with microspore-tubes *st*. (× 100.) *B*, microspore-tube (× 200) with closed, *C* with empty antherid. *D*, antherozoids (× 600). (After Pringsheim.)

The development of the multicellular *embryo* from the fertilised oosperm has been very carefully followed out in Salvinia. The first segmentation is by a nearly vertical wall (at right angles to the surface of the prothallium) into two somewhat unequal portions, each of which

again divides by a septum nearly at right angles to the first one. Further divisions then take place. Out of the posterior of the first two segments (the one immediately beneath the mouth of the archegone) is formed the *foot* of the young plant, by which it is attached to and derives its nutriment from the prothallium. From the anterior of these two segments is derived a peculiar foliar structure, differing from all the subsequent leaves, the *cotyledon* or *scutiform leaf*, by the growth of which the terminal bud of the stem becomes directed downwards. No root whatever is produced. Azolla is stated to have a second cotyledon. Both stem and root (in Azolla) are developed from a single apical cell, which is rounded in front and pointed below. The primary meristem-layers are differentiated, as in Flowering Plants, into plerome, periblem, and dermatogen.

The mature sporophyte differs considerably in appearance in the two genera, but always floats on the surface of the water. The very short stem is erect or horizontal, and the branching of both stem and root (in Azolla) is monopodial. The root, stem, and leaf-stalk are each traversed by a single concentric 'vascular' bundle of simple structure, containing spiral and annular tracheides. The leaves of Azolla are very crowded, and are placed in two rows on the dorsal side of the stem ; but in some species they have the appearance of standing in four rows. They are of delicate membranous texture, and are always deeply bifid, one lobe being submerged and the other floating. The floating lobe of each leaf has a remarkable cavity, covered by a double epidermal layer, with the exception of a narrow orifice which opens into the cavity. This cavity is formed during the growth of the leaf by a more rapid growth of the epiderm than of the subjacent tissue, and is itself clothed with an epidermal layer. The cavities are frequently occupied by well-developed colonies of Nostoc-filaments. Salvinia is remarkably heterophyllous (see figs. 7, 8). The first leaf of the young plant is the *scutiform* or *peltate leaf* already mentioned, which is produced near the base of the stem. It is coriaceous in texture and sagittate in form. Next are produced, also singly, two ovate *aërial leaves*. All the subsequent leaves are arranged in whorls of three, two of which are aërial, with flat, ovate, or orbicular lamina, floating on the surface of the water ; while the third or *submerged leaf* at once branches into long slender filiform segments, which hang down into the water and perform the function of roots. The leaves of adjacent whorls are placed alternately, so that the mature plant has two rows of ventral submerged, and four rows of dorsal aërial leaves. Each leaf has a single definite apical cell in Salvinia, but not in Azolla. The leaves of both genera are furnished with stomates, which, according to Strasburger, differ considerably, both in structure and

appearance, from those of Flowering Plants. Those of Salvinia are remarkably small, and are inserted about halfway up the epidermal cells, which are eight or nine times their height. Air-pores occur also in the submerged leaves. The very simple roots of Azolla are of endogenous origin. The root-cap originates from a single cell; in A. Caroliniana (Willd.) the cap is eventually thrown off, leaving the root-tip naked.

The sporanges are enclosed in unilocular *sporocarps*, formed two together or in larger numbers; in Salvinia on the youngest teeth of the submerged leaves, in Azolla on the pendent submerged lobe of the deeply bifid leaves, and only on the lowermost leaf of each shoot. The leaf-segment which is destined to become fertile first of all develops into a *columel* or *placenta*, to which the sporanges are attached. An annular wall, the rudiment of the *indusium*, then becomes elevated from the base of the columel, eventually overtops its apex, closes up, and thus forms the wall of the sporocarp. The sporocarp of Salviniaceæ is therefore a metamorphosed portion of a leaf, and corresponds to a sorus in the Hymenophyllaceæ (Filices), with the difference that in the latter the envelope remains open in the form of a cup, while in the former it closes completely over the sorus, as in Cyathea (Filices). The indusium is much more strongly developed than that of ferns, and completely envelops the sorus; it consists of two layers of cells, the walls of which are, in Azolla, strongly lignified in the upper part. Each sporocarp contains one kind of sporange only; but both kinds always occur on the same individual, and may even spring from the same metamorphosed leaf. In Salvinia the megasporanges are considerably larger than the microsporanges, and the number of the latter in a sporocarp is greater (see fig. 7). In Azolla the number of microsporanges in a sporocarp is about forty, while the female sporocarps contain only a single megasporange, and consequently only a single megaspore, enveloped first in the wall of the sporange, and then in the greatly hardened indusium. The microsporanges are nearly globular capsules, with long slender pedicels, the wall consisting, when mature, of a single layer of cells. The megasporanges are pear-shaped, with much shorter and



FIG. 12.—Fertile shoot of *Azolla filiculoides* Lam., with two female sporocarps, *a* (×27). (After Strasburger.)

stouter pedicels, and arise from the apex of the columel. In the formation of the pedicels of the megasporanges longitudinal cell-division takes place, as well as transverse. The mode of formation of the spores within each kind of sporange has already been described in general terms, after the preliminary separation of a single external layer of cells which develops into the wall of the sporange. The sixty-four microspores appear to be disposed without any arrangement in the cavity of the microsporange. A large nucleus lies at the end of the megaspore which is nearest the apex of the sporange. Before fertilisation both kinds of sporange become detached from their pedicels, and are carried



FIG. 13.—Massula of *Azolla Caroliniana* Willd. (×240). (After Strasburger.)

up to the surface of the water in the spring by the surrounding masses of Algæ. The epispore then splits above the apex of the megaspore into three lobes, between which the emerald-green prothallium forces itself, and impregnation is effected. In Azolla the epispore assumes a still more striking form. In the microsporanges it has the appearance of a large-celled tissue, and breaks up into two or more spherical masses called *massulæ*, each of which envelops a number of microspores, and has a distinct coat. In some species, but not in all, the surface of these masses is covered with hair-like appendages, barbed at the apex, the *glochids*, by means of which, after emerging from the sporange, and when floating on the surface of the water, they attach themselves to the

floating megaspores. The roundish megaspore, which does not nearly fill up the sporange, is completely covered by a very thick warty layer of hardened frothy mucilage, its epispore, which projects far above the apex, and separates, in its upper part, into either three or nine large pear-shaped masses of the same substance, terminating in a tuft of fine threads. These bodies constitute a *floating apparatus* for the megaspore, the fine threads floating on the surface of the water, and suspending beneath them the *float-corpuscles*, either three in number or more numerous, containing abundance of air-cavities, and the megaspores, to which the microspores are attached by their glochidiate processes.

The two genera of Salviniaceæ, Salvinia and Azolla, each include but a small number of species, all annual plants, widely distributed over the globe, especially in its warmer regions. Those of Azolla form green or red floating patches of considerable size, with the habit of a Jungermannia. No economical application is known of either genus.

Order 2.—Marsileaceæ.

The female prothallium attains here a much smaller degree of development than in the Salviniaceæ. It arises within the apical papilla of the megaspore, the protoplasm of which breaks up into several cells, which remain for a time unclothed with cellulose, and only subsequently constitute a tissue containing a small quantity of chlorophyll. Even after this the prothallium still remains for some time completely enclosed within the apical papilla of the megaspore, being covered by the epidermal layers of the apex of the spore itself, and shut off from the spore-cavity within and below by a *diaphragm* which is attached to the internal coat of the spore. By the further growth of the prothallium the epidermal layers of the apical papilla are broken through, and the dorsal ridge of the prothallium projects into the *funnel* formed by the absence at this spot of the thick outer layers of the epispore. The diaphragm subsequently arches convexly, and the prothallium is pushed further outwards, but still lies as a hemispherical mass in the funnel-shaped opening.

In those species which have hitherto been examined each prothallium produces only a single archegone. Even before the prothallium breaks through the megaspore, the large *central cell* may be recognised in it, covered only by four cells arranged in a cross, which form at the same time the apex of the prothallium. From these are developed the more or less projecting neck and the stigmatic cells of the archegone. As in Salviniaceæ, a neck-canal-cell is separated from the central cell, which pushes up between the neck-cells, as well as a smaller ventral canal-cell;

the lower and larger portion of the protoplasmic cell-contents contracting into an *oosphere*. If the archegone remains unfertilised the prothallium continues to grow into a comparatively large chlorophyllous structure with rhizoids.

The male prothallium and antherids are reduced to a still more



FIG. 14.—*Marsilea salvatrix* L. Longitudinal section through megaspore, prothallium, and embryo. *am*, starch grains; *i*, inner coat; *ex*, epispore; *c*, space beneath diaphragm; *pt*, prothallium; *wh*, its rhizoids; *a*, archegone; *f*, foot; *w*, root of embryo; *b*, cotyledon; *sl*, mucilaginous envelope of megaspore. (After Goebel, × 60.)



FIG. 15.—*Marsilea salvatrix*. *A*, prothallium *pt*, projecting from ruptured coat *r* of megaspore; *sl*, layers of mucilage forming funnel, with antherozoids. *B*, vertical section of prothallium *pt*; *o*, oosphere; *a*, stigmatic cells. (After Goebel, greatly magnified.)

rudimentary condition than in the Salviniaceæ. The contents of the microspore divide into three cells, one of which (the prothallium) remains sterile, the other two constituting the antherid. The contents of each of these two cells again divide into sixteen antherozoid-mother-cells. From the nucleus of each of these is formed an antherozoid;

these bodies are therefore developed entirely within the microspore, while the microspores themselves are set free completely from the sporange. As in the Salviniaceæ, the whole of the contents of the mother-cell is not used up in the formation of the antherozoid ; a portion remains behind in the form of a roundish turbid lump consisting of protoplasm and starch-grains, which gradually becomes clearer, and attaches itself, in the form of a vesicle, to the antherozoid, which in Pilularia becomes soon detached, but in Marsilea remains attached to the antherozoid during the greater portion of the period of 'swarming.' When the antherozoids are mature the exospore of the microspore bursts at its apex, and the endospore swells up into a hyaline bladder,



FIG. 16.—*Pilularia globulifera* L. Longitudinal section of megaspore. *a*, coat of spore ; *b*, *c*, *d*, the three layers of the epispore. (After Luerssen, magnified.)



FIG. 17.—*Marsilea salvatrix*. Microspore discharging antherozoids. *ex*, exospore ; *dl*, endospore ; *zz*, antherozoids; *yy*, their vesicles with starch-grains. (After Goebel, ×350.)

which finally bursts to allow of the escape of the antherozoids with their vesicles. In Pilularia the antherozoid consists of only four or five coils with a few vibratile cilia ; in Marsilea it is of considerable length, the shape of a corkscrew, and consists of twelve or thirteen coils, the vibratile cilia being also of great length. The antherozoids collect, in large numbers, in the funnel-shaped depression of the epispore of the megaspore above the prothallium (see fig. 15), and force themselves, through the neck of the archegone, to its central cell.

In its early stages the development of the *oosperm* or impregnated oosphere corresponds to that of Salviniaceæ. After becoming invested with a cell-wall of cellulose, the first segmentations give rise to the

parent-cells of the first root, of the young stem, of the first leaf or cotyledon, and of the foot by which the young embryo is attached to the prothallium. In Marsilea a second cotyledon is formed from the fourth octant of the lower half of the embryo. The layer of tissue surrounding the central cell becomes double after impregnation ; a few grains of chlorophyll are formed in it, and its outer cells develop into long rhizoids, which are especially luxuriant if no fertilisation has taken place.

The sporophyte of the Marsileaceæ differs very widely in external form in the two genera ; but its internal structure agrees in its essential features with that of the Salviniaceæ. The stem, root, and leaves all originate from a single apical cell, which always divides into three rows. The stem is procumbent on damp soil or at the bottom of stagnant water, and is traversed by a single central 'vascular' cylinder filled with fundamental tissue, each bundle consisting of a central xylem with spiral or scalariform tracheides, surrounded by a phloem with large sieve-tubes and sieve-plates, and the whole enclosed in a brown sclerenchymatous bundle-sheath, composed of a single layer of cells with wavy lateral walls. A single 'vascular' bundle traverses each root and leaf-stalk ; in the lamina of the leaf of Marsilea this branches into a dichotomous venation. The fundamental tissue abounds, in both genera, in large schizogenous intercellular cavities, as is usually the case with water-plants. Those of Marsilea form a complete intercommunicating system. In those of Pilularia are remarkable spiral hairs. The leaves develop basifugally, as in Salviniaceæ ; they are formed in two alternating rows on the dorsal side of the stem ; but, as in Salviniaceæ, it is not every segment of the stem that produces a leaf. In this respect the Rhizocarpeæ agree with Filices, and differ from Equisetaceæ and Muscineæ. The leaves are circinate in vernation, resembling in this respect true ferns only. Tannin-sacs occur in the petiole. In Marsilea (see fig. 6) all the leaves except the first, which is filiform and destitute of a lamina, have a very long slender petiole and a four-lobed lamina ; they are larger when growing in water than on dry land. M. quadrifolia (L.) has stomates on both surfaces of the aërial, on the upper surface only of the floating leaves. The stomates are depressed in the tissue of the leaf by the growth of the adjoining epidermal cells over the guard-cells. The mesophyll of the aërial leaves is characterised by the presence of rows of sclerenchymatous cells. In Pilularia the petiole is elongated and quill-like, and entirely destitute of a lamina (fig. 5).

The *sporocarp* or conceptacle of the Marsileaceæ is an even more complicated structure than that of the Salviniaceæ. In Pilularia it is a roundish, shortly-stalked capsule springing from the axil of a leaf-stalk

on the procumbent stem. According to Juranyi, each sporocarp is the result of the coalescence of two segments of bifid leaves. The wall of the sporocarp is very thick and hard, and consists of several layers of cells forming a sclerenchymatous tissue. It is divided by vertical walls into compartments, varying from two to four in the different species. Each compartment has, at least in its young state, an opening at the apex, and is therefore not of endogenous origin, but rather a depression in the surface. In each compartment there is, on the side which forms its outside wall, a cushion-like *placenta*, formed from superficial cells, and resting on a 'vascular' bundle. To this placenta are attached a number of stalked sporanges of both kinds, constituting a *sorus* ; the megasporanges are chiefly below, and the microsporanges above. The remainder of each compartment is occupied by a delicate thin-walled parenchyme. When mature the sporocarp splits from the apex downwards into as many valves as it has compartments ; and each sporange dehisces by the expansion of the gelatinous mass resulting from the dissolution of the tapetal cells.



FIG. 18. -Transverse section of sporocarp of *Pilularia globulifera*. *mi*, microsporanges ; *ma*, megasporanges ; *e*, wall of sporocarp ; *g*, parenchyme. (After Goebel, magnified.)

The sporocarps of Marsilea are capsules with somewhat the shape of a bean, a longer or shorter pedicel, and a very hard sclerenchymatous wall. They spring, usually in clusters, from the petiole of an ordinary leaf. The pedicel runs along the dorsal edge of the sporocarp, and gives off lateral veins right and left which branch dichotomously and run to the ventral edge. The ripe sporocarp has a bilaterally symmetrical structure, and is divided by transverse walls into two rows of compartments, each of which has, when young, a narrow opening on the ventral side. Each compartment contains a single sorus, consisting of a few megasporanges in the centre, with a larger number of microsporanges on each side. As in Pilularia, a large portion of the cavity of each compartment is occupied by a succulent parenchyme.

The development of the sporanges commences, in both genera, with the swelling up of some of the epidermal cells of the part which ultimately becomes the placenta. These cells divide several times

obliquely—not horizontally, as in Salviniaceæ—into three rows of segments, until ultimately a convex septum cuts off a triangular apical cell, which at length becomes the tetrahedral *archespore*. From this is separated the mantle-layer of *tapetal cells* ; further divisions take place both in these and in the rows of cells of which the wall of the sporange is composed, and the archespore divides by successive bipartitions into sixteen spore-mother-cells, each of which produces four spores in the ordinary way. The pedicel of the sporange consists at first of three rows of cells, the number being subsequently increased by further longitudinal divisions. The tapetal cells become gradually disorganised, and form a granular mucilage, filling up the interstices between the mother-cells of the spores, which is subsequently employed in the production of the *epispore* or gelatinous envelope of the spore. The differentiation of the two kinds of spore now commences. In the microsporanges all the sixty-four microspores reach maturity, each special mother-cell or rudimentary spore becoming invested, while still within the mother-cell, with its permanent cell-wall, while the walls of the sixteen mother-cells disappear. In the megasporanges, on the other hand, one of the four special mother-cells in each of the sixteen tetrads displays at first a greater vigour of growth than the other three. Of these sixteen sister-cells fifteen gradually become abortive, only one reaching maturity and developing into a perfect megaspore. During their development and disappearance all the rudimentary spores are furnished with spiny protuberances, by which they are attached to one another. As the megaspore increases in size its coat becomes hard and brown ; and it is ultimately invested by a gelatinous *epispore* consisting of three distinct layers (fig. 16). The innermost of these is a mucilaginous coat, which is often folded, and ultimately forms a papilla above the apex of the mature spore. Outside this is a thicker layer of a soft prismatic substance, and external again to this a third still thicker but less clearly organised envelope. The two outer layers are wanting at the apex of the spore, where there is a funnel-shaped depression exposing the papilla belonging to the innermost layer of the epispore. Down this *funnel* the antherozoids pass to impregnate the oosphere within the archegone produced on the prothallium within the apical papilla of the megaspore.

The processes by which both kinds of spore escape from the very hard shell of the sporocarp are very remarkable. In Pilularia globulifera (L.) the ripe sporocarp lies above or beneath the surface of the damp soil. It splits from its apex downwards into four valves, and exudes a tough hyaline mucilage derived from the parenchymatous tissue within each compartment. This mucilage accumulates on the ground ; and into it both kinds of spore escape after the rupture of the

sporanges. Fertilisation takes place within the drop of mucilage, which then gradually disappears, and the impregnated megaspore lies on the damp ground, to which it becomes attached by the rhizoids put out from the prothallium until the first root of the embryo penetrates into the soil. In Marsilea the processes are somewhat similar. The excessively hard, almost stony, shell of the sporocarp gives way slightly at its ventral edge as it lies in water, and the water penetrates into the interior. This causes the succulent parenchymatous tissue in each compartment to swell up, and splits the shell along the whole of the ventral edge into two valves. Between these valves the contents are gradually forced out; the compartments still remain closed, each enclosing a sorus, and are attached in two rows to the cartilaginous cushion or *sorophore* which ran along the ventral edge of the sporocarp, and which now becomes detached at one end, and exposed in the form of a string many times longer than the sporocarp itself ; by the absorption of water it has increased enormously in size, to something like 200 times its original dimensions, and the sori are thus placed at a considerable distance from one another. Ultimately the walls of the sori or original compartments of the sporocarp disappear ; the walls of the sporanges burst, both kinds of spore escape, and fertilisation is effected on the damp soil.



FIG. 19.—Sporocarp of *Marsilea salvatrix*. *A*, transverse section. *B*, swollen and bursting, showing sorophore (×2½). *C*, sporocarp (natural size), with sorophore fully extended, and sori attached. *D*, section of sorus (×6); *ids*, indusium; *mik*, microsporanges; *mak*, megasporanges. (After Hanstein.)

All the species of both genera of Marsileaceæ are marsh or aquatic perennial plants, mostly natives of the warmer temperate and tropical countries. The number of species of Pilularia is small, of Marsilea about forty. The leaves of Marsilea, when growing in the air, display a sensitiveness to light, folding up in the evening and expanding in the morning, similar to those of many Leguminosæ and other Flowering

Plants. The sporocarps of M. Drummondii (Br.), and probably of some other species, are eaten by the natives of Australia under the name of nardoo.

LITERATURE.

Bischoff—Die Rhizocarpeen und Lycopodiaceen, Nuremberg, 1828.
Mettenius—Beitr. zur Kenntniss der Rhizocarpeen, 1846; Linnæa, 1847, p. 260; and Beitr. zur Botanik, Heft 1, 1853; Plantæ Tinneanæ.
Meyen—Nov. Art. Acad. Cæsar-Leopold., vol. xviii., pt. 1, p. 253.
Hofmeister—Ueb. Keimung der Salvinia, Abhandl. Sächs. Gesell. Wiss. 1857, p. 665.
Pringsheim—(Salvinia) Jahrb. f. wiss. Bot., vol. iii., 1863, p. 484.
Hanstein—Ueb. eine neuholländische Marsilia, Monber. Berl. Akad., 1862, p. 183; Befruchtung u. Entwickelung der Marsilia, Jahrb. f. wiss. Bot., vol. iv., 1865, p. 107; Pilulariæ generatio cum Marsilia comparata, Bonn, 1866.
Braun—(Marsilia and Pilularia) Monber. Akad. Wiss. Berlin, 1870, p. 653, and 1872, p. 668.
Russow—Vergleich. Unters., Petersburg, 1872; and Hist. u. Entwick. d. Sporenfrucht v. Marsilia, Dorpat, 1877.
Strasburger—Ueber Azolla, Jena, 1873.
Juranyi—Ueb. d. Entwick. d. Sporangien u. Sporen v. Salvinia, Berlin, 1873; and (Pilularia) Sitzber. Ungar. Akad. Wiss., 1879 (see Bot. Centralbl., vol. i., 1881, p. 207).
Berggren—(Azolla) Rev. Sc. Nat., 1881, p. 21.
Heinricher—(Spores of Salvinia) Sitzber. Akad. Wiss. Wien, vol. lxxxv., 1882, p. 494.
Goebel—(Pilularia) Bot. Zeit., 1882, p. 771.

Class II.—Selaginellaceæ.

This class is composed of two genera only, Selaginella (Spring) and Isoëtes (L.), resembling one another in the general facts of their life-history, but differing widely in external appearance, and each constituting a monotypic order. We have, again, as in Rhizocarpeæ, two kinds of spore; the megasporanges and microsporanges are of very similar appearance, and are produced in connection with the leaves. The female prothallium, produced within the megaspore, is a more completely endogenous structure than in any other class of Cryptogams, and is altogether destitute of chlorophyll. From the occurrence in both genera of a foliar structure known as the *ligule*, the term 'Ligulatæ' is sometimes given to the class; but the character is unsatisfactory, and it will be best to treat the two orders *Selaginelleæ* and *Isoëteæ* separately.

Order 1.—Selaginelleæ.

In the genus Selaginella, the sole representative of the order, the prothallium appears to be already completely formed by the time the megaspore is mature, but occupies only the apical portion of the cavity of the spore; the basal portion is still filled by an undifferentiated mucilaginous protoplasm, which subsequently develops into a cellular tissue, the *secondary prothallium*, or, as it is termed by some writers, the *endosperm*. In some species, at least, this structure is separated by a diaphragm from the true prothallium. The prothallium always produces a number of archegones, sometimes as many as thirty, which arise in centrifugal succession on the exposed portion of the prothallium, the one formed first being at the apex. The archegone originates by division of a superficial cell in a direction parallel to the surface; the outer of these two then divides into four cells, and each of these again breaks up by tangential division into two. These form the *neck* of the archegone, which therefore consists of four rows, each composed of two cells. The lower of the two original cells becomes the *venter* of the archegone, and puts out a slender prolongation between the neck-cells, which is separated as the *neck-canal-cell*. Another very small portion is subsequently separated



Fig. 20.—*A—F*, stages in the division of the microspore of *Selaginella caulescens* Spr. *v*, sterile cell; *G*, antherozoid (× 1400); *H*, vertical section of megaspore of *S. Martensii* Spr.; *p*, prothallium with three archegones; *end*, endosperm; *e*, exospore (× 165). (After Pfeffer.)

as the *ventral canal-cell*, and the protoplasm of the larger and lower portion rounds itself off into the *oosphere*. The two canal-cells then deliquesce into mucilage, leaving an open passage for the entrance of the antherozoids.

The microspores, spherical orange or bright red bodies, remain in a dormant condition through the winter, and undergo further development in the spring. The contents then divide, first of all by a transverse wall of cellulose near one end into two cells of very unequal size. The smaller one of these does not divide further, and remains sterile ; it is regarded as the last degraded vestige of a male prothallium. The contents of the larger of the two cells, which may be regarded as an *antherid*, then break up into from four to eight primordial cells, and each of these divides again into four mother-cells of antherozoids ; but it is uncertain whether all the cells are fertile. The *antherozoids* are coiled up into a helix, and are furnished at the anterior end with two long fine cilia. The antherozoid is in all cases derived from the nucleus of the mother-cell ; a central vacuole, invested with a delicate membrane, often remains attached to its posterior end during 'swarming.' The 'swarming' condition continues for about half to three-quarters of an hour.



FIG. 21.—Formation of embryo and suspensor in *S. Martensii*, showing order of formation of dividing walls. (After Pfeffer.)

In the formation of the embryo the first division of the oosperm differs from that in Rhizocarpeæ and in Filices in taking place at right angles to the axis of the archegone. It is thus divided into two superposed halves, from the lower of which is developed the *embryo* itself ; from the upper half a structure almost peculiar to this order, consisting

of a small number of large cells, known as the *suspensor*, or pro-embryo, from its apparent homology with the structure which goes by this name in Flowering Plants. By the elongation of the suspensor and the compression and absorption of the adjacent cells, the lower portion of the oosperm is forced into the endosperm, from which it appears to derive nutriment, dividing, both previously and subsequently, into a small-celled tissue. This tissue undergoes a large amount of differentiation before the embryo emerges from the megaspore, the rudiments of all the principal parts of the sporophyte making their appearance at this early stage. The mother-cell of the embryo divides into two by an oblique wall. From one of the two cells thus formed originate the *stem* and one of the *cotyledons*, from the other the *foot* and the other cotyledon. The rudimentary stem has a two-edged *apical cell*, from which segments are cut off alternately right and left. An inner mass of tissue soon becomes differentiated as the procambium of the axial 'vascular' bundle, the peripheral tissue as dermatogen and periblem. The stem-bud, or *plumule*, with its first leaves (subsequent to the cotyledons), finally grows erect from the apex of the megaspore as the embryo increases in length. The formation of the first *root* begins later between the foot and the suspensor; its apical cell is formed from an inner cell of the older segment; the first layer of its root-cap originates from the splitting into two layers of the overlying dermatogen; the later layers of the root-cap are formed from the apical cell of the root.

The sporophyte differs greatly in appearance in the two orders. In Selaginella the stem is always slender, erect or procumbent, with distinct internodes, and lengthening rapidly by monopodial branching, which very often has a dichotomous appearance from the vigorous growth of the lateral branches. These lateral branches, with their ramifications, frequently develop in a single plane, giving the system the appearance of a compoundly pinnate leaf; all the branches and leaves displaying a distinctly dorsiventral character. The stem has an epiderm composed of elongated prosenchymatous cells, without stomates, but containing a considerable amount of chlorophyll in remarkably large grains. The fundamental tissue consists of elongated thin-walled cells with oblique septa, fitting closely together without intercellular spaces, and endowed with the power of long-continued growth both in length and diameter. In the absence of small intercellular spaces the stem of Selaginella resembles that of mosses; but, on the other hand, each 'vascular' bundle is surrounded by a large air-cavity, traversed by *trabecules*, rows of cells connecting the bundle with the surrounding fundamental tissue. The entire cortex has a tendency to become dark brown with age from partial sclerosis. When

the cells of which these trabecules are composed are round, they form a loose spongy parenchyme surrounding the bundle, and sharply differentiated from the firm compact fundamental tissue. The stem has one or more cauline 'vascular' bundles, which may be traced in the



FIG. 22.—*Selaginella inæqualifolia* Spr. *A*, branch (natural size); *B*, microsporange; *C*, megasporange (greatly magnified).

procambial condition to the apex of the stem close beneath the apical cell and above the youngest leaves ; the separate bundles which descend from the leaves (leaf-trace-bundles) only unite with the cauline ones at a later period. The bundles are usually ribbon-shaped, and are concen-

tric and closed. In the centre is the xylem, consisting chiefly of scalariform and reticulate tracheides; this is completely surrounded by the



FIG. 23.—Transverse section through leaf of *S. inæqualifolia*. *ch*, chlorophyll-bodies; *co*, upper epiderm; *eu*, lower epiderm; *sp*, stomates; *l*, air-cavity surrounding vascular bundle and traversed by trabecules. (After Goebel, magnified.)

FIG. 24.—Transverse section of stem of *S. denticulata* Lk. *b*, air-cavity surrounding 'vascular' bundle. (After Goebel, magnified.)

thin-walled phloem. The primary elements of the xylem, very narrow spiral tracheides, are formed at the angles of the bundle, and from them

the development and lignification of the tracheides advance centripetally. The external layer of phloem is itself surrounded by two or three parenchymatous layers, constituting a *bundle-sheath*, belonging to the fundamental tissue, but within the large air-cavity. The mode of apical growth varies in the different species. In some the apex of the stem is occupied, as in Isoëtes, by a group of equivalent merismatic cells ; while in others there are two co-ordinate apical cells side by side, or a single one, which may be two-sided or three-sided.



Fig. 25.—*S. inæqualifolia* ; transverse section of stem (×150). (After Sachs.)

The leaves are simple and unbranched, and are traversed by a single 'vascular' bundle. They are always of small size, resembling those of Lycopodium, awl-shaped and acuminate, or ending in a delicate awn, and usually with a cordate base. The greater number of species are heterophyllous, the sterile leaves having two different forms ; those on the ventral or shaded side of the obliquely ascending stem are larger than those on the dorsal side exposed to the light. They are always in four rows, one dorsal and one ventral leaf forming a pair. On the upper side of the leaf near its base is the peculiar structure known as the *ligule*, from the presence of which the class has sometimes been called 'Ligulatæ.' The fertile leaves are uniform in size, and differ somewhat in shape from the sterile, forming a compact square terminal spike. The sporange springs from the upper surface below the ligule. In some species the epiderm is alike on the two sides of the leaf ; in others it differs. The epidermal cells contain chlorophyll, as is the case in ferns, and frequently have beautifully serpentine lateral walls ; in some species they are so greatly thickened that the cell-cavity disappears alto-

gether. The chlorophyll, both in the epidermal cells and in the mesophyll, is collected into large lumps, in which are grains of starch. Stomates occur in the under, rarely also in the upper surface. The mesophyll consists of a loose spongy parenchyme ; when the leaves are very small it is developed only as a single layer surrounding the central 'vascular' bundle, and is altogether suppressed near the margins, where the upper and lower epiderm are in actual contact.

True roots occur in all known species belonging to the order. In some species of Selaginella a structure known as the *rhizophore* intervenes between the stem and the root. The rhizophores resemble roots in general appearance, but are destitute of a root-cap. They may spring either from the dorsal side of the stem only, near the base of a branch, bend round and then grow downwards, or two may spring from each fork, one on the dorsal, the other on the ventral side, the former of which usually remains undeveloped in the form of a small protuberance, while the latter grows to the normal size. Their origin is very near the growing point, and they appear to be formed in the same way as the branches. Unlike the roots, they are exogenous structures. After apical growth has ceased, the end of the rhizophore, which is still very short, swells up into a spherical form ; its cell-walls become thicker, and the first rudiments of the true root originate in the interior of the swelling, but do not break through the surface until the rhizophore has increased considerably in length by intercalary growth, and its swollen end has penetrated the soil, where its apical cells deliquesce into mucilage, through which the true roots reach the ground. In some species the rhizophores are frequently transformed into leafy shoots, which at first manifest some deviations from the normal structure of aërial shoots, but afterwards present the ordinary structure, and may even bear sporanges. The rhizophore is not, however, universal in Selaginella. In many species the roots spring directly from the lowest fork of the stem, and branch monopodially before they reach the ground. They originate, like those borne by the rhizophores, near the growing point. All the roots branch copiously, the planes of the successive branchings crossing one another at right angles. They have a single apical cell, but this soon ceases to give off segments, and the subsequent increase in length is chiefly due to intercalary growth.

Both kinds of sporange are shortly stalked nearly spherical capsules (fig. 22), closely resembling those of Lycopodium in appearance and structure, except in their being heterosporous. They are collected into dense spikes at the extremity of somewhat metamorphosed leafy shoots. The lower portion of each spike in some species consists of megasporanges, the upper portion of microsporanges, and the former may be

reduced to only one. Each fertile leaf or *sporophyll* subtends only a single sporange, which is borne on the stem above the leaf-axil. The sporanges are of considerable size in proportion to that of the leaf, and are formed from a group of superficial cells at the growing point of the stem. They make their first appearance as flat, afterwards spherical or club-shaped, swellings, completely covered by the epiderm, which subsequently forms, by tangential divisions, the wall of the sporange, composed of three layers. By subsequent growth the sporange comes to be placed in the axil or even on the base of the leaf. The 'vascular' bundle of the leaf passes beneath the sporange without sending a branch into it. As in the other heterosporous families, the two kinds of sporange present no differentiation in their early stages. The *archespore* is the terminal hypodermal cell of an axial row. This divides into the *sporogenous tissue* surrounded by the layer of *tapetal cells* formed from the innermost of the layers of cells into which the wall of the sporange divides. In the megasporanges one of the spore-mother-cells grows more vigorously than the rest, which gradually abort. In this

FIG. 26.—*A*, fertile branch of *S. inæqualifolia* (half natural size). *B*, longitudinal section of upper part, showing microsporanges and megasporanges. (After Goebel, magnified.)

FIG. 27.—Section of megasporange of *S. inæqualifolia*, showing double wall of sporange, layer of tapetal cells, and megaspores. (After Goebel, magnified.)

privileged cell are formed four spores, the number usually present in the mature megasporange. The microspores are formed in the same way as in the other heterosporous families. After falling out of the sporange they frequently adhere together in fours. The microspore has three coats—*endospore*, *exospore*, and *epispore*—of which the innermost is composed of cellulose. The coat of the megaspore is also treble, and the epispore is not unfrequently beautifully granulated and spiny. The dehiscence of both kinds of sporange is caused by the unequal contraction of the epidermal cells. The microsporanges are 2-valved, the megasporanges 3-4-valved.

The genus Selaginella (Spring) includes above 300 species, spread over the whole globe, but most abundant in the tropics. Many species resemble Lycopodium very closely in habit, but are more moss-like, and the leaves generally more delicate ; in others the stem is erect, and they reach the magnitude of small shrubs. Several species are favourite objects of cultivation from the beautiful metallic lustre of the leaves. They are readily propagated non-sexually, a small fragment of the stem producing a new plant if kept warm and moist on loose earth, owing to the production of adventitious roots in the angle formed by the branching of the 'vascular' bundle of the leaf from that of the stem. No economical use is known of any species either of Selaginella or Isoëtes.

Order 2.—Isoëteæ.

In the single genus Isoëtes the general phenomena of the life-history correspond to those of Selaginella, but with some important differences. Some weeks after the escape of the megaspore from the decaying megasporange its cavity becomes filled, by free-cell-formation, with a number of naked primordial cells, which gradually fill up the whole cavity of the endospore, and then become converted into a cellular tissue by the investment of each with a cell-wall of its own. The endospore at the same time thickens, and separates into several layers with a finely granular structure. The epispore, or outer layer of the coat of the megaspore, now splits at its apex by a three-rayed fissure, exposing the endospore, which also subsequently disappears, and a portion of the spherical prothallium is thus laid open. At its exposed apex appears the first archegone, which is followed by others if the first is not fertilised. The archegones resemble those of Selaginella, except that each of the rows which constitute the neck is composed of four instead of two cells.

The microspores are yellowish grey, and of the form of the quadrant of a sphere. The antherozoids are very long, slender, and attenuated

at both ends, where they are provided with two tufts of very long cilia ; in some species they are remarkably large. Their period of 'swarming' does not last more than about five minutes.

The stem of Isoëtes is distinguished by its extraordinarily restricted

FIG. 28.—*Isoëtes lacustris* L. (natural size).

growth in length, and the complete absence of branching, as well as by a remarkable secondary increase in thickness. It is completely covered by the bases of the leaves, leaving no part exposed. Its upper portion has the form of a shallow funnel, with the apex depressed in its centre.

The long-continued increase in thickness which distinguishes this genus alone among Vascular Cryptogams—except possibly Botrychium (Filices)—is dependent on an internal layer of meristem which surrounds the axial 'vascular' cylinder, and continually produces new layers of paren-



FIG. 29.—*A*, megaspore of *Isoëtes acustris* L. *B*, prothallium; *a*, archegone (× about 50). (After Hofmeister.)

chyme on the outside. This takes place especially in either two or three directions, so that a corresponding number of projecting masses of tissue are formed, which slowly die off on the outside; and between them lie the same number of furrows meeting on the ventral side of the stem, which has hence the appearance of a laterally elongated plate or

FIG. 30.—*I. lacustris*. *A—D*, microspore, showing stages in formation of antherid and antherozoids (× 580). *v*, sterile cell; *a—d*, stages in formation of antherozoid (× 580); *e*, *f*, mature antherozoid (× 700). (After Millardet.)

disc. In these furrows are produced a large number of rows of roots in acropetal succession. In the stem is a single cauline axial bundle composed of short reticulate and spiral tracheides, surrounded by a rudimentary phloem without sieve-tubes. From this axial bundle there

proceeds a branch into each leaf and one into the root. The layer of meristem which surrounds the axial bundle increases chiefly in the centrifugal direction, fresh layers thus formed replacing the outer ones, which continually die off. The secondary long-enduring increase in thickness of the stem is chiefly due to increase in thickness of the cortical tissue, though new xylem-elements are also produced. The mode of apical growth differs from that in most species of Selaginella. There is no single apical cell, the apex of the stem being occupied by a group of equivalent merismatic cells.

FIG. 31.—Longitudinal section of stem of *I. lacustris*. b—b^8, leaves; r^1—r^{10}, roots: the ligules are shaded (× 30). (After Hofmeister.)



FIG. 32.—Longitudinal section through lower portion of leaf of *I. lacustris* (diagrammatic). *L*, ligule; *Ind*, indusium; *Sp*, microsporange; *Tr*, trabecules; *Gf*, vascular bundle of sporophyll. (After Goebel.)

The leaves of Isoëtes are very elongated, cylindrical, and quill-shaped, and are arranged in a complicated phyllotaxis on the very short stem. They are segmented into a basal portion, the *sheath* or *glossopode*, and an apical portion, the *lamina*. The sheath is nearly triangular in form with a very broad insertion, and does not completely embrace the stem. It is convex behind and concave in front, where it bears the sporange in a large depression known as the *fovea*; the margin of this depression rises in the form of a thin membranous outgrowth, the *veil* or *indusium*, which, in many species, extends above and beyond the sporange. Above

the fovea, and separated from it by a ridge called the *saddle*, is a smaller depression, the *foveola*, the lower margin of which forms a lip-like structure, the *labium*, and from its base rises a narrow membranous acuminate structure, the *ligule*, with a cordate base, and usually projecting above the foveola. The sheath passes above into the lamina, which is narrow and thick, almost cylindrical, but flattened in front, contains chlorophyll, and is traversed by four wide longitudinal air-cavities, segmented by transverse septa. A rosette of these fertile leaves or *sporophylls* is produced annually, but between these whorls are alternate whorls of *phyllades*, or imperfect leaves, consisting, in the submerged species, of only a small lamina with no sheath, while in the terrestrial species they are reduced to mere scales. Stomates occur in the paludose and terrestrial, but not in the submerged species. Scattered spiral tracheides are found in the parenchymatous base of the leaf. The fundamental tissue, which



FIG. 33. Development of microsporange of *I. lacustris*. *t*, tapetal cells; *Tr*, trabeculæ: the archespore and sporogenous cells derived from it are shaded. (After Goebel, magnified.)

is not separated from the single 'vascular' bundle by a bundle-sheath, has a strong tendency to become sclerenchymatous, especially beneath the epiderm and in the sheath. The very simple bundle which occurs in each leaf is stated by Russow to be collateral, the xylem and phloem lying side by side.

The roots spring from the furrows of the stem, and resemble, in structure and mode of branching, those of Selaginella. There is no rhizophore.

The *sporanges* of Isoëtes do not make their appearance until the third year after germination. Each sporophyll bears only a single sporange, which is undoubtedly a product of the leaf, and is situated below the ligule in the fovea, to which it is attached by a narrow base. The outer leaves of the fertile rosette produce megasporanges only, the inner leaves microsporanges only. Both kinds of sporange originate from a group of cells at the base of the leaf. The archespore is derived

from a hypodermal layer of cells. In the formation of the microsporange the archespore-cells elongate in a direction at right angles to the surface, and divide by transverse walls. Some of these rows of cells are then arrested in their growth, lose their abundant protoplasm, and divide into elongated tabular cells constituting the *trabecules*, which cross the sporange from the dorsal to the ventral side. The remaining cells develop into the mother-cells of the microspores, an external layer having been previously separated as tapetal cells. In the development of the megasporange the processes are the same as far as regards the formation of the tapetal cells and the trabecules ; the mature sporange may contain either four or a much larger number of megaspores. The mode of development of the megaspores presents perhaps the closest analogy to that of the secondary embryo-sacs of Gymnosperms that occurs in any order of Vascular Cryptogams ; and the same remark applies to the formation of the microsporanges and pollen-sacs. Both kinds of sporange are indehiscent, the spores escaping only by the decay of the tissue.

In both kinds of spore the epispore is frequently granulated, tuberculate, or echinate ; and in some species there are two kinds of microspore differing from one another in this respect.

One or two species of Isoëtes display the phenomenon of *apogamy* in various degrees. In extreme cases the formation of the megasporange is arrested at a very early stage, and its place supplied by a vegetative shoot, which becomes detached and develops into an independent plant.

The number of species of Isoëtes is about fifty, the greater part inhabitants of the warmer portions of the globe. They somewhat resemble Pilularia in general habit. Some species are aquatic and entirely or partially submerged, other paludose, and a very few terrestrial ; and they present corresponding differences in the structure of their tissue, presence of stomates, &c.

Literature.

Von Mohl—(Stem of Isoëtes) Linnæa, 1840, p. 181.
Braun—Ueber Isoëtes, Monber. Berlin Akad. Wiss., 1863.
Hofmeister—Entwick. d. Isoëtes lacustris, Abhandl. Sächs. Gesell. Wiss., 1865.
Pfeffer—Entwick. d. Keims Selaginella, in Hanstein's Bot. Abhandl., iv., 1871.
Tchistchakoff—(Isoëtes) Nuov. Giorn. Bot. Ital., 1873, p. 207.
Bruchmann—Wurzeln v. Lycopodium u. Isoëtes, 1874.
Hegelmaier—Bot. Zeit., 1874, p. 481.
Goebel—(Apogamy of Isoëtes) Bot. Zeit., 1879, p. 1.
Mer—(Sporange of Isoëtes) Compt. Rend., xlii., 1881, p. 310 ; and Bull. Soc. Bot. France, 1881, pp. 72, 109.
Kienitz-Gerloff—(Embryo of Isoëtes) Bot. Zeit., 1881, pp. 761, 785.
Vines—(Isoëtes) Annals of Botany, vol. ii., 1888, p. 117.

ISOSPOROUS VASCULAR CRYPTOGAMS.

Class III.—Lycopodiaceæ.

The Lycopodiaceæ are a comparatively small group of plants comprised in only four genera, differing from one another greatly in habit, but agreeing in the prevalence of a dichotomous rather than of a monopodial mode of branching in both stem and root, though this is by no means universal. Growth is effected by a group of equivalent cells in the growing point, never (except in Psilotum, Sw.) by a single apical cell. The leaves are always of small size and entirely undivided ; in Psilotum they are reduced to mere scales, and this genus is also entirely rootless ; while in Phylloglossum (Kze.) the underground stem is tuberous. The sporanges and spores are of one kind only ; the spore produces on germination (where this has been observed) a green or colourless prothallium, which carries on a much more independent existence than is the case in the heterosporous orders, and bears, in the cases which have been examined, both archegones and antherids. The position of the sporanges varies. In the Lycopodieæ it corresponds to that of the Selaginelleæ, on the upper side of the base of the leaf, or they are crowded on special erect branches ; and here the sporanges are unilocular ; while in the Psiloteæ they are plurilocular, and are grouped on the main stem or on short lateral branches. Further details are best described under the heads of the two orders into which the Lycopodiaceæ may be divided. In the monœcious prothallium the Lycopodiaceæ approach the Ophioglossaceæ ; while in the structure of the sporophyte they display a remarkable resemblance to the heterosporous Selaginellaceæ.

Order 1.—Lycopodieæ.

In this order are included two genera of very different habit : Lycopodium (L.), with nearly 100 known species ; and Phylloglossum (Kze.), with only one. The form and appearance of the oophyte vary greatly even in the different species of the typical genus Lycopodium. The germination of the spores of L. inundatum (L.) has been described as follows by de Bary :—The endospore bursts in the form of a nearly spherical vesicle through the exospore, which splits into three valves ; the germinating filament which originates in this way then divides by a septum into a basal cell, which undergoes no further change, and a larger apical cell, which divides into two rows of segments ; each segment further divides by a tangential wall into an inner and an outer cell, so that

the young prothallium now consists of an axial row of four short cells, the basal and apical cells, and two lateral rows. The cells contain a few grains of chlorophyll. The formation of the sexual organs was not observed. The mature oophyte of L. annotinum (L.) presents several important differences. The prothallium is underground and of a yellowish-white colour, destitute of chlorophyll, and consists of a tuberous mass with cushion-like ridges on the upper side and a few small rhizoids. On its upper side and completely imbedded in the tissue are a number of antherids, consisting of cavities covered by one or more layers of cells, and containing a large number of mother-cells of antherozoids. The antherozoids themselves appear to be minute bodies consisting of only a few coils, and probably with two cilia. The archegones have not been actually observed, but are evidently borne on the same prothallium

FIG. 34.—Young plant of *Lycopodium annotinum* L. *p*, prothallium; *w*, root (natural size). (After Fankhauser.)



FIG. 35.—*A*, prothallium of *Lycopodium cernuum* L.; *t*, tuberous outgrowth (× 25). *B*, young plant of *L. cernuum* (×2). (After Treub.)

as the antherids, and in close contiguity with them, apparently on the upper side, in the depressions between the ridges. Only one archegone appears to be fertilised on each prothallium. The young sporophyte has no foot, its place being supplied by a tuberous swelling with root-hairs. A very different type of prothallium is presented, according to Treub, by L. cernuum (L.). It consists of a short cylindrical axis half immersed in the soil, containing chlorophyll in its exposed portion, and putting out rhizoids from its lower end. The upper extremity bears a tuft

of small leaf-like lobes, beneath which are the archegones and antherids on the same prothallium, and buried in its tissue. Each antherid arises from a single superficial cell, which divides by a transverse septum into an outer stigmatic cell, subsequently splitting up into three, and a basal cell in which the antherozoids are formed. The archegone has a very short neck, consisting of three rows of cells. While the prothallium displays greater differentiation than is the case elsewhere in Vascular Cryptogams, the embryo or young sporophyte is, on the contrary, of very simple structure, and entirely parenchymatous. A *cotyledon* is formed at one end, but there is no primary root, or other differentiation of organs. In L. Phlegmaria (L.) the prothallium is cylindrical, without chlorophyll, and branches freely. The rhizoids proceed from a superficial layer, which also branches laterally, by which means fresh prothallia are constantly being formed without any production of sexual organs. The archegones and antherids are borne on the upper surface of the prothallium, and are always accompanied by *paraphyses*, barren tubular cells of rare occurrence among Vascular Cryptogams. The



FIG. 36.—Prothallium of *L. Phlegmaria* L. Bifurcate branch bearing antherids, *a*, and paraphyses, *p* (× 90). (After Treub.)



FIG. 37.—*Lycopodium clavatum* L. *A*, sporangiferous branch (natural size); *B*, sporange and subtending leaf (greatly magnified); *C*, spore, showing lines of fissure (still more magnified).

antherids are scattered or in groups, and produce biciliated antherozoids. The archegones appear later than the antherids on the thickened extremities of the same branches. They project above the surface of the prothallium, and have from three to five canal-cells. In the formation of the sporophyte it is possible that we have a transition between Vascular Cryptogams and Muscineæ. The oosphere divides by a wall vertical to the axis of the archegone into two cells, of which the one nearest the neck becomes the *suspensor*, while the other is the mother-cell of the *embryo*. The first root is but slightly endogenous. The cells of the prothallium of all known species of Lycopodium are liable to be infested by an endophytic Pythium, the zoospores of which have very probably been taken for antherozoids. The oophyte of Phylloglossum is unknown. Of the development of the sporophyte of the Lycopodieæ very little is known in its early stages.



FIG. 38.—*Phylloglossum Drummondii* Kze. *l*, leaves; *r*, roots; t_1, old tuber; t_2, new tuber; *a*, sporangiferous axis (× 3). (After Bower.)

In the typical genus Lycopodium the sporophyte resembles Selaginella in habit. The stem of most species is procumbent, extending in the case of L. clavatum (L.) to several feet, and putting out here and there a few roots into the soil. Less vigorous branches rise erect, and are sporangiferous. The procumbent species display a tendency to bilateralness, especially in the structure of the axial 'vascular' bundle. In other species the much shorter stem grows erect, and puts out roots from its lower portion, which in some cases grow downwards through the tissue of the stem, emerging only as a tuft at its base. In some of these erect species, especially in the tropics, the stem is stout and shrubby. One or two species of Lycopodium have climbing stems; a few are epiphytic. The sporophyte of Phylloglossum, its only known condition, has a striking resemblance to the embryonic condition of Lycopodium. The erect unbranched stem is very short, rising into a slender scape, which bears at its extremity a spike of sporanges, and at its base a rosette of long subulate leaves; otherwise the stem bears only a few very rudimentary leaves. The plant is reproduced by adventitious shoots consisting of a tuber with a leafless rudimentary bud.

As far as has been at present observed, the growing end of the stem of Lycopodium has no single apical cell, and the same is true of the leaves and of the roots. The growing point of the stem corresponds

closely with that of Gymnosperms. It is composed of a small-celled primary meristem, in which no differentiation can be detected into



FIG. 39.—Transverse section of stem of *L. annotinum*. O, epiderm; A R, outer cortex; J R, inner cortex; S, sclerotised fundamental tissue; P, vascular bundle-sheath; H, xylem; B, phloem-portion of axial vascular cylinder; Bsp, leaf-traces (magnified).

dermatogen and periblem, the rudiments of the 'vascular' bundle penetrating nearly to its apex. In some species it projects in the youngest leaves in the form of a low cone ; in others the apex is flat. As in Flowering Plants, the leaves and rudiments of the shoots do not arise from single cells, but from groups of cells which include both the outermost and the subjacent layers of the primary tissue of the growing point.

The branching of the stem of Lycopodium is either monopodial or dichotomous in its origin ; but in the latter case one of the bifurcations usually greatly exceeds the other in vigour of growth. The branches are never placed in the axils of leaves, as in Flowering Plants, but usually arise from the stem above a leaf, but without any definite relation to it. In several species two new growing points of equal strength appear side by side on the flat apical surface, and continue to develop dichotomously. In others the rudiment of the new branch takes the form of a lateral protuberance on the greatly elevated cone of growth. On the stems of many species the small leaves are so closely packed that the internodes are completely suppressed.

The internal structure of the stem of Lycopodieæ presents several peculiarities. The cells of the fundamental tissue are sometimes uniformly thin-walled, but usually the inner layers in particular have thicker walls, and the cells are prosenchymatous, or even have their walls strongly sclerotised, reminding one of the sclerenchymatous layer in ferns; but they are never coloured brown. The axial 'vascular' cylinder is separated from the cortical fundamental tissue by a strongly developed bundle-sheath, composed of from one to three layers of cells. Air-cavities and mucilage- and gum-passages sometimes occur in the fundamental tissue of the stem and the leaves. The 'vascular' bundles themselves present a striking peculiarity in Lycopodium, forming in the stem and root a single axial cylinder, usually with a circular outline. In this compound bundle are plates or bands of xylem, which are either completely isolated, or coalesce in various ways so as to form figures which are divided into two similar halves by an axial longitudinal section. The cylinder may therefore be described as displaying a bilateral symmetry. If transverse sections are made at different heights in the stem, the xylem presents different figures, in consequence of the bands anastomosing in their course. The elements of these xylem-bands are, like those of ferns, tracheides pointed at both ends, and increasing in breadth towards the interior, the most common form of thickening being pitted or scalariform rather than annular or spiral ; the latter are found only at the outer edges of the bands. The whole mass of xylem-bands is surrounded by a narrow-celled phloem, containing, in the larger species, sieve-tubes. Between the outer edges of the xylem-bands and the

periphery lie the bast-like cells known as 'protophloem-elements.' Within the bundle-sheath the phloem is surrounded by several layers of larger cells, corresponding to the phloem-sheath of ferns. Though the sclerotised tissue is much less developed than in ferns, the axial 'vascular' bundle is, in the stouter species of Lycopodium, surrounded by a ring of fibres composed of several layers. The axial bundle is cauline, and may be followed out, in a rudimentary condition, to very near the apex. In Phylloglossum the short stem is traversed by a single narrow bundle, which is very weak, and has no scalariform, only a few spiral and annular tracheides.

The roots of Lycopodium originate on the outside of the axial cylinder; their internal structure is similar to that of the stem. In the erect species they have their origin at a considerable height in the stem, whence they strike downwards through its fundamental tissue, in which they sometimes even branch dichotomously, and emerge in the form of a tuft at the base of the greatly thickened stem. In the creeping and climbing species they emerge separately, and dichotomise in the soil in intercrossing planes. The epiderm of the root is often strongly cuticularised. In Phylloglossum the underground portion of the stem consists of two ovoid tubers of different age (see fig. 38), which are destitute of the least trace of 'vascular' bundles. From above these tubers spring a few adventitious roots, which are of endogenous origin, do not branch, and each of which has a single concentric axial bundle.

The leaves are very small in most species of Lycopodium, and invariably narrow, simple, and sessile; sometimes with a long apiculus. They are sometimes adpressed to the stem with the exception of the free apex; more often they are entirely free. In some species the form and size of the leaves vary greatly even on the same individual plant, and these heterophyllous species often display more or less of a bilateral structure. The phyllotaxis is sometimes verticillate, sometimes spiral, or both arrangements occur together in the same species. In the verticillate species the leaves are either decussate, or in whorls of three, four, or more; on creeping stems they are usually placed on a transverse zone oblique to the axis; and the number of leaves in a whorl varies even on the same branch. The small and extremely variable divergences of the leaves in the species with spiral phyllotaxis are very remarkable. Each leaf is always penetrated by a single central 'vascular' bundle without any lateral branches; it is of very simple structure, and is in connection with the axial cylinder of the stem. In L. albidum (Bak.) the leaves are membranous, and quite destitute of chlorophyll. The epiderm is provided with stomates, either on the under surface only or on both surfaces, and frequently collected into groups. The fundamental

tissue is sometimes furnished with air-cavities and gum-passages, usually in connection with the 'vascular' bundle. In some of the heterophyllous species these occur only in the sporangiferous or fertile leaves. In Phylloglossum the leaves all spring from the base of the scape (see fig. 38); they are narrow and subulate, about half an inch in length, and penetrated by a single 'vascular' bundle. They are colourless in their basal half, green in their apical half, and have stomates only in the green part. In a large number of species of Lycopodium all the leaves serve the purpose of nutrition only, and the sporanges are borne in the axils of ordinary leaves. But in the remaining species of Lycopodium, and in Phylloglossum, the leaves which subtend the sporanges are greatly modified, being of a membranous texture and colourless. In these species the sporanges with their subtending leaves are usually collected into spike-like 'inflorescences,' which may be short, erect, bifurcate branches, as in L. clavatum, or an elongated naked scape, as in Phylloglossum.

The *sporanges* of Lycopodium are seated each on the base of a leaf which has frequently undergone more or less metamorphosis (see fig. 37, *B*); by displacement they may subsequently become axillary. They are kidney-shaped, and are attached at their broader side by a short thick pedicel. They are unilocular, and dehisce by a fissure across the apex in the longest diameter. In all the Lycopodiaceæ the outer walls of the epidermal cells of the sporange are composed of pure cellulose, while the inner and side walls are lignified. Dehiscence takes place by the outer face of these cells contracting more than their inner face in dry air. The small and numerous spores are sphero-cubical, the exospore being marked in a variety of ways. On germinating the exospore splits by three fissures which meet in a point at the apex of the spore, the endospore projecting between the three valves thus formed. The sporange originates as a prominence from a group of superficial cells at the base of the leaf. The original cells from which it is formed are few in number; the central one of these gives rise to the *archespore*. The wall of the sporange ultimately consists of from two to four layers of cells; the innermost of these forms the layer of *tapetal cells*. The mother-cells of the spores become separated from one another, and invest themselves with very thick cell-walls; from each is developed four spores, and the exospore becomes elevated into warts, spines, &c., before the walls of the mother-cells have become absorbed. In Phylloglossum the sporanges are also unilocular, and are placed in the axil of short triangular apiculate metamorphosed leaves; and a large number are collected into a spike-like 'fructification' at the extremity of a naked scape. They dehisce by a vertical longitudinal fissure. The spores

are excessively minute, and have three radiating lines meeting at the apex. Their germination is unknown.

In most species of Lycopodium vegetative propagation takes place by means of axillary bulbils, which become detached; and in some, adventitious buds are also produced lower down on the stem. L. cernuum produces similar gemmæ or bulbils on the root. Phylloglossum is propagated by the lateral budding of its underground tubers, in a manner somewhat similar to our native species of Orchis.

The species of Lycopodium are scattered over the whole globe from the polar to the equatorial regions, the greater number growing on elevated ground or in swamps; some are epiphytic. The monotypic Phylloglossum Drummondii (Kze.) is a native of swamps in Australia and New Zealand. Several species of Lycopodium have an ancient use as cathartics. The spores are used in the manufacture of pills, and have the property, from the large quantity of oil which they contain, of keeping the hands dry when dipped in water. Those of L. clavatum are collected in large quantities, especially in Northern Germany, for pyrotechnic purposes. The British species are popularly known as 'club-moss' and 'stag's-horn moss.'

Order 2.—Psiloteæ.

This order is composed of the two very small tropical genera Psilotum (Sw.) and Tmesipteris (Bernh.), of the latter of which very little is known, it never having been examined in the living state.

The stem is erect, and is penetrated, in Psilotum, by a cauline 'vascular' bundle of very simple structure, which is circular on transverse section, and is surrounded by a bundle-sheath. It always branches dichotomously. Psilotum is entirely rootless, the function of roots being performed by remarkable underground branches of the stem, which dichotomise like the aërial shoots. These underground shoots have a three-sided apical cell, and are entirely destitute of a root-cap. Those only which are nearest the surface have a few whitish subulate rudimentary leaves; these may turn upwards, develop chlorophyll, and become ordinary aërial shoots. Those branches which strike deeper into the soil are slenderer, and the rudiments of leaves are reduced to groups of a few cells which remain buried in the tissue, not projecting above the surface. They resemble true roots in their single axial 'vascular' cylinder. Psilotum triquetrum (Sw.) produces minute gemmæ or bulbils, which remain dormant for a time, and from which the plant not unfrequently makes its appearance apparently spontaneously in orchid- and palm-houses. The leaves of Tmesipteris are erect, elliptical,

FIG. 40.—*Psilotum triquetrum* Sw. *A*, fertile branch (natural size); *B*, portion of the same (magnified); *C*, smaller portion and sporange (still more magnified).

and apiculate, and are penetrated by a single 'vascular' bundle; those which subtend the sporanges are much smaller, and apparently deeply bifid, in consequence of their becoming connate at their base in pairs. In Psilotum the leaves are reduced to mere scales without any 'vascular' bundle.

The sporanges of the Psiloteæ differ from those of the Lycopodieæ in not being formed in connection with the leaves, and in being pluri-locular. They are collected into spikes which are formed at the growing point of a primary shoot. In Tmesipteris each spike usually consists of two sporanges only, situated in the fork between two connate or one bifid fertile leaf; they are oblong and bilocular, and dehisce by two vertical slits; the spores are very minute, oblong, and curved. In Psilotum the sporanges are collected into groups of three or four on special short lateral branches, each in the axil of a rudimentary leaf, and forming a loose spike. They are turbinate in form, and are divided into three compartments, less often into two or four; each locule dehisces by a vertical fissure. The germination of the spores and the oophyte generation are entirely unknown in the order.



B A

FIG. 41.—*Tmesipteris tannensis* Bernh. *A*, portion of branch (natural size); *B*, sporange and subtending leaves (magnified).

Psilotum consists of two species, natives of the tropical regions of both hemispheres, having the appearance of small branching nearly leafless shrubs; Tmesipteris of a single known species, epiphytic on the trunks of tree-ferns, with a pendulous habit, in the Southern Hemisphere.

LITERATURE.

Spring—Monograph des Lycopodiacées in Mém. Acad. Roy. Belgique, 1842 and 1849.
Cramer—(L. Selago) in Nägeli u. Cramer's Pflanzenphys. Unters. Heft i., 1855.
De Bary—(Germination) Naturf. Gesell. Freiburg, 1858.
Mettenius—(Phylloglossum) Bot. Zeit., 1867, p. 97.
Payer—Botanique Cryptogamique, 1868.
Juranyi—(Psilotum) Bot. Zeit., 1871, p. 177.
Hegelmaier—Bot. Zeit., 1872, pp. 798 *et seq.*, and 1874, p. 773.

Fankhauser—(Prothallium) Bot. Zeit., 1873, p. 1.
Strasburger—Bot. Zeit., 1873, pp. 81 *et seq.*
Bruchmann—Ueb. Wurzeln v. Lycopodium u. Isoëtes, 1874.
Beck—(Prothallium) Oesterr. Bot. Zeitschr., 1880, p. 341.
Bertrand—(Psilotum) Compt. Rend., xcvi., 1883, pp. 390, 518; (Phylloglossum) do., xcvii., 1883, pp. 504 *et seq.*
Solms-Laubach—(Psilotum) Ann. Jard. Bot. Buitenzorg, 1884, p. 139, and 1886, pp. 217, 233.
Treub—(Prothallium) Ann. Jard. Bot. Buitenzorg, 1884, p. 307, and 1886, p. 87; see Nature, xxxi., 1885, p. 317, and xxxiv., 1886, p. 145.
Galloway—(Spores) Bull. Torrey Bot. Club, 1885, p. 55.
Bower—(Phylloglossum) Proc. Roy. Soc., xxxviii., 1885, p. 445.
Bruchmann—(do.) Bot. Centralbl., xxi., 1885, pp. 23, 309.
Treub—(Prothallium) Ann. of Bot., i., 1887, p. 119.
Goebel—(Prothallium) Bot. Zeit., 1887, pp. 161, 177.

Class IV.—Filices.

Ferns (under which term the Ophioglossaceæ are also included in popular language) are by far the most numerous and best known class of Vascular Cryptogams. In some families, however, as the Marattiaceæ and Schizæaceæ, much yet remains to be made out with regard to the history of development, and their exact position in the circle of affinity must remain for a time doubtful.

The germinating spore develops into the prothallium by the bursting of the cuticularised exospore, and the rapid growth and division of the contents of the endospore into a plate of cells. Before germination the contents of the spore become invested with a new cellulose membrane. But the tabular prothallium does not always result directly from the contents of the spore. In the Hymenophyllaceæ the spore undergoes division, even before the rupture of the exospore, into three cells, one of which only attains great development, dividing by transverse septa, and branching until it greatly resembles the protoneme of a moss; the flat prothallia then springing from lateral shoots. In most of the Polypodiaceæ, which include by far the greater number of the genera of ferns, and in the Schizæaceæ, the contents of the spore develop directly into a short segmented filiform protonemal structure, which expands at the apex into a cordate or reniform plate of tissue, consisting at first of only a single layer of cells. If a single apical cell is present, it soon disappears, and is replaced by a growing point situated in a depression at the anterior end of the prothallium, behind which a cushion, several layers in thickness, is formed by tangential cell-

divisions. The prothallium is most commonly monœcious, though the sexual organs may not appear at the same time, and is strictly bilateral or dorsiventral, the result, according to Leitgeb (Sitzber. Akad. Wiss. Wien, lxxx., 1880, p. 201), of the action of light. The archegones



FIG. 42.—Germination of prothallium of fern, with exospore still attached. *a*, *b*, *Dicksonia antarctica* Lab. (× 240); *c*, *d*, *Aspidium filix-mas* Sw. (× 120). (After Luerssen.)

are found exclusively (except in Marattiaceæ) on the under side of the cushion. Rhizoids are produced in large numbers on the under side of another part of the cushion; the antherids also on the under side among the rhizoids, or less often on the margin. In Gymnogramme

leptophylla (Desv.) the prothallium is many-lobed, and the archegones and antherids are produced on separate conical tuber-like outgrowths from its under side, which penetrate into the soil, where they are perennial, and may give birth by budding to new prothallia, while the sporophyte generation is annual. The prothallium is occasionally, in the Hymenophyllaceæ, reduced to a single row of cells terminating in an antherid, or even to a single cell. Campbell has detected continuity of protoplasm in the cells of the prothallium of Struthiopteris germanica (L.). In the Osmundaceæ the prothallium springs directly from the spore without any intermediate protoneme, a plate of cells being formed on germination by longitudinal and transverse divisions; the first rhizoid is formed out of a posterior cell. The ribbon-shaped prothallium of Osmunda (L.) is characterised by the presence of a midrib composed of several layers of cells running along its whole length. The archegones are produced on the under surface on this midrib; the antherids either on the margin or on the under surface with the exception of the midrib. An approach towards a higher type of organisation is indicated by the tendency of the prothallium to become dicœcious in the Osmundaceæ, and in Struthiopteris (L.). All the spores from the same sporange sometimes produce male prothallia, *i.e.* such as bear antherids only, the



FIG. 43.—Under side of prothallium of *Aneimia Phyllitidis* Sw. *sh*, cushion, with archegones; *a*, antherids and rhizoids (× 25). (After Bauke.)



FIG. 44.—Antherid of *Adiantum capillus-Veneris* L., in different stages. *p*, prothallium; *a*, antherid; *s*, antherozoids; *b*, vesicle with starch-grains (× 500).

archegones being produced later, and in smaller numbers, on female prothallia ; or the same prothallium may produce first antherids and subsequently archegones, when it may be termed proterandrous. This is remarkably the case also in Gymnogramme. In Cystopteris fragilis (Bernh.) (Polypodiaceæ) Campbell states that there are two kinds of prothallium, a smaller male and a larger hermaphrodite. The prothallium of ferns is sometimes propagated vegetatively by the production of adventitious shoots from single marginal cells, which become detached and form independent prothallia. This takes place especially in Hymenophyllaceæ and in Osmunda, but occurs also in Polypodiaceæ,

FIG. 45.—Archegone of *Adiantum capillus-Veneris*, in various stages. *A*, *B*, *C*, *E*, in longitudinal, *D*, in transverse section ; *h*, neck ; *sl*, canal-cells converted into mucilage ; *s*, ventral canal-cell ; *e*, oosphere ; in *E* divided into a 2-celled embryo (× 800). (After Goebel.)

abundantly in Gymnogramme (see Cramer, Denkschr. Schweiz. Naturf. Gesell., 1880). The prothallium of Vittaria (Sm.) produces peculiar stalked bulbils.

The *antherids* of ferns are small papilliform projections on the under side or margin of the prothallium (very rarely on the upper side), produced among the rhizoids, and of similar origin, *i.e.* from a single superficial cell ; in the Hymenophyllaceæ they are produced also on the protonemal filaments. The protuberance becomes separated by a septum from the parent superficial cell, and then sometimes divides at once into the parent-cells of the antherozoids. But more often the

papilla divides first of all into a central cell surrounded by a single layer of peripheral cells. These last are barren, but contain chlorophyll; while the central cell divides still further, each derivative nearly cubical cell then producing a flat spirally-coiled antherozoid contained within a vesicle, or 'special parent-cell.' In no case is the number of antherozoids produced by a single antherid very considerable. The function of the peripheral cells appears to be to absorb water violently when the antherid is mature, in consequence of which they swell up considerably and rupture the central cells, thus causing the escape of the parent-cells of the antherozoids. From each of these is then discharged, by the bursting of its delicate cell-wall, an *antherozoid* consisting of a flat band of protoplasm coiled spirally three or four times, and bearing at its anterior end a number of fine cilia (fig. 44). To its posterior end is frequently attached for a time a vesicle containing starch-grains, which is probably the remains of the special parent-cell of the antherozoid. As in other Vascular Cryptogams, the body of the antherozoid appears to be formed from the nucleus of the mother-cell, the cilia from the cell-protoplasm.

FIG. 46.—Archegone of *Pteris serrulata* L. at the moment of the expulsion of the mucilage (× 350). (After Strasburger.)

The *archegones* are produced on the under side of the cushion of the prothallium, but in much smaller numbers than the antherids. Like them, each archegone is derived from a single superficial cell, which at first bulges only slightly, and is first divided into three cells by two tangential walls. The lowermost of these three, or *basal cell*, divides further, and takes its share in the formation of the *venter*, or swollen part of the archegone, which is altogether imbedded in the prothallium. The outermost of the three cells develops into the neck-wall, or outermost wall of the *neck* of the archegone, dividing at first into four cells, from which the four rows of cells which constitute the neck are formed by oblique septa. Since the neck grows more rapidly on the anterior side, *i.e.* the side nearest to the apex of the prothallium, and hence becomes convex on that side, the number of cells is also larger in the

anterior rows of the neck, the usual number being six, while there are seldom more than four in the concave posterior side. From the middle one of the three primary cells arises the whole of the axial row of cells of the archegone, consisting of the *central cell* and the *canal-cells*. During the development of the neck this middle cell becomes pointed upwards, and forces itself between the neck-cells ; this pointed portion becomes divided off by a septum, and now forms the single *neck-canal-cell*, which lengthens as the neck lengthens. The large central cell now breaks up into an upper and smaller *ventral canal-cell* and a much larger lower cell, the protoplasmic contents of which subsequently become rounded off, and constitute the *oosphere*. According to Campbell, the ventral canal-cell is wanting in Struthiopteris germanica (L.). The walls of the canal-cells swell up and become converted into mucilage, and finally this thin mucilage, together with the protoplasm of the canal-cells, is expelled from the open neck. The antherozoids are retained by the mucilage, and collect in large numbers before the archegone ; a number of them force themselves into the canal of the neck, and of these some eventually reach the oosphere, and coalesce with it, entering it at a light-coloured spot near the neck, which is termed the *receptive spot*. After impregnation the neck closes up. It is very rare for more than one archegone to be fertilised on the same prothallium, and the enormous majority of prothallia perish without producing any sporophyte generation.

The ordinary course of the alternation of generations is occasionally interrupted by *apogamy* or by *apospory*, the suppression respectively of the oophyte or of the sporophyte generation. The former has been observed especially in Pteris serrulata (L. fil.), the latter in particular varieties of Athyrium filix-fœmina (Bernh.), and of Polystichum angulare (Willd.). In apogamy the non-sexual fern-plant springs directly from the prothallium without the intervention of a fertilised archegone. In apospory a prothallium is produced on the surface of the frond, either in the locality where the sorus would normally be found, or less often elsewhere, and may assume unusual forms, sometimes that of a solid cylindrical body, but bears normal archegones and antherids. Bower classifies the various forms of substitutionary or correlative growths connected with the suppression of the sporophyte generation under three heads, viz.—(1) simple prolification ; (2) sporophytic budding ; (3) apospory. The first hardly occurs among ferns. The second is illustrated by the familiar formation of bulbils in species of Asplenium (L.), Cystopteris (Bernh.), &c., in which the formation of the buds cannot be directly correlated with arrest of spore-formation. In apospory we get a more or less complete sporal arrest, but this may vary in degree. In some instances

the substitutionary growths which accompany the arrest of spore-formation are restricted to the sporanges themselves. These are replaced by 'pseudo-bulbils' of a pear-like form, presenting but little resemblance to ordinary prothallia, but demonstrating their oophytic character by producing antherids. In other examples the prothalloid growths are by no means restricted to the sporange; they may either arise from the sorus itself, or may appear at points quite distinct from the sori, and even on fronds which bear no sori at all. There is here a distinct transition from



FIG. 47.—Apogamous shoot of *Pteris serrulata*, on the under side of the prothallium *p*; *b*, first leaf; *v*, apex of stem; *w*, rudiment of first endogenous root (× 80). (After de Bary.)

FIG. 48.—Prothalloid growth of *Polystichum angulare* Willd. var. *pulcherrima*, originating from surface of frond; *arch*, archegone (× 10). (After Bower.)

sporophyte to oophyte without the intervention of spores. Comparatively little is known about the oophyte generation in the Hymenophyllaceæ, but it would appear as if apogamy were a very common, perhaps even normal occurrence in some species of Trichomanes (Sm.); and here the two phenomena have even been observed on the same individual, the oophyte and sporophyte generation succeeding one another without the production of either spores or sexual organs.

The fertilised oosphere or *oosperm* becomes immediately invested

with a cell-wall of its own, and develops by cell-division into the *embryo*, from which springs the young sporophyte, commonly known as the fern-plant. The first division-wall in the oosperm is always nearly vertical; and two others follow, at right angles to it and to each other, dividing the oosperm into octants. From the anterior of the two original halves are derived the growing point of the *stem*, and the *cotyledon* or first leaf; from the posterior half the *foot* by which it is attached to the prothallium, and the first *root*. Until the differentiation of the first leaf and the fixed root, the embryo remains imbedded in the surrounding tissue of the prothallium, which grows with its growth. The primary root is always small; in the Hymenophyllaceæ it disappears early, and in many

FIG. 49.—*Asplenium decussatum*; adventitious bud, *k*, already rooting (natural size).



FIG. 50.—Young sporophyte of *Adiantum capillus-Veneris* still attached to prothallium *p*; *b*, first leaf; *w'*, *w''*, 1st and 2nd root; *h*, rhizoids of prothallium (× 30). (After Goebel.)

species of Trichomanes no subsequent roots are formed, their place being supplied by underground branches.

The mature fern varies in size from that of the 'filmy ferns,' species of Hymenophyllum not above an inch in height, with delicate moss-like habit, to the stately 'tree-ferns' of the Southern Hemisphere (Cyatheaceæ and Dicksonia, L'Herit.), fifty or sixty feet in height. The stem is either ascending and vertical, or creeping on or beneath the surface of the soil, or occasionally scandent (Lygodium, Sw.), often very short with undeveloped internodes, and the leaves so crowded that frequently no portion of the stem—then often called a *caudex*—remains exposed; while in the creeping and climbing species the leaves are often

separated by long internodes. The ultimate roots are always adventitious ; that is, there is no predominant axial root which is a prolongation downwards of the main axis of the plant, as in many Dicotyledons and some Monocotyledons. They are usually very numerous, especially in tree-ferns, arising in acropetal succession, and completely clothing the lower part of the stem, or, where this is suppressed, the leaf-stalks, as in the case of the common 'male fern' (Aspidium filix-mas, Sw.). The ultimate branches of the root are furnished with a root-cap as in Flowering Plants. The leaves, or, as they are more commonly called, 'fronds,' are invariably stalked, and are remarkable in many species, especially when they attain a large size, for the great extent to which subdivision of the lamina is carried; in some tree-ferns they attain a length of from six to ten feet. In the filmy ferns they consist, as in Muscineæ, of only a single layer of cells penetrated by distinct 'vascular' bundles. Stomates, similar in structure to those of Flowering Plants, occur abundantly both on the under side of the leaf and on the leaf-stalk, except in the Hymenophyllaceæ. The leaves exhibit very little metamorphosis compared to those of Flowering Plants. Most usually all the leaves are alike in form and extent of division, and even nearly so in size ; but in some species only certain of the leaves, *sporophylls*, are fertile, and these then show a more or less well-marked difference from the barren leaves, as in our 'hard fern' (Lomaria spicant, Desv.) and 'parsley fern' (Cryptogramme crispa, R. Br.). In the 'elk's-horn fern' (Platycerium alcicorne, Desv.), commonly grown in cultivation, the leaves are alternately broad thallus-like barren plates, closely applied to the surface on which the plant grows, and long erect dichotomously branched fertile leaves. The leaves (except the first, which spring from the prothallium) are *circinate* in vernation, both the principal rachis and the midrib of the pinnæ (when the leaf is pinnate) being rolled up on their upper side—owing to the more rapid growth of the cells on the under than those on the upper surface—and only slowly unroll as the development of the leaf advances. The young leaves, and the rachis and petiole of mature leaves, are generally more or less completely clothed with *pales* or ramenta, flat brown scale-like trichomes or outgrowths of the epiderm. These are often glandular, and sometimes contain crystals of oxalate of lime. They serve as a protective mantle to the young stem and leaves, and also as a reservoir of moisture. Capitate, glandular, and other more ordinary kinds of hair are also of frequent occurrence. In the typical ferns the sporanges are also trichomic in their origin. They are collected into groups or *sori* (fig. 58), usually formed in connection with a 'vascular' bundle. In unilamellar leaves these sori are placed on the edge, in all others almost invariably on the under side of the leaf, especially of its

apical portion, and they then assume a great variety of shapes—circular, reniform, crescent-shaped, linear, or they are concealed beneath the revolute margin of the leaf. The sorus may or may not be covered by a membrane called the *indusium*, an outgrowth of the epiderm. In the

FIG. 51.—*Asplenium Adiantum-nigrum* L.; rhizome with fronds showing circinate vernation (natural size). *a*, under side of fertile pinnule (magnified).

Cyatheaceæ they assume the form of a cup; in the Hymenophyllaceæ they are situated at the extremity of a vein at the apex or margin of a pinna. In some instances, as in our native 'flowering' or 'royal fern' (Osmunda regalis, L.), the sori completely consume, in the course of their

development, the parenchyme of the fertile (apical) part of the leaf, giving the appearance of a panicled or thyrsoid inflorescence. The *sporange* is usually stalked, and has an elliptical form, or that of a battledore or racket-bat. The sides are commonly thin and membranous, and the sporange dehisces either longitudinally or transversely, generally from the elasticity of an *annulus* or ring of brown thick-walled cells running along or across it. The position of this annulus, and its more or less complete development or entire absence, are useful characters in the subdivision of the class. In the Marattiaceæ the sporanges are of altogether different origin, being developed from hypodermal masses of cells; and transitional forms occur between the two. The *spores* are minute, very commonly reniform, or often nearly cubical, resembling pollen-grains in structure, usually furnished with two coats, an *exospore* and *endospore*, the latter of which is sometimes double, and the former generally marked with papillæ, reticulations, &c.



FIG. 52.—Diagram of tip of leaf of *Ceratopteris thalictroides* Brongn. *S*, apical cell; *L*, lateral lobe of leaf. (After Kny.)

A more minute description must now be given of the structure and peculiarities of the various organs.

The great distinguishing feature which characterises the development of the stem of ferns, as contrasted with that which occurs in all Flowering Plants (Gymnosperms and Angiosperms), is the presence of a single *apical cell*, from which the whole of the growing point or apical meristem originates, and which may therefore be recognised as the parent-cell of the whole of the tissue subsequently formed. This apical cell is usually wedge-shaped in creeping stems with a bilateral structure, a three-sided pyramid in erect or ascending stems. The growing apex of the stem is frequently completely hidden in the youngest leaf-bud, but in other species there is a considerable intervening space. In some Hymenophyllaceæ leafless prolongations of the stem assume the appearance and the function of roots. As contrasted with Flowering Plants, especially Exogens, the stem of ferns is characterised by the small extent to which it branches;

and this is true not only of the erect columnar stem of tree-ferns, but also of the creeping or erect stem of smaller species. Axillary branching is very rare, if it ever actually occurs ; the terminal branching is always dichotomous, never sympodial. The fundamental tissue of the stem and leaf-stalk consists, in many species, entirely of thin-walled parenchyme. In others, and especially in tree-ferns, portions of it undergo a change in the great thickening and brown colouring of the cell-walls, the cells becoming at the same time prosenchymatous. In this *sclerenchyme* of the fundamental tissue the sclerosis may take place in individual isolated cells ; more often the cells so affected are united into conspicuous bands or sheaths. In many Polypodiaceæ and Osmundaceæ the entire cortex assumes eventually a dark colour. In the common brake (Pteris aquilina, L.) two thick sclerenchymatous bands of this description lie between the inner and outer 'vascular' bundles, while another continuous layer immediately underlies the epiderm. The firmness and solidity of the stem of tree-ferns are mainly due to strongly developed sclerenchymatous cylinders which form complete sheaths surrounding the 'vascular' bundles. The 'vascular' bundles themselves are always *closed* or destitute of cambium ; in the stem, except in Osmunda, and usually in the leaves, they are *concentric*, consisting of a central xylem-portion entirely enveloped in a layer of phloem ; in the stem and leaves of Osmunda, and in the leaves of some other ferns, they are *collateral*, the xylem and phloem portions lying side by



FIG. 53.—Transverse section of 'vascular' bundle of *Polypodium leiorrhizum* Wall. *p*, fundamental tissue ; *s*, sclerenchymatous sheath ; *b*, phloem ; *h*, xylem (× 200). (After Luerssen.)



FIG. 54.—Transverse section of stem of *Pteris aquilina* L. *r*, epiderm ; *p*, fundamental tissue ; *pr*, sclerenchymatous sheath ; *ig*, vascular bundle ; *ag*, outer network (somewhat magnified).

side. Besides a few narrow spiral tracheides, lying at definite points of the transverse section, the xylem consists mainly of *scalariform tracheides*, *i.e.* of tracheides with bordered pits which usually have the appearance of transverse clefts, their ends being mostly obliquely truncate or fusiform and pointed. *True vessels* occur but rarely in ferns ; *e.g.* in Pteris aquilina and in the rhizome of Athyrium filix-fœmina, where they are also scalariform. Between the tracheides lie narrow thin-walled cells which contain starch in winter. In the phloem, in addition to narrow parenchymatous cells, are sieve-tubes with well-developed sieve-plates, but forming true callus only in a small number of cases ; and at the circumference narrow bast-like thick-wall prosenchyme. Each individual bundle is usually immediately enclosed in a single distinct layer of narrower cells, the *vascular bundle-sheath* or *endoderm* ; this layer, which probably originates from the fundamental tissue, displays a strong tendency for its walls to become brown and suberised. In very young stems and those which permanently remain very slender, as in many Hymenophyllaceæ, there is a single axial bundle. But in stouter stems and leaf-stalks the central bundle is replaced by a network of anastomosing bundles, presenting, in typical cases, a cylinder of considerable diameter, by which the fundamental tissue is separated into an outer *cortical* and an inner *medullary* portion ; but isolated scattered bundles also arise in addition. The principal bundles which constitute this cylindrical network mostly have the form of broad plates with the margins curved outwards, each surrounded by its thick firm brown sclerenchyme-sheath ; they usually present the appearance of an interrupted ring near the periphery, but in Osmundaceæ the ring is more continuous. From the margins of these cauline bundles spring the more slender filiform bundles which pass into the leaves, the number of openings in the meshes of the cauline 'vascular' cylinder corresponding to that of the leaves. In the leaf-stalk the bundles may either run separately or may coalesce into plates. Terletzki (Pringsheim's Jahrb., 1884, p. 452) has detected continuity of protoplasm between the cells in the parenchyme of the rhizome in several species of fern ; the intercellular spaces also contain protoplasm which is in connection with that of the cells. In the aquatic genus Ceratopteris the stem contains large air-cavities. Elongated tannin-sacs occur in the parenchyme of the stem and leaf-stalk of many ferns, especially in the neighbourhood of the 'vascular' bundles. Gum- and mucilage-cells are also of frequent occurrence. Incrustations of calcium carbonate are not infrequent on the leaves. Round stalked glands occur in the fundamental tissue of the stem and leaves of Aspidium filix-mas.

The *leaves* of ferns stand either in two rows on the stem, or less often in a single dorsal row, or the phyllotaxis is more complicated and spiral : they are never opposite or whorled. Each leaf originates from a superficial cell of the growing point distinguished by a stronger swelling of its outer cell-wall. The petiole is the first part of the leaf formed ; the lamina then begins to appear at its apex, and itself develops from the base to the apex. In many ferns the leaves of the mature plant are characterised by the extraordinary slowness of their development. In old plants of Pteris aquilina the formation of the leaf commences fully two years before it begins to unfold ; at the commencement of the second year only the leaf-stalk is as yet in existence ; during the summer of the second year the lamina begins to develop at the apex of the rod-like petiole, and may be found hidden beneath the long hairs in the form of a minute disc. It then begins to bend downwards at its apex, and continues for a time in a pendent condition. It is only in the spring of the third year that the elongation of the leaf-stalk forces the lamina above the surface of the soil, and that the latter begins to unfold. In Aspidium filix-mas the development is almost as slow. The basifugal development of the lamina itself is also extremely slow in many ferns, the lower portions having long been fully formed while the apex is still unfolding. In several genera, as Gleichenia (Sm.) and Mertensia (Willd.), a periodical interruption occurs of the apical growth, this intermittent development even extending over several years. In Lygodium the primary pinnæ remain unfolded after the formation of a pair of pinnæ of the second order, while the rachis of the leaf grows without limit and resembles a twining stem, climbing in some cases to the height of fifty or one hundred feet, the pinnæ themselves presenting the appearance of leaves. In the anomalous Ceratopteris thalictroides, which grows in water, the ultimate segments of the leaves are swollen up in a pod-like manner. Goebel has shown (Ann. Jard. Bot. Buitenzorg, vii., 1887) that in the heterophyllous ferns, such as Platycerium alcicorne and several species of Polypodium (L.), one form of leaf is specially adapted, directly or indirectly, for the supply of nutriment to the plant, and is sometimes furnished with special aquiferous tissue. The stipular structures of the Marattiaceæ are quite peculiar among Vascular Cryptogams. The leaves of ferns not unfrequently display a tendency to branch dichotomously at the apex, but in other cases the branching appears to be monopodial. The leaf-stalk has frequently the power of producing adventitious leaf-buds (see fig. 49). In Struthiopteris germanica these develop into long underground stolons covered with scale-leaves, which at length rise above the surface, and bear at their apex a rosette of ordinary leaves. Nephrolepis (Sch.) is also furnished

with remarkable stolons, the extremities of which swell up into tubers ; but it is uncertain whether they belong to the stem or the root. In other cases adventitious leaf-buds are borne on the upper surface of the lamina or in the axils of the leaves. In Woodwardia radicans (Sm.) and some other species, the long drooping leaves may root in the soil and put up new shoots. The veins in the leaves of the great majority of ferns do not anastomose, but divide repeatedly dichotomously. Their 'vascular' bundles differ, in some cases, from those of the stem in being collateral, the xylem facing the upper, the phloem the under surface of the leaf.

With the exception of some species of Trichomanes, all ferns have true *roots*, characterised, like those of Vascular Cryptogams generally, and of Flowering Plants, by the presence of a true *root-cap*, composed of several layers of cells, and by the tendency of the epidermal cells to develop into long unsegmented filaments or *root-hairs*, by which the nutriment is absorbed from the soil. These proceed in acropetal succession from the creeping underground stem or rhizome, or, where the stem is erect, very short, and densely covered with leaves, from the leaf-stalks. In tree-ferns the lower part of the erect stem is densely covered with a thick felt-like envelope of slender roots, which give a broad base to the stem. Like the stem, the tissue of the root develops from a single apical cell. The branching is always monopodial. The lateral rootlets arise in acropetal succession on the outside of the primary 'vascular' bundle. The root is traversed by a single axial cylinder formed by the coalescence of 'vascular' bundles.

The epiderm of the leaves of those ferns which are not unilamellar differs in no essential respect from that of Flowering Plants, but its cells contain a larger quantity of chlorophyll. The epiderm of the under side only is abundantly provided with stomates, which are usually of quite the ordinary structure. In some cases, as Aneimia (Sw.), they present the peculiar appearance of the two guard-cells being entirely enclosed within a single epidermal cell. Those of Kaulfussia (Bl.) (Marattiaceæ) are very large, and of peculiar structure.

The *sporanges* of ferns are rounded, ovoid, or pear-shaped capsules, seated on long stalks in the Polypodiaceæ and Cyatheaceæ, sessile in the other orders. When mature, the wall of the capsule consists, except in the Marattiaceæ, of a single layer of cells, a particular row of which, running longitudinally, transversely, or obliquely round the capsule, usually undergoes special development, and is known as the *annulus* ; but the annulus may be entirely wanting, as in the Marattiaceæ, or its place may be taken by the special development of an apical or lateral group of cells, as in the Osmundaceæ. Where there is

an annulus the sporange dehisces by a fissure at right angles to it ; in the Marattiaceæ it opens by an apical pore. The dehiscence is due to unequal contraction in drying of the unequally thickened portions of the cell-walls of the annulus. On one side of the sporange of a considerable number of ferns are two, three, or four cells of peculiar form, with lignified cell-walls, the *lip-cells*, between which the dehiscence always begins, and which appear to guide its direction. These cells together are sometimes called the 'stomium.' Among the spo-



FIG. 55.—Sporange of *Aspidium filix-mas*, showing annulus *an*, lip-cells *lc*, and paraphyse attached to stalk. (After Kündig, greatly magnified.)



FIG. 56.—Development of sporange of *Asplenium Trichomanes* L. *s*, archespore; *r*, annulus (× 550). (After Goebel.)

ranges are frequently slender segmented filaments, or *paraphyses*. In some Polypodiaceæ there is a single (rarely more than one) outgrowth from the stalk of the sporange, resembling a capitate hair, and sometimes septated internally. It is regarded as a paraphyse, and may probably be an undeveloped sporange. The entire *sorus* may be covered by the recurved margin of the leaf, or by a true *indusium* belonging to the epiderm, or by a false indusium, consisting of an outgrowth of the hypodermal tissue, composed of several layers of cells. In Enterosora (Bak.), from British Guiana (Polypodiaceæ), the sporanges spring from the base of spherical chambers in the under surface of the

leaf, which open only by a narrow slit. Although the sori usually originate on the veins of a leaf, and only on the under side of the lamina, this is not always the case. In Olfersia they spring from both surfaces of the lamina by the side of the midrib ; and in this and other species of Acrosticheæ from the mesophyll as well as the veins. In the Hymenophyllaceæ they are enclosed in a cup-like indusium, and are attached to the apex of the veins, which projects beyond the margin of the leaf. The spot on the fertile vein which bears a sorus is known as the *placenta* or receptacle. In the Polypodiaceæ, and probably also in the Cyatheaceæ, each sporange originates from a single epidermal cell, which swells up considerably, and is cut off from the rest of the leaf by a septum. This mother-cell of the sporange subsequently divides by another septum into a basal cell which develops into the pedicel, and an apical cell which becomes the capsule. The *pedicel* usually consists ultimately of three rows of cells produced by longitudinal and transverse divisions. The nearly hemispherical mother-cell of the capsule first divides, by four successive oblique walls, into four parietal and a nearly cubical central cell, the *archespore*. In the parietal cells further divisions follow at right angles to the surface ; while from the archespore are formed four tabular segments parallel to the parietal cells, which again divide by walls vertical to the surface into one or two layers of *tapetal* or *mantle-cells*, constituting together the *tapete*. The row of cells in the wall of the sporange which constitute the annulus are the result of divisions at right angles to the surface of the sporange ; their outer walls bulge out and project above the surface. The tapetal cells ultimately disappear, and the whole of the space within the wall of the capsule is occupied by a fluid, in which float the mother-cells of the spores formed by successive bipartitions of the archespore, and normally sixteen in number in the Polypodiaceæ and Schizæaceæ. In these families the formation of the spores takes place in the following way. Each of the sixteen spore-mother-cells divides into four by two successive bipartitions of the protoplasmic contents, preceded by corresponding divisions of the nuclei. Sixty-four spores are thus normally formed in each sporange. They then invest themselves with a cell-wall, which is usually double, consisting of an inner coat, or *endospore*, generally but not always composed of cellulose, and sometimes itself consisting of two layers, and an outer brown cuticularised *exospore*, provided with ridges, papillæ, warts, or other elevations. Leitgeb states that in Osmunda and some other genera there is no true endospore ; while in Onoclea (L.), according to Campbell, there are three coats. The spores of the Osmundaceæ and Hymenophyllaceæ contain chlorophyll. Another mode of spore-formation, more analogous to what takes place

in the formation of the pollen of Flowering Plants, occurs sometimes in the two orders just named, and invariably in the Cyatheaceæ, Osmundaceæ, and Hymenophyllaceæ. Each spore-mother-cell divides at once by cell-walls of cellulose into four compartments, sometimes called 'special mother-cells'; within each of these compartments the spores become invested with their permanent cell-walls, both the walls of the original mother-cells and their septa being then absorbed. The spores formed in this way have a rounded cubical shape, while those produced in the mode first described are bilateral, and very commonly kidney-shaped. In the Schizæaceæ and Osmundaceæ the sporanges are not strictly trichomic in their origin, being formed, before the differentiation of the epiderm, each from a single cell, which may be regarded as the rudiment of a leaf-branch. In both these orders the number of spores produced in a sporange is much larger than in the Polypodiaceæ; in this and in other respects they manifest an approximation to the Marattiaceæ and Ophioglossaceæ. The sporanges of Marattiaceæ are altogether endogenous in their origin, being developed from an internal mass of tissue, and are destitute of an annulus. On the dehiscence of the normal sporange the spores are at first attached to the annulus, and are detached and thrown to a distance by its sudden return to its original position. The spores of many Polypodiaceæ retain their vitality and power of germination for a long period, and require a longer or shorter period of rest before germination; those of the Hymenophyllaceæ, on the other hand, often begin to germinate while still enclosed in the sporange. In Scolopendrium (Sm.), according to Beck (Verhandl. Zool.-Bot. Ges. Wien, 1879), the exospore does not burst, but decays at the spot where the germinating filament emerges.

The non-sexual propagation of ferns takes place chiefly by means of the adventitious buds already described, which appear on the lamina or petiole of the leaf (fig. 49). As a normal phenomenon it is, however, confined to a small number of species known in cultivation as viviparous or bulbiferous ferns, such as Asplenium bulbiferum (Forst.) and Cystopteris bulbifera (Bernh.). The occasional vegetative propagation of the prothallium has also been already described.

Ferns are distributed over the whole globe, from the equator to the arctic zone; and, from the ease of their culture and the beauty of their forms, are favourite objects of cultivation. They are most abundant in moist warm climates, and hence enter largely into the composition of all insular floras. In the tropics a large number of species are epiphytic; and it is only there, and in the islands of the Southern Hemisphere, that they attain the size of tree-ferns. One or two species are annual, and a single one, Ceratopteris thalictroides, is aquatic, while a very few have a

climbing habit. The number of species which are at the present time applied to any economical purpose is extremely small. The common brake, Pteris aquilina, is largely used throughout Northern Europe for forage purposes, and for the stuffing of rough beds and pillows. The so-called 'male fern,' Aspidium filix-mas, has a very ancient repute as a vermifuge, and is still occasionally employed for that purpose, as also are several other species to a less extent in different parts of the globe, either for a similar purpose or as astringents and mucilages. In several species, especially tropical, the stem contains sufficient starch to be esculent. The number of known species of ferns is at present estimated at about 3,000, but it is constantly and rapidly increasing. They are arranged in a comparatively small number of genera, the limits of which are often extremely difficult to define.

LITERATURE.

Mettenius—Filices Hort. Bot. Lipsiensis, 1856.
Hofmeister—Abhandl. Sächs. Gesell. Wiss., vol. v., 1857 ; and Pringsheim's Jahrb. wiss. Bot., 1863, p. 278.
Wigand—Botanische Untersuchungen, 1854.
Newman—History of British Ferns, 4th ed., 1865.
Reess—(Sporange) Pringsheim's Jahrb. wiss. Bot., 1867, p. 217.
Strasburger—(Fertilisation) *ibid.*, 1869, p. 390.
Kny—(Antherid) Monber. Akad. Wiss. Berlin, 1869, p. 416 ; and Pringsheim's Jahrb. wiss. Bot., 1869, p. 1.
Hooker and Baker—Synopsis Filicum, 1868.
Goebel—(Prothallium of Gymnogramme) Bot. Zeit., 1877, pp. 671 *et seq.*
Weiss—(Bundle-sheath) Flora, 1880, p. 119.
Cramer—(Fertilisation) Denkschr. Schweiz. Naturf. Gesell., 1880.
Haberlandt—(Vascular Bundles) Sitzber. Akad. Wiss. Wien, lxxxiv., 1881, p. 121.
Lachmann—(Root-organs) Compt. Rend., xcviii., 1884, p. 833 ; and ci., 1885, p. 592.
Klein—Bot. Zeit., 1884, pp. 577 *et seq.*
Schmidt—(Dehiscence of Sporange) Flora, 1885, pp. 451, 471, and 1887, pp. 177, 202 ; and Ber. Deutsch. Bot. Ges., 1886, p. 396.
Campbell—(Prothallium) Bot. Gazette, 1885, p. 355.
Thomæ—(Leaf-stalk) Pringsheim's Jahrb. wiss. Bot., 1886, p. 99.
Goebeler—(Pales) Flora, 1886, pp. 451 *et seq.*
Gardiner and Ito—(Mucilage-cells) Ann. of Bot., i., 1887, p. 27.
Campbell—(Struthiopteris germanica) Mem. Boston Soc. Nat. Hist., 1887, p. 17.
Kündig—(Sporange) Hedwigia, 1888, p. 1.

The following classification of the Filices into families or orders follows partly the plan proposed by Mettenius, partly that adopted in Hooker and Baker's 'Synopsis Filicum.' It must, however, be distinctly borne in mind that the divisions are of very unequal value. The first three are closely allied to one another ; the Hymenophyllaceæ show con-

siderably greater divergence ; the affinity of Osmundaceæ and Schizæaceæ with the typical ferns is more remote ; while the Marattiaceæ exhibit so many points of divergence that some high authorities have removed them altogether from the Filices.

Order 1.—Polypodiaceæ.

This order includes by far the largest number of genera and species, and may be regarded as the typical family of ferns. The sporanges arise out of single epidermal cells, and have usually long pedicels ; they

Fig. 57.—*Alsophila aculeata* Klotzsch, a tree-fern (reduced).

have an incomplete vertical annulus, and consequently dehisce transversely. The sori vary greatly in size and form, and usually consist of a large number of sporanges. They are seated on the under side of the divided or undivided leaf, upon the veins ; much less often (Acrosticheæ) also on the upper surface or in connection with the mesophyll ; they are usually covered by an indusium in the Asplenieæ, Aspidieæ, and Davallieæ, but not in the Acrosticheæ or Polypodieæ. In the great majority of species all the leaves are ultimately fertile; but in some genera (Gymnogramme (Desv.), Lomaria (Willd.), Platycerium (Desv.), &c.)

there are species with barren and fertile leaves, differing from one another in habit and in degree of segmentation ; in other heterophyllous

FIG. 58.—Sori of Polypodiaceæ (variously magnified). *a, b, Adiantum; c, Lindsaya; d, e, Blechnum; f, g, Cystopteris; h, i, Davallia; k, l, Cheilanthes; m, n, Pteris; o, p, Woodwardia; q, Scolopendrium; r, s, Asplenium; t, Aspidium; u, v, Woodsia; w, Gymnogramme; x, z, Polypodium.* (After Luerssen.)

ferns (species of Polypodium) both forms of leaf are fertile. A few of the Polypodieæ are arborescent (Dicksonia L'Herit.).

Principal genera :—Adiantum (L.), Cheilanthes (Sw.), Pellæa (Link), Pteris (L.), Lomaria (Willd.), Blechnum (L.), Woodwardia (Sm.), Doodia (R. Br.), Asplenium (L.), Scolopendrium (Sm.), Aspidium (Sw.), Nephrodium (Rich.), Lastrea (Presl), Nephrolepis (Sch.), Polypodium (L.), Nothoclæna (R. Br.), Gymnogramme (Desv.), Hermonitis (L.), Vittaria (Sm.), Acrostichum (L.), Platycerium (Desv.), Onoclea (L.), Woodsia (R. Br.), Ceterach (Adans.), Dicksonia (L'Herit.), Davallia (Sm.), Cystopteris (Bernh.), Lindsaya (Dry.).

Order 2.—Cyatheaceæ.

This order is not distinguished from the Polypodiaceæ by any very clear lines of demarcation. The sporanges are epidermal and shortly stalked ; they have a complete oblique eccentric annulus. The sori are seated on a receptacle or placenta which often projects considerably ; they are naked, or are more often enclosed in a cup-shaped indusium or 'involucre,' which sometimes forms a closed cup opening by a transverse fissure ; the sporanges are densely crowded in the sorus. The greater number of the tree-ferns belong to this family (fig. 57). The leaves are often very large (five to ten feet in length), and usually compoundly pinnate, forming a rosette at the summit of the columnar unbranched arborescent stem, which is densely covered with aërial roots, especially in its lower portion, and is marked in a diamond pattern by the scars of fallen leaves. In some species, in addition to the ordinary cylinder of 'vascular' bundles, there are a number of accessory bundles distributed through the medulla and cortex, forming a delicate open network. Crystals of calcium oxalate are not uncommon in the epidermal cells.

Fig. 59.—Sorus of *Cyathea*, with open indusium (magnified).

Principal genera :—Cyathea (Sm.), Hemitelia (Br.), Alsophila (R. Br.).

Order 3.—Gleicheniaceæ.

The sporanges are epidermal and sessile, with a broad complete transverse annulus running nearly horizontally, and hence with vertical dehiscence. The sori are naked, on the under side of ordinary leaves, and usually consist of only three or four sporanges. The stem is a slender creeping rhizome ; the leaves are usually very large, and with peculiar buds or 'innovations' on the lamina.



Fig. 60.—Sporange of *Gleichenia* (magnified).

Principal genera :—Gleichenia (Sm.), Mertensia (Willd.).

Order 4.—Hymenophyllaceæ.

The oophyte generation is known but in a very few species of Hymenophyllaceæ Where it has been observed (some species of Hymenophyllum and Trichomanes) it differs from that of other ferns, and is usually filiform, closely resembling the protoneme of a moss, but somewhat coarser. Antherids appear to be produced at the middle of these filaments, and archegones at their extremity. But apogamy is much more common in the Hymenophyllaceæ than in any other family of ferns, and it is doubtful whether it does not even occur regularly in some species. Bulbils or gemmæ are produced abundantly on the prothallium, consisting of a small number of cells, and borne on pedicels or *sterigmas*. They germinate with extreme slowness. In Trichomanes pyxidiferum (L.) the prothallium is frequently an aposporous growth, derived from imperfect arrested sporanges, or even from cells of the placenta. The archegones are borne on peculiar structures, known as *archegoniophores*, massive outgrowths of the prothallium, each archegoniophore bearing either a single archegone or several. The archegoniophore is usually a multicellular structure, and the venters of the archegones are imbedded in its tissue. T. alatum (Sw.) is habitually apogamous, and is possibly never reproduced sexually. Aposporous prothallia spring in great numbers from all parts of the frond, often quite independent of the sori, and are more flattened and ribbon-like in structure than those of most Hymenophyllaceæ. They produce large numbers of stalked gemmæ. Archegones have never been observed in this species, and the antherids are imperfect, and apparently functionless The spores are multicellular before germination. The archegones differ from those of other ferns in having a perfectly straight neck.



Fig. 61.—Archegoniophore of *Trichomanes pyxidiferum* L., bearing five archegones, *ar*, of different ages (× 175). (After Bower.)

The sporanges have a complete horizontal or oblique annulus (incomplete in Loxsoma, R. Br.), and hence dehisce by a vertical fissure. They are borne on a prolongation of the fertile vein, the *columel*, which projects beyond the margin of the leaf, and is enclosed in the cup-shaped indusium. The columel or placenta elongates by intercalary growth, and the sporanges are produced on it in a spiral line in basifugal succes-

sion. In Hymenophyllum (L.) and Trichomanes (Sm.) the sporanges are sessile and biconvex, and are attached to the columel by one of their convex faces; the annulus, projecting in the form of a cushion, separates the two convexities, and is usually oblique, dividing the circumference into two unequal portions: in Loxsoma they are pear-shaped and distinctly stalked. Paraphyses occur only in a few species of Hymenophyllum. The mesophyll of the leaf consists, in the two larger genera, of only a single layer of cells, and the leaves have hence a filmy and moss-like appearance; but in Loxsoma there are several layers of cells, and the leaf is then provided with stomates. The stem is generally creeping and mostly very slender, and is penetrated by a single axial 'vascular' bundle.

FIG. 62.—*Hymenophyllum tunbridgense* (natural size).

FIG. 63.—*Hymenophyllum*; sporanges exposed (magnified).

Many species of Trichomanes are rootless; and the stem is then densely clothed with root-hairs, and slender ramifications of the stem assume the appearance and function of true roots. Even the ordinary branches of the stem have often been long formed before their leaves emerge from a rudimentary condition. In some species of Trichomanes the fructification is confined to special fertile leaves.

The Hymenophyllaceæ include but three genera, of which Loxsoma comprises a single species only, of creeping habit, native of New



FIG. 64.—*A*, germinating spore and prothallium of *Hymenophyllum*; *B*, *C*, *D*, stages in development of prothallium (× 120). (After Luerssen.)

Zealand; Hymenophyllum and Trichomanes nearly 100 species each, exceedingly delicate and graceful ferns, growing mostly on the trunks of trees and damp rocks, often within reach of the spray of waterfalls, in the moister and warmer parts of the globe. The smaller species of Hymenophyllum are known as 'filmy ferns.' The Hymenophyllaceæ may be regarded as the simplest and are probably the oldest family of ferns, and possibly form a connecting link between the Muscineæ and the Vascular Cryptogams.



FIG. 65.—Prothallium of *Osmunda regalis* L. *a*, antherids; *w*, rhizoids; *v*, growing point. (After Goebel.)

LITERATURE.

Mettenius — Ueber die Hymenophyllaceen, 1864.

Janczewski and Rostafinski — (Prothallium) Mém. Soc. Nat. Sc. Cherbourg, xix., 1875.

Goebel — (Germination) Ann. Jard. Bot. Buitenzorg, vii., 1887, p. 57.

Prantl — Untersuchungen zur Morphologie der Gefässkryptogamen, Heft 1.

Bower — Annals of Botany, vol. i., 1887, pp. 183 and 269.

ORDER 5.—OSMUNDACEÆ.

The prothallium of the Osmundaceæ is characterised by its strong tendency to propagate itself vegetatively, by means of adventitious shoots, and is commonly dicœcious, springing directly from the spore. It is usually ribbon-shaped, with a well-defined midrib

The sessile or shortly stalked, roundish, but unsymmetrical sporanges are not strictly epidermal in their origin. They bear on one side of their apex a modified annulus in the form of a group of cells of peculiar form, and dehisce vertically on the other side. Todea (Willd.) presents no difference between the fertile and sterile leaves; while in Osmunda (L.) the fructification has the appearance of a continuous or interrupted panicle,

FIG. 66.—*Osmunda regalis* L. Portion of frond (natural size).

FIG. 67.—Mucilage-gland from *Osmunda regalis* (magnified). (After Gardiner.)

from the entire absorption of the mesophyll of the fertile part of the leaf. In some species of Todea the leaf consists of only a single layer of cells. In Osmunda there are

FIG. 68.—Sporange of *Osmunda* (magnified).

abundant mucilage-cells at the base of the leaf-stalk. The 'vascular' bundles of the stem are collateral, as contrasted with the concentric bundles of typical ferns, and their course bears more resemblance to that in Gymnosperms and in Dicotyledons. In the structure of the leaves, and in the structure and development of the growing point, Osmundaceæ exhibit a transitional condition between the typical ferns

and the Marattiaceæ. The young leaves (in Osmunda cinnamomea, L., and Todea superba, Col.) present the remarkable peculiarity of their apex being occupied by a well-marked triangular-conical apical cell.

The family includes only a very small number of species, comprised in the two genera Osmunda (L.) and Todea (Willd.). Osmunda regalis (L.), growing in bogs, with remarkably coriaceous leaves, is our 'royal fern' or 'flowering fern.'

Fig. 69.—*Lygodium palmatum* Sw. (reduced).

Literature.

Bower—Proc. Roy. Soc., xxxvii., 1884, p. 42; and Quart. Journ. Micr. Sc., 1885, p. 75.
Gardiner and Ito—Annals of Botany, vol. i., 1887, p. 27.

Order 6.—Schizæaceæ.

The ovoid or pear-shaped sessile sporanges are not strictly epidermal in their origin. The apex of the sporange is occupied by a cap-like zone of cells of peculiar form, and the dehiscence is vertical. In the genera Aneimia (Sw.) and Schizæa (Sm.) the fertile leaves have the paniculate appearance of an Osmunda. In Schizæa and Lygodium (Sw.) the sporanges are seated in two rows on the under side of very narrow pinnæ; and in Lygodium each sporange is enclosed in a pocket-shaped indusium. In Aneimia the two lowermost pinnæ form a long-stalked panicle, from which the mesophyll has disappeared. In Aneimia and Lygodium the sporanges spring originally from the margin of the leaf, but are eventually placed in the course of development on its under side. In Mohria (Sw.) they are placed on the back of the leaf, and are concealed by its recurved margin. The differentiation of tissues, both in the mesophyll of the leaf and in the 'vascular' bundles, is very slight. The stem is in general but feebly developed, and seldom branches; the leaf-stalk is penetrated by only a single 'vascular' bundle. The peculiar position of the stomates in some species of Aneimia has already been described.

The family of Schizæaceæ comprises a small number of species, included in five genera, of which Mohria (Sw.) and Trochopteris (Gardn.) are monotypic. Aneimia (Sw.) and Schizæa (Sm.) resemble the Osmundaceæ in their paniculate appearance. All the species of Lygodium (Sw.) present the appearance of climbing stems, from the peculiar structure and mode of growth of the leaves already described. The family is confined to the warmer parts of the globe.

LITERATURE.

Prantl—Untersuchungen zur Morphologie der Gefässkryptogamen, Heft 2, 1881; also in Engler's Jahrb. 1881, p. 297.

ORDER 7.—MARATTIACEÆ.

The Marattiaceæ differ more widely from the typical ferns than any of the families hitherto described, and are by many authorities separated from them into a distinct class. But the point of divergence most relied on, the endogenous origin of the sporanges, has lost much of its significance since it has been known that the Osmundaceæ and Schizæaceæ present connecting links in this respect. The aërial flat prothallium, the circinate vernation and general structure of the leaf, and the ultimate structure of the sori, present so many points of contact with the other orders of Filices, that it seems most desirable at present to retain them as an aberrant order of the class.

The prothallium, the development of which is very slow, is a dark green elliptical or heart-shaped plate of tissue, lying on the surface of the soil, consisting of one or more layers of cells, and with a projecting hemispherical cushion on the under side. Its development has been followed out in Marattia (Sm.) and Angiopteris (Hoffm.). Chlorophyll is formed in the spore as soon as it begins to germinate. The exospore bursts, and the first cell of the prothallium projects as a papilla between the two lobes. After a considerable time the first division takes place at right angles to the direction of growth, and the first rhizoid is formed. Further divisions follow rapidly, and the prothallium soon becomes a cellular tissue, and is distinguished from that of typical ferns by its deep green colour and by its moderately thick cuticle. One of the cells first formed becomes an apical cell, from which fresh cells are formed until the prothallium has assumed its ultimate cordate form. The antherids make their appearance, after a period of some months, on both sides of the prothallium, but especially on the ventral cushion. Their structure differs in some respects from that of typical ferns. From a single superficial cell are produced a central cell, two upper cells, and a triangular stigmatic cell, which is thrown off when the antherid is

mature. Within the central cell are produced from twenty to two hundred antherozoids, each in its own mother-cell. Both antherids and archegones are deeply sunk in the tissue of the prothallium. The archegones, only the uppermost part of the neck of which appears above the surface, are formed on the ventral cushion, very rarely on the upper side of the prothallium. Their development presents no very special features.

The sporophyte generation has, when mature, the habit and appearance of an ordinary fern. The stem is usually erect and short, with tuberous base, never attaining a greater height than from one to two feet; less often (Kaulfussia, Bl.) a creeping underground rhizome. It resembles the stem of Ophioglossaceæ and Isoëteæ in never branching; that of true ferns in being densely covered, when erect, with leaves as well as with roots, so that no portion is left exposed. The growing point has a single apical cell, and is concealed in a terminal rosette of large leaves. The fundamental tissue is everywhere traversed by long rows of tannin-cells; and lysigenous mucilage-cells abound in the petioles and in the parenchyme of the pith and cortex of the stem; they anastomose freely, and are continuous from the stem into the roots. The sclerenchymatous tissue, so characteristic of the parenchyme of the stem of typical ferns, is but feebly developed, or is altogether wanting, in the Marattiaceæ. The 'vascular' bundles are concentric, and resemble those of true ferns. A central xylem, composed of wide scalariform tracheides, is surrounded by the phloem; the bundle-sheath is wanting in the bundles belonging to the stem and leaves, but is present in those of the root. The bundles bend from the stem into the leaves in the ordinary way. The stem and rachis of the leaves are not covered with pales, as in true ferns, nor are they completely glabrous, as in the Ophioglossaceæ.



FIG. 70.—Base of leaf-stalk of *Marattia* cut through. *st*, stipule; *c*, commissure; *v*, anterior, *h*, posterior wing (natural size). (After Sachs.)

The leaves are thick, coriaceous, and very large, attaining in some species a length of from five to ten feet. They have a long and very stout petiole, which is channelled on the upper side; the lamina unfolds very slowly, and is either simply or doubly pinnate, less often palmate or digitate. They are furnished at their base with appendages peculiar to the Marattiaceæ among Vascular Cryptogams, the *stipules* or auricles. While still in the bud the leaves are rolled up in a circinate manner, and are entirely enveloped in the large stipules until the lamina unfolds. The pair of stipules belonging to each petiole form an anterior and a posterior chamber, separated from one another by a longitudinal wall termed the *commissure*. In the posterior chamber is the rolled-up leaf

to which the stipules belong, the two posterior wings of the stipule being folded together behind it ; while the chamber formed by the anterior wings encloses all the younger leaves. The stipules remain succulent, not merely as long as the leaves last, but even after the lamina has fallen off ; and adventitious buds are not unfrequent upon them. The leaf-stalk is articulated immediately above the stipule ; the leaf always becomes detached at this articulation by a smooth scar, leaving behind the base of the leaf-stalk with its succulent stipules. The primary and secondary pinnæ are attached to their respective rachis by similar articulations ; and at each articulation is a *pulvinus* or cushion, containing collenchymatous tissue. In the mesophyll of the leaf occur, in all the genera, outgrowths of the cell-walls bounding the intercellular spaces and projecting into them. Where the intercellular spaces are small these outgrowths have the form of humps and cones ; in larger spaces they elongate into long slender filaments which present a superficial resemblance to the hyphæ of Fungi, but are quite solid, and consist of cuticularised cellulose. They are found also less abundantly in the leaf-stalk, stem, and root. Layers and bundles of sclerenchyme occur in the mesophyll, but are only feebly developed and of a light colour. The leaves of Kaulfussia are characterised by the presence of remarkably large stomates on the under side, formed in the ordinary way, but distinguished by the great size of the orifice, and by the guard-cells forming a narrow ring, and being surrounded by two or three layers of epidermal cells, which are also arranged in a ring. There are only two semicircular guard-cells, and the structure of these organs is very different from that of the stomates of Marchantia, to which they bear a superficial resemblance. Lenticels occur in the leaf-stalk of many species. Spherocrystals have been found in the mesophyll and leaf-stalk of Marattia cicutæfolia (Kaulf.) and Angiopteris evecta (Hoffm.).

The roots arise endogenously from immediately beneath the growing point of the stem. They strike obliquely downwards through the succulent parenchyme of the stem, penetrating the network of the 'vascular' bundles, with which they may easily be confounded, and generally emerging from a leaf-stalk. They are of a lighter colour, greater thickness, and more delicate texture than those of true ferns, approaching those of Ophioglossaceæ. After entering the soil they branch copiously and apparently monopodially.

The sporanges are produced in large numbers on the under side of ordinary leaves, each being developed, not from a single cell, but from a group of cells. They are situated on the veins, and usually form two rows of sori, which cover the lateral veins either for their whole length or only near the margin of the leaf ; in Kaulfussia they are placed on

fine anastomosing branches between the lateral veins. The placenta or receptacle on which each sorus is seated is a cushion-like outgrowth of the vein. The sporanges are altogether destitute of an annulus ; the wall always consists of several layers of cells. In Angiopteris the sporanges which make up a sorus are quite distinct, ovoid, and sessile, and dehisce by a vertical fissure on the inner side. In all the other genera they are more or less confluent, and the entire boat-shaped sorus is then known as a *synange* ; but each sporange still dehisces separately by a vertical slit on its inner side ; or, in Danæa, by an apical pore. The coalescence is most complete in Kaulfussia, where the circular sorus has the appearance of a plurilocular basin. The sorus is usually surrounded by flat lobed hairs of epidermal origin, forming a kind of



FIG. 71.—*A*, under side of leaf of *Angiopteris caudata* ; *B*, of *Marattia* ; *s*, sori ; *C*, sorus of *Marattia* cut through, showing open sporanges. (After Goebel.)

involucre. The true indusium is sometimes altogether wanting. The sporanges originate from the tissue of the leaf itself. The placenta is first formed as a cushion-like outgrowth of the fertile vein, partly from the epiderm, partly from the subjacent tissue. On this originate, in Angiopteris, the separate sporanges as papillæ, each composed of a number of cells. In Marattia, however, while the two rows of sporanges are distinct, those of each row are confluent from the first, but each has its own archespore. The primary mother-cells of the spores are formed at an early period within the archespore.

The spores are formed in fours within their parent-cells, and resemble in general character those of typical ferns. Two different forms of spore sometimes occur in the same species, but they present no difference on

germination. The wall of the spore is composed of three layers ; the surface is covered by minute wart-like spines.

Vegetative propagation takes place with great facility in some species of this family. In Marattia cicutæfolia the leaves, or even the stipules, have only to be cut into small pieces, and placed in damp soil or in a bottle, when a number of adventitious buds will be developed in connection with the 'vascular' bundles.

The Marattiaceæ comprise only a very small number of species, almost entirely confined to the tropics, and included in the four genera Angiopteris (Hoffm.), Marattia (Sm.), Danæa (Sw.), and Kaulfussia (Bl.). With the exception of the stipules, and the great thickness of the leaves, they have quite the habit of ordinary ferns.

LITERATURE.

De Vriese and Harting—Monographie des Marattiacées, 1853.
Mettenius—Ueber den Bau von Angiopteris, 1863.
Luerssen—Mittheil. aus dem Gesammtgebiete der Bot., vol. i., Heft 3, 1872 ; and Bot. Zeit., 1872, p. 768 ; and 1873, pp. 624 and 641.
Rüssow—Vergleichende Untersuchungen, 1872, p. 185.
Holle—Bot. Zeit., 1875, p. 215.
Jonkman—Bot. Zeit., 1878, p. 129 ; and Archives Néerland., 1880.
Schenk—Ber. Deutsch. Bot. Gesell., 1886, p. 86.

Class V.—Ophioglossaceæ.

This small but very well-defined group, although popularly included with the Filices under the common denomination of ferns, differs from them in several important points of structure, in some of which a connecting link is furnished by the Marattiaceæ. The prothallium is underground and destitute of chlorophyll, exhibiting a similarity to that of Lycopodiaceæ rather than to that of true ferns. The stem rarely attains more than a few inches in height, and usually does not branch ; it is remarkable for its slowness in growth. It contains no sclerenchymatous layers. The leaves are not circinate in vernation, and the petiole is furnished at the base with lateral outgrowths, which have been compared to the stipules of Marattiaceæ. The venation is dichotomous or reticulate, and generally inconspicuous. The sporanges are completely endogenous in their origin, and are never borne on the under side of the green leaf, but on a separate branch of the leaf, altogether destitute of green parenchyme, and form a compound 'fructification,' resembling in appearance a spike or panicle : there is no annulus. The

'vascular' bundles of the stem are collateral; the primitive elements of the xylem are usually scalariform tracheides.

The oophyte generation has been observed in only a few species. In

FIG. 72.—*Ophioglossum vulgatum* L. *A*, p'ant (natural size); *B*, portion of sporophyll (magnified).

Ophioglossum (pedunculosum, Desv.) it has at first the form of a small round parenchymatous tuber, entirely buried in the soil and destitute of chlorophyll, from which is subsequently developed a cylindrical vermiform shoot, which grows erect by means of a single apical cell,

and very rarely branches, and then but slightly. The apex of this shoot finally appears above ground, develops chlorophyll, becomes lobed, and ceases to grow. When fully developed, the *prothallium* consists of an axial bundle of elongated cells surrounded by shorter parenchymatous cells; its upper surface is clothed with rhizoids. Its length sometimes amounts to as much as two inches, though generally it is much shorter; its breadth is always very small. In Botrychium (Lunaria, Sw.) the prothallium is a minute light brown or yellowish white ovoid mass of firm cellular tissue, subterranean and destitute of chlorophyll, and producing rhizoids from all sides. In both genera the prothallium is monœcious.



FIG. 73.—*Botrychium Lunaria* Sw. (nat. size).

The *archegones*, *antherids*, and *antherozoids* do not differ materially in structure from those of Filices.

FIG. 74.—Longitudinal section of prothallium of *Botrychium Lunaria*, showing archegones and antherids (× 50). (After Hofmeister.)

Neither archegones nor antherids are limited in their production to any particular part of the prothallium. In Botrychium (Sw.) the antherids are cavities in the tissue chiefly of the upper side of the prothallium, and the archegones are produced in their immediate vicinity; in Ophioglossum (L.) they project slightly above the surface. Before opening to discharge the antherozoids, they are covered by a few epidermal layers of cells. The mode of formation of the antherozoids resembles that of Marattiaceæ. Their mother-cells originate from repeated divisions of one or two cells of the inner tissue lying beneath these layers. They are comparatively large, and escape through a narrow opening in the layers of cells which originally covered the antherid. The archegone consists of a venter containing the central cell, and a neck composed of four vertical rows, each consisting of two or more cells, and only slightly projecting above the surface. The

venter is completely imbedded in the prothallium, its wall being developed out of the tissue of the latter.

The course of development of the embryo and rudimentary sporophyte has not yet been followed out in a sufficient number of instances to warrant a general description; in those in which it has at present been observed it appears to present some discrepancies.

The short *stem* is erect and entirely glabrous, often with a swollen tuberous base; only in Helminthostachys (Kaulf.) is there a creeping underground rhizome. In only a very few exceptional cases does it branch. The flattened apex of the stem consists of an irregular meristem derived from a single pyramidal apical cell. The fundamental tissue of the stem consists of short nearly cylindrical thin-walled succulent cells, which are longer in the leaf-stalk, interspersed with large intercellular spaces. It exhibits a striking difference from the corresponding tissue of true ferns in the entire absence of sclerenchymatous layers. It is separated by the 'vascular' cylinder into a cortical and a medullary parenchyme. It has an epiderm well provided with stomates, and exhibits sometimes a remarkable development of layers of cork. The 'vascular' bundles form (in Ophioglossum vulgatum, L.) a hollow cylindrical network, from each mesh of which is sent out a leaf-trace. The whole of the tissue which fills up the meshes is frequently transformed into scalariform tracheides, so that considerable lengths of the stem then contain a continuous hollow cylinder of lignified tissue. The bundles of the stem are collateral, the xylem occupying the axial, the phloem the peripheral side; and the structure is the same in Botrychium (Lunaria). Those of the leaf-stalk are four to eight in number, arranged in a circle and separated by fundamental tissue. In Ophioglossum they are collateral, the axial portion consisting of narrow reticulate tracheides, the peripheral portion of a broad band of phloem, containing sieve-tubes; while in Botrychium they are concentric, consisting of a broad band of scalariform or reticulate tracheides surrounded by a thick layer of phloem. There is no bundle-sheath in Ophioglossum, and only a rudimentary one in Botrychium.

The *roots* of Ophioglossaceæ are remarkable from the slight development of the root-cap, and the absence of root-hairs. They spring from the short stem in the midst of the leaf-insertions, and rarely branch, then always monopodially. Like the stem, they originate from a single pyramidal apical cell. The roots of Ophioglossum give rise to adventitious buds.

The *leaves* are always very few in number, often only one on each stem, and the number is uniform in the same species. They are remarkable for the slowness of their growth, which extends, in Botrychium Lunaria, over five years, the leaf only rising above the surface of the

soil at the commencement of the fifth year: where there are several leaves, the points of origin of those of successive years have a spiral phyllotaxis. They are quite simple and entire, simply pinnate, or twice or thrice pinnate. There is always a long petiole, which is furnished with a ligular or sheath-like outgrowth on each side; and the coalescence of these appendages forms a hollow chamber within which the leaf is developed, similarly to what takes place in Marattiaceæ. They are never circinate in vernation. They are of coriaceous texture, and are always quite glabrous, and possess a well defined epiderm furnished with stomates on both surfaces, and in immediate contact with the mesophyll, without any intermediate hypodermal layers. The mesophyll is large-celled and spongy, with large intercellular spaces. The 'vascular' bundles are but feebly developed; they anastomose in Ophioglossum, but only dichotomise in Botrychium and Helminthostachys. In most species all the leaves are fertile; but in Rhizoglossum (Presl) (a section of Ophioglossum) there are both barren and fertile leaves. The leaf divides at an early period into two branches—an outer branch which is sterile, and which alone develops chlorophyllous parenchyme; and an inner fertile branch, the *sporophyll*, which springs either from the base or middle of the lamina or from the leaf-stalk. This branch never has any green parenchyme except in Helminthostachys.



FIG. 75.—*Botrychium Lunaria.* Portion of sporophyll with open sporanges (magnified). (After Luerssen.)



FIG. 76. *Ophioglossum vulgatum.* Portion of sporophyll with closed sporanges, *s*; *g*, 'vascular' bundle (×10).

The *sporanges* resemble in their origin and mode of formation those of Marattiaceæ. They are not formed from a single cell, but from a group of cells in the substance of the sporophyll which are differentiated from the surrounding tissue. The terminal cell of the axial row beneath the epiderm is the archespore from which all the spores are formed; it is surrounded by the layers of mantle-cells constituting the tapete, which are formed out of the epidermal cell immediately above the archespore, and which ultimately disappear. The wall of the sporange, consisting of several layers of cells, is developed from the epiderm, and contains stomates. Strasburger regards each sporange as corresponding homologically to an entire sorus in the Filices—being, in fact, a meta-

morphosed portion of a leaf. They are buried, in an early stage, in parenchyme, which ultimately becomes entirely absorbed, and which is traversed by 'vascular' bundles anastomosing into long meshes. The sporanges require an entire year for their complete development. They have no annulus, and dehisce vertically from unequal tension of the epidermal and hypodermal layers of cells. In the formation of the spores each mother-cell divides into four 'special mother-cells,' with very thin cell-walls ; the protoplasm in each of these becomes invested with a new and firmer cell-wall, and the spores are thus completely formed, and are ultimately set free by the absorption of the walls both of the special mother-cells and of the original spore-mother-cells. The spores are nearly cubical ; the exospore is strongly cuticularised, and is furnished with prominent knobs and ridge.

Vegetative propagation is known to take place only by means of adventitious buds produced on the roots.

The Ophioglossaceæ include only a very small number of species, comprised in the genera Ophioglossum (L.), Botrychium (Sw.), and Helminthostachys (Kaulf.), spread over the whole globe. They are small plants, dying down each year, only a few species attaining the height of more than a foot, with a single or only a very few coriaceous leaves, and a conspicuous 'fructification,' which is simple and spicate in Ophioglossum and Helminthostachys, branched and paniculate in Botrychium. They are of no known economic value. They are represented in our English flora by the 'adder's-tongue' (Ophioglossum vulgatum, L.) and 'moonwort' (Botrychium Lunaria, Sw.).

LITERATURE.

Mettenius—Filices Hort. Bot. Lipsiensis, 1856.
Hofmeister—Abhandl. Sächs. Gesell. Wissenschaften, 1857.
Holle—Bot. Zeit., 1875, pp. 241 *et seq.*
Prantl—Ber. Deutsch. Bot. Gesell., 1883, pp. 155 and 348 ; and Jahrb. Bot. Gart. Berlin, 1884.

Class VI.—Equisetaceæ.

The Equisetaceæ or Horsetails are a very small group of plants, consisting only of a small number of species, arranged in a single genus, of remarkably uniform and peculiar habit. The aërial stems, which are invariably erect or ascending, arise from a creeping underground rhizome, and are characterised by their perfect multilateral symmetry and verticillate branching. The ascending stem is always elongated and slender, fluted with longitudinal furrows and ridges, and is remarkable for the

FIG. 77.—*Equisetum sylvaticum* L. *a*, non-chlorophyllous fertile branch : *b*, chlorophyllous fertile branch ; *c*, barren branch (reduced).

tendency displayed by the epidermal cells to deposit silica in their cell-walls. It is always divided into very distinct nodes and internodes, and is furnished at the nodes with modified foliar organs of a membranous character, the *leaf-sheaths*, the margin of which is split into a number of teeth. The tissue of the internodes is permeated by large intercellular air-canals. At the nodes the stem (except in some species the fertile stem) gives out a whorl of symmetrically arranged branches, which almost precisely resemble the main stem except in their smaller size and simpler structure, consisting of internodes and nodes furnished with leaf-sheaths; but these secondary branches do not usually again branch. The stem and root increase in length by means of a single large pyramidal apical cell, which produces three rows of segments. The 'vascular' bundles of the stem are but feebly developed, and contain but little xylem. Both stem and branches perform the function of leaves, contain chlorophyll, and are provided with stomates. The sporanges are never borne on the branches or leaves, but are collected into spicate or catkin-like 'fructifications,' borne at the extremity either of ordinary vegetative stems, or of special fertile stems which resemble the ordinary stems in structure, but contain no chlorophyll and do not branch. The sporanges spring from the inner side of the *peltate scales* of which these 'fructifications' are composed; they dehisce by a longitudinal fissure, but have no annulus like those of ferns. The spores are distinguished from those of all other Vascular Cryptogams by being enclosed in four distinct coats, the outermost of which ultimately breaks up into four ribbon-shaped strips, and detaches itself from the spore except at the point of junction of these strips, which are termed *elaters*. The elaters are remarkably hygroscopic, absorbing or giving off water with every change in the moisture of the surrounding atmosphere. In consequence of this they are constantly altering their shape, and imparting a somewhat rapid motion to the spores, thus assisting in their dissemination. On germinating the spore gives rise to a strap-shaped prothallium, which has an independent power of growth, and is usually dicœcious. The antherids and archegones differ in no essential point from those of other Vascular Cryptogams. Male and female prothallia are generally produced in close proximity to one another, so that impregnation is readily effected through the agency of moisture.

The *oophyte* generation of Equisetum (L.) springs directly from the spore, which contains chlorophyll. On germinating the spore throws off its outer coats, and changes its form from spherical to pear-shaped. The contents, still clothed in the thin endospore, then divide by a wall, the direction of which is not constant, into two cells of unequal size; according to Stahl the direction of this division depends on the direction of the rays of light. From the smaller of these two cells the chlorophyll

rapidly disappears entirely ; it undergoes no further division, but elongates rapidly into a long hyaline rhizoid. The larger of the two primary cells, which still contains abundance of chlorophyll, divides further by walls, at first in two directions only, into a multicellular plate which increases rapidly by apical growth, and soon branches in one plane. A difference is now set up between the development of the *male* and *female prothallia*. The former remain comparatively small and narrow, and the cell-division continues in the two directions only ; they consist, therefore, permanently of only a single layer of cells, and display but little lobing. Their colour is yellowish green. The female prothallia, on the other hand, grow to a considerably larger size, as much as half an inch in length, are of a deeper green colour, and at an early period form a number of lobes at their anterior portion, which consist of masses of merismatic tissue, cell-division taking place in the tangential as well as the other two directions ; they branch also in the same plane much more abundantly than the male prothallia. The formation of female or male prothallia appears to depend

FIG. 78.—*A*. male prothallium of *Equisetum arvense* L.; *a*, antherids (× 200). (After Hofmeister.) *B*—*E*, antherozoids of *E. maximum* Lam. in different stages of development (× 1200). (After Schacht.)

FIG. 79. Vertical section of lobe of female prothallium of *E. arvense*. *a*, archegones; *h*, rhizoids (× 600). (After Goebel.)

on the greater or less supply of nutriment. The antherids make their appearance about five weeks after germination, the archegones not till much later.

The *antherids* arise at the extremity or margin of the male prothallia. They are first separated off as papillæ by a tangential wall; further divisions then arise, by which they are divided into a large central cell and a single layer of much smaller peripheral 'mantle-cells.' The contents of the central cell then divide into the mother-cells of the *antherozoids*. These escape, still enclosed in the delicate wall of the 'special mother-cell,' by the separation from one another, through the action of water, of the apical cells of the mantle-layer; the expulsion often takes place with considerable force, and is due to the swelling up of the walls of the mother-cells; sometimes they emerge still all united together into a ball. The antherozoids are much more numerous than in ferns; probably several hundreds are formed in each antherid; they are also much larger, being the largest in any class of Cryptogams. Each antherozoid (see fig. 78) is a thread of protoplasm, gradually narrowing from the posterior to the anterior end, where it is coiled spirally, and bears a tuft of very long delicate vibratile cilia. The posterior portion is widened into a thin membranous fin-like expansion, by the undulations of which its motion through the water is greatly assisted, and may continue for many hours. To the posterior portion of the antherozoid is also frequently attached for a time a minute bladder containing starch, which is regarded by some as the wall of the special mother-cell, by others as a vesicle contained in it. The body of the antherozoid appears to be formed from the nucleus of the mother-cell, its cilia from the cell-protoplasm.



FIG. 80.—Portion of female prothallium of *E. sylvaticum*, with projecting archegone, *ar*. (After Buchtien, greatly magnified.)

The *archegones* are formed on the under (shaded) side of the thick lobed portions of the female prothallia, a lobe being usually situated immediately beneath an archegone, and assisting in its impregnation by retaining water. By the continued growth of the subjacent tissue they are ultimately pushed on to the upper surface, and hence the direction of growth of the archegone is the opposite of that of ferns. Otherwise

the mode of development and structure of the archegones of Equisetum differ in no essential respect from those of other Vascular Cryptogams. The basal cell is wanting, and the neck-canal-cell does not extend the whole length of the neck. The lower portion of the neck and the venter remain completely imbedded in the tissue of the prothallium, while the outermost or stigmatic cells of the neck lengthen greatly, and ultimately bend outwards, giving to the archegone, when ready for impregnation, the appearance of a miniature four-armed anchor.

Although the prothallium of Equisetum is normally dicœcious, it is not very uncommon to find a few archegones on male, and a few antherids on female prothallia. The abnormal sexual organs then generally make their appearance later than the normal ones.

The mode of formation of the *embryo* from the impregnated oosperm is essentially the same as in other classes of Vascular Cryptogams. The first division-wall is vertical to the axis of the archegone, and therefore parallel to the surface of the prothallium ; by subsequent walls a division takes place into octants. Of the four quadrants of the upper half, one gives birth to the rudimentary *stem*, with a triangular pyramidal apical cell, while from the three others proceed two *cotyledons*, which at a very early period unite in their growth with the first leaf which proceeds from the apex of the stem. From the lower quadrants of the octant are formed the *foot* and the first *root*. The first few stems of the sporophyte generation are successively thicker, and with a larger number of teeth in their leaf-sheaths, until ultimately mature stems are produced from perennial underground rhizomes.

The *stem* of Equisetum always consists of very distinct more or less elongated internodes, which are hollow, but are closed above and below at the nodes by transverse septa or *diaphragms*. The cortex of each internode is continued upwards above its upper node into a *leaf-sheath*, which embraces the base of the next internode above, and is split at its margin into *teeth*, varying from three or four to a considerable number. From each tooth of the leaf-sheath a 'vascular' bundle descends vertically as far as the next node. The teeth at each node always alternate in position with those of the leaf-sheaths belonging to the nodes next above and next below it, and each descending bundle branches at the node into two short diverging arms, each with its own xylem and phloem, by means of which it unites with the two adjacent bundles of the next internode below, where they descend into it from the sheath-teeth in which they originate. In addition to the large *central cavity* in the axis of each internode, the cortical tissue contains a number of much smaller cavities running vertically through the internode, the *lacunæ* or *vallecular canals*, which alternate with the 'vascular' bundles, and are there-

fore intermediate between the sheath-teeth ; these canals are of lysigenous origin—that is, they result from the disappearance of masses of cells. Each bundle also contains a longitudinal air-cavity, or *carinal canal* (sometimes called the 'essential air-cavity'). The outline of the stem always shows a number of alternate *ridges* and *furrows* ; the ridges correspond to (or are on the same radii as) the 'vascular' bundles, the furrows to the cortical lacunæ. This general description applies equally to the primary vegetative stems, the fertile unbranched stems, and the underground rhizomes ; the branches of the barren stems have no axial cavity or cortical lacunæ. In those species where the fertile differ in structure from the barren shoots, this appears to result from an arrest of development of the latter. The difference consists essentially in the absence of chlorophyll, the suppression of branching, and the absence of stomates, as well as in the greater development of the leaf-sheaths. It is possible to induce artificially the fertile shoots of E. arvense (L.) to put out green branches from the lower internodes, chlorophyll being also formed in the main stem.



FIG. 81.—*E. maximum*. *A*, portion of stem (natural size). *i*, *i'*, internodes ; *h*, central cavity ; *l*, cortical lacunæ ; *S*, leaf-sheath ; *a*, *a'*, *a''*, branches. *B*, longitudinal section of rhizome (×2). *K*, transverse diaphragm ; *h*, *h*, cavities ; *g*, 'vascular' bundle ; *l*, cortical lacunæ ; *S*, leaf-sheath. *C*, transverse section of rhizome. *D*, union of 'vascular' bundles of two internodes, *i*, *i* ; *K*, node. (*B*—*D*, diagrammatic.)

The branches always spring from within the leaf-sheath at its base, each branch in the space between two teeth ; they therefore correspond in number to the 'vascular' bundles of the stem, and are always arranged in a whorl. The same description applies to the roots. The number of leaf-teeth and bundles is always smaller on the secondary than on the primary axes, and these, as a rule, do not again produce branches. In some species even the primary stem never branches. The whorl in which the branches stand is, however, not a true, but a false whorl—that is, the phyllotaxis originally shows a regular spiral one-third divergence ; but by subsequent unequal growth the insertions

come ultimately to stand on a level. On the rhizomes the ridges and furrows of the outer surface are generally less well marked, and the axial cavity of the internode is sometimes wanting ; but the vallecular and carinal canals are always present, and play an important part in the diffusion of air through the tissue. The aërial stems, both barren and fertile, are usually completely formed in miniature during the preceding year within the underground bud, and their rapid growth after they appear above the soil is mainly due to the great elongation of the internodal cells. The ascending stem and all other aërial parts of the plant are always entirely destitute of hairs ; while the rhizomes and the underground leaf-sheaths are frequently covered with a felt of root-hairs.

The firmness and strength of the slender aërial stem are not due, as in ferns, to the 'vascular' bundles, but mainly to the siliceous epiderm and the firm hypodermal tissue. The epiderm, consisting of a single layer of elongated cells, is provided with *stomates* in the green, leafy, aërial stem, but not usually in the colourless fertile stem, or in the rhizomes. In most species of Equisetum the stomates lie in one or more longitudinal rows in the furrows of the stem only ; but in E. arvense, according to Miss E. A. Southworth ('American Naturalist,' 1884, p. 1041), also on the ridges. Stomates also occur on the leaf-sheaths. The stomates either have their orifice on a level with the surface (*Equiseta phaneropora*) or considerably depressed below it (*Equiseta cryptopora*) ; in the latter case they frequently do not open directly into the surrounding air, but are situated in the hypodermal tissue, beneath a 'false stomate,' or pore in the epiderm. The stomates (fig. 82) differ from those of other classes of vascular plants in being formed of two pairs instead of a single pair of guard-cells. Strasburger (Beitr. zur Entwickelungsgesch. der Spaltöffnungen, in Pringsheim's Jahrb., vol. v., p. 297) terms the lower pair 'subsidiary cells' of the true stomate. All the cells of the epiderm, even the guard-cells of the stomates, have their outer walls or cuticle strongly silicified ; and these deposits of silica frequently project above the surface in the form of fine granules, bosses, rosettes, rings, transverse bands, teeth, or spines. On the guard-cells they usually have the form of ridges radiating from the orifice. Beneath the epiderm, except on the deciduous fertile stems, bundles or layers of firm thick-walled cells generally constitute a sclerenchymatous *hypodermal tissue*, which is especially developed in the elevated ridges of the aërial internodes. On the underground stems both epiderm and hypoderm frequently assume a beautiful brown-red colour. In addition to silica, analysis of the ash of Equisetaceæ (Dieulafait, 'Compt. Rend.,' vol. c., 1885, p. 284) shows the presence

of a large amount of sulphates, and the total absence of alkaline carbonates.

A large portion of the parenchymatous fundamental tissue is destitute of chlorophyll; it is only in the vegetative aërial shoots that there is any considerable development of chlorophyllous tissue, and then it is usually situated in the furrows beneath the stomates. The 'vascular' bundles of Equisetum are collateral, and are much less strongly lignified than those of ferns. They lie in a circle between the medulla and the cortex, between and somewhat within the cortical canals, and necessarily run parallel to one another. As will be seen from the description given above, each bundle is the result of the coalescence of two branches, one of which originates in the leaf-sheath, while the other develops in the



FIG. 82.—Stomate of *E. hyemale* L. (× 390). *A*, front view; *B*, transverse section of stem, showing side view; *C*, siliceous residuum after maceration. (After de Bary.)

internode itself, from above downwards. At the angle where the two arms meet, the formation of tracheides begins in each; the lower end of each bundle unites by two lateral branches with the two next bundles, one on each side, of the next lower internode; and the bundles are therefore of the description known as 'common.' Their course resembles more that of the bundles of most Dicotyledons and Conifers than that of ferns. Each bundle is traversed longitudinally on its axial side by a carinal canal, occupying the place of the first tracheides, which have become absorbed; right and left of this, in the mature bundle, are reticulate, annular, and scalariform tracheides; on the outside is the phloem-portion of the bundle, consisting of a few wide

sieve-tubes and narrow cambiform cells. In most species (*e.g.* E. arvense) a *general bundle-sheath*, or *pleromc-sheath*, consisting of a single layer of cells, encloses the entire circle of bundles, as in most Flowering Plants; while in others (E. limosum, L., and littorale, Kuhlw.) each individual bundle is enclosed in a separate *special bundle-sheath*, as in ferns. In the colourless fertile shoots they bend out into the pedicels of the peltate scales of the 'fructification,' as they do into the leaf-sheaths.

The growth of the stem takes place through the activity of a single large pyramidal *apical cell* with an inverted triangular base. There is no other group of plants which exhibits such a well-defined single apical cell or exclusively apical growth. Normally the terminal bud never branches, branching taking place solely by lateral buds produced at the nodes. It was formerly thought that the Equisetaceæ display the only known example of lateral branching being due to the formation of endogenous buds; but recent researches have shown (in E. arvense) that these lateral buds are not of endogenous origin, but originate from a single superficial cell of the growing point of the stem in the ordinary way. The segments resulting from the first divisions of the apical cell lie in three straight rows, and are arranged in a spiral divergence of one-third.

The *roots* are produced in whorls at the nodes of the underground stem, in direct connection with lateral buds, or, under favourable conditions, at the nodes of the aërial stems. They are furnished with a root-cap, increase by the segmentation of a single apical cell, and are penetrated by an axial 'vascular' bundle surrounded by a large air-cavity formed by the coalescence of intercellular spaces owing to the absorption of intermediate cells. The weak bundle, in which the tracheides are but feebly developed, is concentric, with the xylem in the centre. The secondary or lateral roots, which arise in acropetal succession on the primary root, differ in their origin from those of ferns and other Vascular Cryptogams. In these latter it is the innermost layer of cells belonging to the fundamental cortical tissue immediately surrounding the axial 'vascular' bundle that becomes differentiated into the bundle-sheath or endodermal layer, within which lies the pericambium of the bundle itself; and the lateral roots originate from the innermost layer of the cortex separated by the pericambium from the bundle. In the roots of Equisetaceæ the pericambium is wanting, and its place is to a certain extent supplied by the innermost cortical layer, from which the lateral roots spring, and therefore in close contact with the periphery of the axial bundle. The bundle-sheath itself is, in the Equisetaceæ, formed from the row of cortical cells next to the innermost row, and not from the innermost row itself, as in other Vascular Cryptogams.

The *sporanges* of the Equisetaceæ are collected into terminal spicate 'fructifications' of a cone-like or catkin-like character, resembling nothing else among existing Vascular Cryptogams. These are borne at the extremity either of the ordinary green vegetative stems, whether branched (E. palustre, L.) or unbranched (E. hyemale, L.), or of special fertile stems (E. arvense, pratense, Ehrh., maximum, Lam.), which are then always simple, even when the barren stems are branched, and are usually stouter, nearly or quite destitute of chlorophyll, and with much larger leaf-sheaths and coarser teeth. As already mentioned, these can be artificially converted into vegetative stems; and occasionally deciduous fructifications are borne at the extremity of the ordinary green branched stems, in species which normally produce special fertile stems (E. arvense). The sporanges are not, like those of typical Filices, trichomic or epidermal in their origin; their development closely resembles that in the Marattiaceæ. They are endogenous outgrowths of peculiarly metamorphosed leaves, the *peltate scales* or *sporophylls*, arranged, like the branches, in whorls. Intermediate between these and the uppermost whorl of ordinary leaf-sheaths there is (in E. maximum) a whorl of barren but more or less modified leaf-sheaths, forming a small annular girdle, the *involucre* or *annulus*. The whorls of sporangiferous scales, of which a number are formed above this involucre, make their first appearance as similar annular girdles, projecting but slightly from the stem, but gradually forming a hemispherical cushion. This cushion finally breaks up into a number of plates, the surface of which is parallel to that of the stem; and these, by mutual pressure, become polygonal and usually hexagonal; each plate or disc is attached to the stem by a slender pedicel at right angles both to its surface and to that of the stem. On the inner surface of these



FIG. 83.—*A*, upper part of fertile stem of *E. maximum* (natural size); *b*, leaf-sheath; *a*, annulus; *x*, sporophylls and their stalks. *B*, sporophylls (× 6); *sg*, sporanges. (After Goebel.)

discs are developed the sporanges, from five to ten on each disc, arising at first as small multicellular projections. The archespore is the terminal cell of a hypodermal row on the under side of the sporophyll, the sporogenous tissue resulting from its division. The mantle-cells are formed in the same way as in Ophioglossaceæ, but are not so sharply defined. The mother-cells of the spores are connected together in groups of fours or eights, and float freely in the fluid which fills the sporange. The mode of formation of the spores affords an exceedingly good illustration of the production, by free-cell-formation, of new cells within a mother-cell. When division is about to take place, the protoplasm first becomes perfectly clear, the nucleus disappears, and a number of granules arrange themselves in the form of a disc. The protoplasm then again becomes turbid, with the exception of two clear spots, one on each side of the disc, which are the rudiments of the fresh nuclei. These, however, after a time again disappear, and their place is taken by four smaller nuclei, each of which is surrounded by a number of the granules which formed a portion of the original disc. Round these nuclei the cell-protoplasm begins now to collect into four separate masses, which gradually become globular ; and these are the special mother-cells of the spores after each has become invested with a very delicate coat of cellulose. This process, which has a remarkable analogy to the formation of the pollen in Flowering Plants, especially in Coniferæ, does not vary in any essential point from that in the other orders of Vascular Cryptogams : but it has been followed out with the greatest minuteness and success in Equisetum (limosum, L.).

The mature sporange dehisces by a longitudinal fissure on its inner side facing the sporophyll. The mechanism of the bursting is similar to that of the anther of Flowering Plants, and results from the unequal contraction of lignified and of non-lignified portions of the wall, which is furnished with annular or spiral thickenings to its cell-walls. The various coats of the spores are formed while still within the mother-cell. The first formed is the outermost, a non-cuticularised coat capable of swelling, which becomes gradually detached, and finally splits into two bands, the *elaters*, which remain attached to the spore only at one point, in the centre of each, where they meet, while the distal ends of each are dilated into a flattened spathulate form. When the spore escapes from the sporange the four elaters are stretched out nearly straight ; when moistened they roll up, owing to their unequal lignification, covering up the spore almost entirely, as they did at first before becoming detached. The second coat is more or less cuticularised, and on germination also raises itself in folds from the innermost coat, which is closely applied to the contents of the spore, and is again differentiated into an outer

granular cuticularised *exospore*, and an inner *endospore*, composed of unchanged cellulose. So strong is the hygroscopic property of the elaters, that, even if lightly breathed on, the spores of Equisetum are seen under the microscope to be in active motion, from constant changes in the humidity of the air. The spores contain chlorophyll, and, in consequence, retain their vitality only for a very few days, and germinate in a few hours after being placed in favourable conditions. In this respect they show a striking contrast to those of ferns.



FIG. 84.—Stages in the development of spore of *E. limosum* (× 800). 1 and *e*, outer coat from which the elaters are formed; 2, 3, inner coats. (After Goebel.)

FIG. 85.—Spore with elaters extended (magnified).

FIG. 86.—*E. limosum* L. Rhizome and tubers.

The only mode of vegetative propagation known in the Equisetaceæ is by the production of *tubers* on the rhizomes and on the underground portions of the erect stems; they are peculiarly modified internodes, filled with starch and other food-materials, and may remain dormant for years. The buds, especially those produced at the lower nodes of the erect stem, also have the power of retaining their vitality for a considerable period in a rudimentary condition; and, when they vegetate, develop into branches of great vigour. Tomaschek ('Oesterr. Bot. Zeitschr.,' 1881, p. 245) induced prothallia of Equisetum to hibernate by growing them

in a warm situation, in which condition they attained a large amount of independence, and propagated freely by budding.

The number of known species of Equisetum, commonly known as 'horsetails,' does not exceed 20 or 25 ; they are most numerous in the temperate regions, decreasing in number both towards the pole and the equator, and are very rare in the Southern Hemisphere. The stem is always very slender, and seldom exceeds two or three feet in height, though E. giganteum (L.) reaches 20 to 40 feet in the tropics, with a climbing habit. Most of the species prefer loose sandy or gravelly soil in damp situations ; several grow in marshes or standing water. The erect stems are mostly annual, but in a few species they endure for several years : while the rhizome is always perennial, and frequently attains great size both in depth and in lateral extension. The species are all remarkably similar in habit, differing chiefly in the presence or absence of special fertile stems, the position of the stomates, and the degree of branching ; but the classification of the species into two distinct groups of 'homophyadic' and 'heterophyadic' is not a natural one. Each species is also characterised by a special arrangement of the 'vascular' bundles, and of the air-cavities as seen in a transverse section of the stem. In external form, but not in internal structure, they call to mind Ephedra among Gymnosperms, and Casuarina among Angiosperms. The large amount of silex deposited in the epiderm renders several species, especially E. hyemale, useful for scouring purposes, and they are popularly known under the name of 'Dutch rushes.'

Literature.

Cramer—in Nägeli u. Cramer's Pflanzenphys. Unters., vol. iii., 1855.
Sanio—(Epiderm and Stomates) Linnæa, 1857–8, p. 385.
Duval-Jouve—Hist. Nat. des Equisetum, 1864.
Rees—(Apex of stem) Pringsheim's Jahrb. wiss. Bot., 1867, p. 209.
Milde—Monographia Equisetorum, 1867.
Pfitzer—(Bundle-sheath) Pringsheim's Jahrb. wiss. Bot., 1867, p. 297.
Famintzin—(Buds) Bull. Acad. Sc. St. Petersburg, vol. ix., 1876.
Campbell—(Prothallium) Amer. Natural., 1883, p. 10.
Leclerc du Sablon—(Sporange) Bull. Soc. Bot. France, 1884, p. 292 ; and Ann. Sc. Nat. (Bot.), vol. ii., 1885, p. 5.
Goebel—(Fertile shoots) Ber. Deutsch. Bot. Gesell., 1886, p. 184.
Buchtien—(Oophyte) Uhlworm and Haenlein's Biblioth. Bot., Heft 8, 1887.

Fossil Vascular Cryptogams.

Fossil remains or impressions of plants are found in all the stratified geological formations from the Upper Silurian to the latest. Of the Thallophytes that must certainly have existed in the seas from which the oldest fossiliferous strata were deposited, the traces are, as might be expected, few and doubtful; and it is certain that many markings that have been claimed under this category do not belong to the vegetable kingdom at all. The remains of Vascular Cryptogams make their first appearance in the Upper Silurian, and are remarkably abundant in the Devonian and Carboniferous formations. During these periods the arborescent vegetation of the globe consisted entirely of Vascular Cryptogams and Gymnosperms, no remains that can be referred with certainty to Angiosperms being known earlier than the Permian formation. The structure and mode of reproduction in Vascular Cryptogams seem to have been remarkably uniform from the earliest times to the present. The remains found in the fossil state belong, of course, exclusively to the sporophyte generation; but these indicate not only that nearly every class of Vascular Cryptogams now in existence was represented in the Carboniferous period, but also that none of the primeval forms of vegetable life at present discovered presented characters differing very widely from existing types.

Fossil Rhizocarpeæ.

The fossil remains that can be referred, with any degree of certainty, to the Rhizocarpeæ are very scanty. A few leaves found here and there have been described by their discoverers, under the names Marsilidium (Schenk) and Sphenoglossum (Emm.), as representing genera nearly allied to Marsilea; and capsules presenting an external resemblance to the sporocarps of Pilularia and Marsilea have been found in the Eocene.

The Salviniaceæ are represented with much more certainty in the Miocene, impressions of leaves found in various beds belonging to that series being indistinguishable from those of Salvinia. More doubt attends the identification of fructifications referred to this order. Strasburger and Solms-Laubach think it possible that certain minute echinate bodies found in calcareous nodules in the Carboniferous, described under the names Traquairia (Carruth.), Zygosporites (Will.), and Sporocarpon (Will.), the first of which is regarded by its discoverer as a

Radiolarian Rhizopod, may be massulæ of Azolla. Sir W. Dawson (Bull. Chicago Acad. Sc., 1886, p. 105) refers organs of fructification obtained from the Devonian (=Erian) in Canada and the northern United States—and previously described, under the name Sporangites (Daws.), as sporangia of Lycopodiaceæ—to a genus nearly allied to Salvinia, which he calls Protosalvinia ; but, inasmuch as they are borne on Lepidodendron scales, this explanation is inadmissible. Sir W. Dawson believes the megaspores of Rhizocarpeæ to be the chief cause of the highly bituminous character of the shales in which these bodies are found.

Fossil Selaginellaceæ.

Remains of arborescent vegetation more or less nearly allied to the typical Selaginellaceæ of the present day occur in extraordinary abundance in the older fossiliferous strata. Of these the most abundant and best known families are the Lepidodendreæ and the Sigillarieæ.

Of the Lepidodendreæ the stems are known as *Lepidodendron*, and the fructification occasionally found in organic connection with the



Fig. 87.—*A*, *B*, *C*, portions of surface of stem of different species of *Lepidodendron* (natural size) ; *D*, single cushion (magnified). (After Solms-Laubach.)

branches, as *Lepidostrobus*. The fructification is distinctly heterosporous ; and although, in a large number of Lepidostrobus cones, microspores only have been detected, this is unquestionably either because the portion containing the megaspores has not been preserved, or possibly because the megasporanges and microsporanges may have been distributed in distinct fructifications—a degree of differentiation unknown in any existing form.

The remains of a large number of species of *Lepidodendron* occur in the coal measures. They were trees, with stems up to ninety feet in height and two feet in diameter, covered with the diamond-shaped scars of fallen leaves. These scars, together with a portion of the leaf-stalk remaining behind in the form of a cushion, occupied the whole surface of the stem. Wherever the internal structure has been preserved, a

central 'vascular' cylinder can be detected, consisting of scalariform tracheides. There is distinct evidence of a secondary growth in thickness. The branching was always dichotomous. The leaves were very similar to those of Lycopodium, and were penetrated by a single 'vascular' bundle.

The fructifications known as *Lepidostrobus* are cone-like structures, not unlike fir-cones in appearance, consisting of densely packed sporophylls. On the upper side of each leaf is a single sporange, often of considerable size. The cones themselves vary in size from that of a hazel-nut to one and a half feet in length. It is seldom that the remains are in a sufficiently perfect condition for the structure of the spores to



FIG. 88.—*A*, transverse section of cone of *Lepidostrobus Brownii* Schimp. : *B*, longitudinal section (after R. Brown); *C*, diagrammatic longitudinal section of portion of cone of *L. ornatus* Hook. (after Hooker); *D*, upper surface of sporophyll (*Lepidophyllum*). (All from Solms-Laubach.)

be made out with certainty ; but in several examples both kinds occur. Where one kind only has at present been detected, it is, in most cases, the microspore, the megaspores being probably in the lower part of the fructification, which has not been preserved or examined. In the megaspores the exospore has three ridges ; there are numerous spores in each sporange. The microspores of L. dabadianus (Schpr.) are connected together in groups of four ; while in L. Brownei (Schpr.) they are in threes.

More or less nearly allied to Lepidodendron are a number of other arborescent genera, among the more striking of which are Ulodendron (Stbg.), Bothrodendron (L. and H.), and Lepidophloios (Stbg.), all from the coal measures.

Although the genus *Sigillaria* is still placed by some writers among Gymnosperms, its true place is undoubtedly near to Lepidodendron in the order Selaginellaceæ; the structure of the stem presents no important difference from that of Lepidodendron, while the fructification known as *Sigillariostrobus* bears a remarkable resemblance to Lepidostrobus.



FIG. 89.—*A*, *B*, *C*, portions of surface of stem of different species of *Sigillaria*; *D*, *Leiodermaria*. (After Solms-Laubach.)

The remains of various species of *Sigillaria* occur in enormous quantities in the coal measures; and they constituted one of the predominant forms of vegetation of the period. The stems rivalled in height and thickness those of Lepidodendron, and were covered, like them, with the scars of fallen leaves in linear series. They were simple or dichotomously branched. The scars are circular, ovate, or hexagonal from mutual compression. In the section known as *Leiodermaria* the cushions which occur in other forms are wanting, and the scars stand out at a considerable distance from one another on the smooth surface of the stem. The leaves, which are occasionally found still attached to the branches, were narrow, linear, and sedge-like, up to as much as one and a half feet in length, with a projecting midrib. According to Van Tieghem the stem of Sigillaria differs from that of the Lepidodendreæ, and indeed from that of all other Vascular Cryptogams, in the leaf-trace bundles being 'diploxylous'—that is, in the central cylinder having an external secondary and centrifugal as well as an internal primary and centripetal xylem. Renault regards the *Rhytidolepidæ*, or Sigillariæ with stem exhibiting raised cushions as well as scars, as Cryptogamic; the *Leiodermarieæ*, or smooth-stemmed Sigillariæ, as Gymnospermic; but this view is not supported by a careful examination of the structure.

Sigillariostrobus, the fructification of Sigillaria, is extremely rare. It was a cone resembling Lepidostrobus, with the sporanges placed singly on the base of the sporophylls. The sporophylls are broadly lanceolate and apiculate. Only one kind of spore has at present been discovered, the megaspores, but it may be regarded as certain that these were associated with a second and smaller kind.

The fossils known as *Stigmaria* are the roots of Sigillaria, the two having been not unfrequently found in connection with one another. They occur in the Devonian and Carboniferous formations. Fragments have been found from twenty to thirty feet or more in length (S. ficoides, Brongn.), cylindrical and unbranched, or the branching always dichotomous, the two branches running in a nearly parallel direction. The surface is smooth, with numerous shallow saucer-like depressions, the scars of the rootlets, some of which are still very commonly found attached to the primary root. The Stigmariæ obviously lengthened



FIG. 90.—*Stigmaria ficoides* Brongn. with rootlets. (After Solms-Laubach.)

exclusively by apical growth. Transverse section shows a hollow 'vascular' cylinder, broken by meshes for the passage of the bundles of the rootlets, and consisting of scalariform tracheides with a central parenchymatous tissue or 'medulla.' In the rootlets the single central bundle consists of a few scalariform tracheides, which leave the cylinder as a triangular bundle, but become circular in the rootlets.

Under the term LYCOPODITES are included a number of fossil forms, the fructification of which is either entirely unknown, or is not in a sufficiently well-preserved state for definite determination. Some of the leafy stems ought possibly to be referred to Coniferæ; others, with leaves of one kind only, perhaps belong to Lycopodiaceæ; while others, with leaves of two different kinds, are Selaginellaceæ. From beds near the bottom of the Carboniferous series there is a species with thick club-shaped terminal fructification, bearing a striking resemblance to Lycopodium Phlegmaria (Lycopodites Stockii, Kidst.). *Ptilophyton*

(Daws.), from the Devonian and Carboniferous formations of Scotland and North America, should also be included here.

The only fossils that can be referred with any degree of certainty to the Isoëteæ are those comprised in the genus *Isoëtites* (Schmp.), from the Miocene, which is scarcely distinguishable from existing Isoëtes. More doubt rests on the true place of Solenites (L. and H.), from the Jurassic, which has been referred with equal probability to Gymnosperms.

Of fossil Psiloteæ the remains are few and uncertain. To this family has been referred *Psilophyton* (Daws.); but the fructification is very



FIG. 91.—Bases of stem of *Sigillaria*, with *Stigmaria* roots attached. (After Solms-Laubach.

aberrant from the existing Psiloteæ, consisting of a pair of pod-like capsules at the end of special branches.

FOSSIL FILICES.

The remains of ferns—or more commonly the impressions of the leaves—are found in all fossiliferous strata from the Devonian onwards. Great difficulty is presented in the classification of fossil ferns by the small fragments in which they are usually found, anything like an entire plant, or even a number of fronds attached to an aërial stem

or rhizome, being extremely rare. The great majority of species appear to have been herbaceous ; or at all events the stems of tree-ferns are not of very common occurrence, even in the coal measures. And although the fructification has frequently been met with, the vast majority of the leaves of which the remains or the impressions have been preserved are barren. The only available system of classification of the greater number of fossil ferns is based on the mode of venation, on which character a number of families have been founded by Brongniart ; but it is doubtful whether this has any great value as a natural system of classification. A form of heterophylly different from anything which occurs among existing ferns is found in a few species from the Carboniferous formation, where, in addition to the normal pinnæ of the frond, themselves again pinnate, imperfect pinnæ of much smaller size and simpler structure are intercalated between them. These imperfect pinnæ, known as *aphlebiæ*, were described as distinct species before their true character was known. Thus Rhacophyllum adnascens (L. and H.) is the aphlebia of Sphenopteris crenata (L. and H.) ; while various so-called species of Cyclopteris are abnormal pinnæ springing from the rachis below the normal pinnæ of Neuropteris. On the whole, the leaves of ferns belonging to the Carboniferous period bear a striking resemblance to those of our own day ; in many cases they might belong to living genera.

FIG. 92.—*Aphlebia*, from the Carboniferous formation. 1, *Sphenopteris crenata* L. and H. ; 2, 3, *Rhacophyllum adnascens* L. and H. (After Schimper.)

The remains of the fructification of fossil ferns that have come down to us lead us to believe that the existing orders of Filices may have been represented in the earlier geological periods ; and none have as yet been

discovered that cannot be referred to some existing type. The prevalent forms appear to have been the Polypodiaceæ, Hymenophyllaceæ, and Marattiaceæ ; this last order having been apparently much more widely distributed and more abundant in the earlier periods than it is now.

The HYMENOPHYLLACEÆ may possibly have been one of the earliest differentiated types. In Palæopteris hibernica (Schmp.) (Cyclopteris

FIG. 93.—A, frond of *Palæopteris hibernica* Schmp. (restored) (÷6) ; B, pinnule (somewhat mag.) ; C, fertile pinna (nat. size) ; D, two cup-shaped indusia attached to the filiform midrib (mag.) ; E, sporanges of a hymenophyllaceous fern from the coal measures (mag.). (After Carruthers.)

hibernica, Forbes), specimens have been found in which all the lower pinnæ are fertile. The pinnule was reduced to a midrib supporting the slender stalks of the bilabiate cup-shaped indusia ; and the stalk is continued into the indusium, to which the sessile sporanges are attached. The texture of the frond was not membranaceous, like that of most existing Hymenophyllaceæ, but was more like that of Loxsoma. On the rachis between the pinnæ are seated single large decurrent pinnules.

The edges of the pinnules are slightly serrate from the numerous dichotomising veins ; the lower part of the stipe is clothed with scales. Sporanges with the characteristic oblique annulus of the Hymenophyllaceæ have also been found in the coal measures by Carruthers ('Geol. Mag.,' Feb. 1872).

FIG. 94.—Fructifications of fossil Marattiaceæ. *A*, *Seftenbergia ophidermatica*; *B*, *Haulea Miltoni*; *C*, *Oligocarpia Lindsæoides*; *D*, *Scolecopteris polymorpha*; *E*, *Asterotheca Sternbergii*. (After Solms-Laubach.)

Remains which can be referred with certainty to the MARATTIACEÆ are not unfrequent. In Scolecopteris (Stur) we have a true synange ; the separate sporanges, arranged on a common elevated receptacle, are linear-ovate with a long free apex, and open by a fissure on the inner side without any trace of an annulus. In Asterotheca (Presl) the circular sorus usually consists of six exannulate sporanges closely adnate to one another, the sori are sessile, and are arranged in a single row on each side of the midrib of the pinna. In Renaultia (Stur) a group of cells occurs in the outer wall of the sporange similar to that in Angiopteris, which may be the rudiment of an annulus. Seftenbergia (Cord.) presents important differences. The sporanges are not collected into sori, but are scattered along the veins of the third order ; each sporange has at its apex a cap-like annulus. It appears to be a connecting link between the Marattiaceæ and Schizæaceæ. Other types of Marattiaceæ are presented by Danæites (Göpp.) and Botryopteris (Ren.).

The remaining types of ferns of the Devonian and Carboniferous, and especially those of more recent periods, present the greatest resem-

blance to POLYPODIACEÆ and CYATHEACEÆ among existing forms. No fructification resembling that of Osmundaceæ has at present been discovered; but *Osmundites* (Carruth.), from the Lower Eocene, has been referred to that order by its discoverer from the peculiarities of the structure of the stem. The Ophioglossaceæ are still unrepresented in palæophytology.

The internal structure of the stem and leaf-stalk of most fossil ferns,



FIG. 95.—Section of *Stemmatopteris* Cord. invested with roots=*Psaronius* Cord. (From a specimen in the British Museum.)

where this can be determined, differs in no important respect from that of living forms. We find the same interrupted ring of 'vascular' bundles, which may be either concentric or collateral, the xylem consisting largely of scalariform tracheides, the same layers of sclerenchyme both in connection with the bundles and beneath the epiderm; evident indications of gum-passages have even been detected. But though this is by far the most common type of structure, a second is displayed in a few rare examples, in which the arrangement of the 'vascular' elements

may be compared to that in the stem of Monocotyledons, as the ordinary arrangement may be to that in the first year in the stem of Dicotyledons. In Stemmatopteris (Cord.) (including Psaronius, Cord., and Zippea, Cord.), from the Bath coal-field, the circumference of the stem is composed of a continuous envelope of sclerenchymatous tissue, within which are perpendicular tracts of 'vascular' tissue not penetrated by meshes. Between these tracts the leaves were given off in perpendicular series, the large single leaf-bundles coming right out from the central parenchyme, in which they existed as well-formed bundles, filling up more or less completely the central cavity (see fig. 95). There is therefore no closed cylinder with central 'medulla' as in ordinary ferns. By some authors it has been proposed to establish the fern-stems which display this character as a separate group under the name Psaronieæ, but there is every reason to identify the stem of Stemmatopteris insignis (Cord.) with the fronds of Pecopteris arborescens (Schl.), which bear fructification indistinguishable from that of Cyathea; and, this character being the more important, the genus must be placed under Cyatheaceæ.

Fossil Equisetaceæ.

Remains of the genus *Equisetites*, evidently very nearly allied to Equisetum, if not identical with it, are found in greater or less abundance in various strata from the Carboniferous to the Tertiary, attaining their maximum development in the Trias. The stems of these fossil horsetails are from one and a half to six inches in diameter, and may have attained a height of from twenty-five to thirty-five feet. They are cylindrical, and are marked with alternate ridges and furrows. At the nodes are tubular leaf-sheaths split at the margin into numerous short teeth, each of which terminates in an elongated bristle; in some species the number of these teeth appears to have amounted to as many as one hundred. The nodal diaphragms are clearly seen in E. arenaceus (Brongn.), the remains of which occur in extraordinary abundance in the Upper Trias; and, in some species at least, the furrows and ridges of each internode are alternate respectively with those of the internodes next above and next below. Remains of rhizomes have been found closely resembling in structure those of Equisetum. Nearly allied to Equisetites are the genera Schizoneura (Schmp.) and Phyllotheca (Brongn.); the latter differing from the type in its spreading sheath-teeth, and in the ridges and furrows of adjacent internodes not being alternate. The fructification of Equisetites has only been found in a very imperfect condition. That of Phyllotheca bears a close resemblance to the cone-

like sporangiferous spikes of Equisetum. It contained spores of one kind only.

The group of CALAMARIEÆ—including the stems known as *Calamites* and *Calamodendron*, and the fruit known as *Calamostachys*—have been separated by some authorities from the Equisetaceæ on the ground of their alleged heterospory, but without sufficient warrant from the facts of their structure as actually observed.

The remains of *Calamites* occur in immense quantities in the Carboniferous strata ; apparently they constituted one of the most important features of the vegetation of that period, disappearing after the Permian. The stems were frequently of gigantic dimensions compared with our existing Equisetums, attaining a height of thirty feet and a diameter of four inches or more. They consist of a hollow central cavity, with a cylinder of tracheides in wedge-shaped bundles, separated at their origin by parenchyme, and alternating at the nodes, where there is a diaphragm or 'phragma.' The leaves, which spring in whorls from each node, do not, as in Equisetites, coalesce laterally into sheaths surrounding the stem. They are narrow and acicular, with a single prominent midrib. At the nodes are occasionally seen saucer-shaped depressions, the scars of the lateral branches, which are sometimes found attached to the primary axis.

FIG. 96.—*A*, *Phyllotheca equisetiformis*; *B*, fructification of *Phyllotheca*. (After Solms-Laubach.)

The growth of the stem is characterised by a considerable secondary increase in thickness ; and, since this phenomenon was formerly unknown among living Vascular Cryptogams, it has induced some authorities to transfer those examples where it occurs, under the name of Calamodendreæ, to Gymnosperms ; but this has been rendered unnecessary from the fact that a secondary growth in thickness occurs also in Lepidodendron and Sigillaria, as well as in Isoëtes ; and is further contradicted by the fact that fructification of an evidently cryptogamic character has been found in organic connection with stems

which must be referred to the same group. To the same family as Calamites belong probably Astromyelon (Will.), and at least some species of Arthropitys (Göpp.). The degree of identity in structure of *Calamodendron* with Calamites is a point on which the best authorities are not yet in agreement.

To the genus *Calamitina* (Weiss) (Asterophyllites, Ren., Calamocladus, Schmp.) belong a number of calamite stems found with the leaves still in connection with them. These are of very peculiar form, consisting of an ovate-lanceolate basal portion, thickened and marked by a central furrow, and a narrowly-lanceolate acuminate apical portion ; the basal



FIG. 97.—Stems of Calamites. (After Weiss.)

portion alone being very frequently preserved. The leaves do not coalesce laterally into sheaths ; on falling off they leave behind whorls of round or ovate scars. Bornia (Brongn.) (Archæocalamites, Stur) is an older fossil occurring in the Devonian formation, differing from the more recent forms in the broad flat longitudinal ribs on the stem not alternating in adjacent internodes.

In *Annularia* (Brongn.), which occurs only in the Carboniferous formation, the leaves are linear-lanceolate, and are penetrated by a single 'vascular' bundle ; those of each whorl are united laterally in their basal portion into a shallow saucer-shaped cup, through which the stem

passes, and from the margin of which spring the linear-lanceolate free portions of the leaves. Annularia has been regarded by some writers as an herbaceous aquatic plant; but there is little doubt that it is the branches and foliage of Calamites.

The fructification of the Calamarieæ, described under the name *Calamostachys*—with which must be identified Volkmannia (Stbg.) and

FIG. 98.—Leaves of *Calamitina*. (After Weiss.)

FIG. 99.—*Archæocalamites radiatus* Stur. (After Stur.)

Bruckmannia (Stbg.)—has not unfrequently been found in organic connection with the stem. Each cylindrical cone-like fructification consists of a number of whorls of sporophylls, but differs from that of Equisetum or of Equisetites in the fertile whorls alternating, in each spike, with barren whorls consisting of a large number of lanceolate acute leaves, free or more or less connate at the base; the free portions completely cover the next upper fertile whorl and the base of the barren whorl

above that ; thus giving a remarkable superficial resemblance to a fir-cone. The sporophylls of each whorl are not united with one another ; they resemble those of Equisetum in their peltate form, and each bears on its under side four sporanges. It is very rarely that the spores are preserved in sufficient perfection for their structure to be made out with certainty. The statement that they are of two kinds, megaspores and microspores, rests on insufficient evidence. Carruthers believes that he has detected in a few cases an outer separable membrane which would



FIG. 100.—*A*, Fructification of *Cingularia typica*; *B*, portion of a barren and of a fertile whorl *C*, upper surface of fertile whorl. (After Weiss.)

unroll itself in the form of elaters. In Palæostachys (Weiss) the sporophylls stand in the axils of the barren leaves. The fructification of *Cingularia* (Weiss) has a very remarkable appearance, from the alternate barren and fertile whorls standing out nearly at right angles to the axis ; the leaves of the barren whorls are connate for about half their length ; the sporophylls are also united in each whorl into a horizontal plate, divided at the margin into truncate lobes ; from the under side of this plate the sporanges hang vertically in radial rows. The remains of Cingularia fructifications are found in large numbers in the coal

measures. Very little is known of their stem, which appears, however, to have resembled that of the Calamarieæ.

More doubt rests on the affinity with the Equisetaceæ of the group of SPHENOPHYLLEÆ. The remains of *Sphenophyllum* (Brongn.) are found in abundance in the Carboniferous formation, but do not come down to more recent times than the Lower Permian. The stem is divided into distinct nodes and internodes, the latter usually marked with conspicuous ridges and furrows, which are not alternate in adjacent internodes, but pass continuously through the nodes. At the swollen nodes are whorls of leaves, with occasional axillary branches. Each whorl appears to consist always of six leaves or of some multiple of six. They are sessile, and obcuneate from a narrow base, sometimes denticulate and bifid at the apex, but are not in any degree connate. Each leaf contains a number of simple or dichotomous 'vascular' bundles. In the centre of the stem is a triangular bundle composed of scalariform tracheides, to which some authorities add spiral tracheides and others with bordered pits; the bundle passes through the node without material change. This is often surrounded by some layers of secondary wood; the greater part of the stem, on transverse section, is occupied by a small-celled parenchyme.



FIG. 101.—1, 2, Fructification and branch of *Asterophyllites*; 3, 4, fructification and branch of *Annularia*; 5, 6, fructification and branch of *Sphenophyllum*. (After Carruthers.)

The fructification of Sphenophyllum consists of cylindrical cone-like spikes resembling those of Calamites. It is composed of whorls

of sporophylls without any intermediate barren whorls ; the separate leaves are often saccate, the narrow apices ascending and covering in an imbricate fashion the bases of the next upper whorl. The position of the sporanges differs materially from that of the other Equisetaceæ. They are comparatively large bodies, lenticular, from 1 to 2·5 mm. in diameter, solitary and sessile in the saccate hollow on the upper surface of the base of the sporophyll. The solid remains have not yet been found in a sufficiently perfect condition for the structure of the spores to be determined with any degree of certainty, but no sufficient evidence of heterospory has been presented.

From the form and position of the sporanges the Sphenophylleæ are placed by some writers under Selaginellaceæ ; but in the general appearance and structure of the vegetative organs they approach so nearly to the Calamarieæ that it seems best at present to place them here until we are better acquainted with the details of the fructification. Stur has recently described and figured specimens with leaves of Asterophyllites at the base, Sphenophyllum-leaves higher up, and terminating in a fructification.

Literature of Fossil Vascular Cryptogams.

Sternberg—Flora der Vorwelt, 1821–1838.

Brongniart—Hist. des Végétaux Fossiles, 1820-1840; (Sigillaria, Stigmaria, and Lepidodendron) Arch. Mus. d'Hist. Nat., 1839.

Lindley and Hutton—Fossil Flora of Great Britain, 1833-1837.

Witham—Internal Structure of Fossil Vegetables, 1833.

Göppert—Systema Filicum Fossilium, 1826 ; Gattungen der fossilen Pflanzen, 1841.

Unger—Genera et Species Plantarum Fossilium, 1840.

Corda—Beitr. zur Flora der Vorwelt, 1845.

Hooker—Vegetation of Carboniferous Period, in Mem. Geol. Survey, 1847.

Brown—Triptosporites, in Linn. Soc. Trans., 1851.

Geinitz—Steinkohlenformation in Sachsen, 1855.

Ludwig—Calamiten-Früchte Palæontographia, 1861.

Goldenberg—Flora Saraepontana, 1862.

Binney—Fossil Carboniferous Plants, Palæont. Soc., 1868–1875.

Schimper—Paléontologie végétale, 1869–1874.

Weiss—Fossile Flora der Steinkohlenformation, &c., 1869 ; Beitr. zur fossilen Flora, 1876–1884.

Carruthers—(Ulodendron and Bothrodendron) Monthly Micr. Journ., 1870 ; (Lycopodiaceæ) 1869 ; (Calamites) Seemann's Journ. Bot., 1867.

Dawson—Fossil Plants of Upper Silurian and Devonian of Canada, 1871–1882 ; Geological Hist. of Plants, 1888.

De Saporta—Paléontologie Française, 1873.

Grand'Eury—Flora Carbonifère du Dpt. de la Loire, &c., 1877.

Renault—Cours de Bot. fossile, vols. i.–iii., 1881–1883.
Williamson—Organisation of Plants of the Coal Measures, Phil. Trans. 1871–1888; (Stigmaria) Palæont. Soc., 1887.
Kidston—(Lepidodendron, Sigillaria, &c.) Ann. and Mag. Nat. Hist., 1885.
Lesquereux—Coal Flora of Carboniferous Formation of Pennsylvania, &c., 1884.
Stur—Carbonflora der Schatzlarenschichter, 1885; Culmflora, 1877–1883.
Van Tieghem—Bull. Soc. Bot. France, 1883, p. 169.
Zeiller—Ann. Sc. Nat. (Bot.), 1884; (Sigillaria) do., 1884, p. 256.
Zittel—Handbuch der Palæontologie, 1879–1885.
Solms-Laubach—Einleitung in die Palæophytologie, 1887 (for which see very copious bibliography).
Schenk—Die fossilen Pflanzenreste, in Schenk's Handb. der Botanik, vol. iv., 1888.

SECOND SUBDIVISION.

MUSCINEÆ.

The line of demarcation between the Vascular Cryptogams and the plants immediately below them in the scale of organisation, the Muscineæ, is a very sharp one, and their genetic relationship to one another presents considerable difficulties. The lower type of structure is, however, chiefly manifested in the vegetative organs. The mode of sexual reproduction which occurs throughout this group corresponds in its most important features with that in Vascular Cryptogams ; and we have here also a division of the life-history of the plants into a *sporophyte* and an *oophyte* generation, a true alternation of generations, although the phenomenon differs in one important point from that which we have seen in Vascular Cryptogams, viz. in almost the whole of the vegetative system belonging to the oophyte instead of to the sporophyte generation. To this we have already seen an approach in Gymnogramme (p. 65). The vegetative system is invariably of small size, and almost entirely destitute of vascular bundles and of all other strengthening tissues. Within the group the boundary line is crossed between Cormophytes and Thallophytes ; and in the lower orders we entirely lose the differentiation of the vegetative organs into cauline and appendicular organs—in other words, into stem and leaves ; the entire vegetative system consisting of an undifferentiated *thallus*. The mature plant is almost invariably terrestrial in habit, and is attached to the substratum by *rhizoids*. The appendicular organs, when present, are minute leaves, which never contain true vascular bundles, and usually consist of only a single layer of cells. We find, however, the first stage towards the epidermal and fibrovascular structures characteristic of the leaves of vascular plants, in a distinct midrib and edging of elongated cells with somewhat thicker cell-walls overlapping one another at the extremities, and partially or altogether destitute of chlorophyll. In one group (Sphagnaceæ) the leaves are composed of cells of two different kinds, small cells containing chlorophyll interspersed among much larger empty cells. The leaves, being usually unilamellar, cannot, of course, be pro-

vided with stomates, though these are frequently present on the organs of propagation ; while one group of thalloid forms (Marchantiaceæ) possess stomates of remarkable and complicated structure. The vegetative propagation of the Muscineæ takes place in several ways : 1st, by *innovation*, *i.e.* by a process of renewal at the apex, while the oldest parts die off behind ; 2nd, by means of gemmæ, stolons, or detached buds ; and 3rd, by the non-sexual production of a thallus or protoneme, a process which will be described presently. The facility of these various modes of vegetative multiplication gives rise to the tufted or cæspitose habit of many species.

Notwithstanding the variety in the development of the vegetative structure, the sexual organs of Muscineæ are remarkably uniform in their main features. The male and female organs are termed respectively, as in Vascular Cryptogams, *antherids* and *archegones*. The mature antherid is a spherical, ellipsoidal, or club-shaped body, with a longer or shorter stalk, the outer layer of its cells forming an enclosing wall, while each of the small and numerous crowded cells in the interior develops an *antherozoid*. These bodies are spirally-coiled threads of protoplasm, thicker at the posterior end, and tapering to a fine point at the anterior end, where they are furnished with two long fine cilia, the vibrations of which set them in constant motion ; they are set free by the rupture of the wall of the antherid at its apex. The archegones, when in a condition capable of impregnation, are flask-shaped bodies bulging from a narrow base, and produced above into a long *neck*. The swollen or ventral portion, the *venter*, encloses one cell much larger than the rest, the *central cell*, from the larger and lower portion of which is developed, after its separation by a horizontal septum, the germ-cell or *oosphere*. Above this central cell is an axial row of cells termed the *canal-cells*, passing through the narrow portion or neck of the archegone, and continued as far as the apical cells, *stigmatic* cells, or lid-cells, which form what is called the *stigma*. These canal-cells are transformed before impregnation into mucilage, which finally swells up and forces apart the four stigmatic cells, an open canal being thus formed, through which the antherozoids reach the oosphere. Notwithstanding the general uniformity in structure of the sexual organs of the Muscineæ, their origin varies greatly. They may originate, in the thalloid forms, below the growing apex, from the superficial cells of the thallus, or on special metamorphosed branches ; in the foliose forms both antherids and archegones may be formed from the apical cell of the shoot, or from segments of it ; and in this case they may take the place of leaves, of lateral shoots, or even of hairs. According to Leitgeb, the order of evolution from the lower to the higher forms of

Muscineæ is indicated by the position of the sexual organs on the vegetative shoots ; as these organs approach nearer and nearer to the apex, the shoot gradually loses its vegetative character, and becomes differentiated into a special fertile branch.

The sexual organs are frequently surrounded by crowded and slightly modified foliar structures, the whole arrangement having then some resemblance to the flower of Phanerogams. Thus, in the thalloid forms, the antherids and archegones are commonly borne on umbrella-shaped outgrowths of the thallus which are not inaptly termed respectively male and female *inflorescences*. In the foliose forms they are frequently arranged at the extremities or laterally on branches, and closely surrounded by small leaves constituting the *perichæte* or perianth, reminding one of the bracts, or even of the calyx, of Flowering Plants, the whole structure forming an hermaphrodite, monœcious, or dicecious 'flower.' They are often accompanied by barren, hair-like cells, termed *paraphyses*.

The first result of the impregnation of the oosphere by an antherozoid is the formation of an ovoid *embryo* by repeated cell-divisions ; this continues to grow at its apex, and finally develops into the fructification here known distinctively as the *sporogone*, the ultimate form of which varies greatly in the different families. In its most perfect form the sporogone is differentiated externally into a slender stalk or *seta*, which penetrates into the base of the archegone, or even into the underlying tissue, and a spore-capsule, called indifferently the *sporange*, *theca*, or *urn*. Along with the spores the spore-capsule sometimes contains (in the Hepaticæ) elongated cells thickened by a single or double spiral band known as *elaters*, which assist in the dissemination of the spores. The mature sporange is, in the highest forms, surmounted by a cap or *calypter*, which becomes completely detached at its base, while the mature lower portion of the archegone encloses the base of the seta in the form of a sheath or *vagine* ; in the lower forms the spore-capsule always remains enclosed in the calypter. The *spores* of the Muscineæ are always formed in fours within the *spore-mother-cells*, which latter are produced within the cavity of the sporange by free cell-formation, from a special layer or layers of cells known as the *archespore*. When ripe they have a double cell-wall, the outermost layer or *exospore* being provided with small excrescences, and the inner layer or *endospore* bursting through it on germination. The contents consist of protoplasm, chlorophyll-grains, starch, and oil.

The thalloid or leafy plant (as the case may be) does not, as a rule, arise immediately from the germinating spore, but, in all the higher forms, only after the previous formation of a colourless confervoid or

filamentous structure, the prothallus or *protoneme*, on which the leafy plant containing chlorophyll arises as a lateral shoot. The Muscineæ present, therefore, an illustration of the phenomenon of alternation of generations ; the sexual generation which intervenes between germination and impregnation, or *oophyte*, consisting of the protoneme (when present), the leafy (or thalloid) plant, together with the sexual organs ; the non-sexual generation intervening between impregnation and germination, or *sporophyte*, consisting of the sporogone only with its spores.

The Muscineæ are divided into two well-marked families, the *Musci* or Mosses, and *Hepaticæ* or Liverworts. In the Musci the immediate result of the germination of the spore is always a protoneme consisting of branched rows of green or colourless cells, and often growing for a considerable time independently, even after it has produced leafy stems by lateral budding. The vegetative structure is always cormophytic, and consists of a filiform stem furnished with two, three, or four rows of leaves, not exhibiting any distinct bilateral structure, and branching monopodially, never dichotomously. The sporogone is only for a time enclosed in the calypter, which is usually eventually ruptured below, the lower portion developing into the *vagine*, while the upper part is elevated above the apex of the sporogone in the form of a cap. The spore-mother-cells are produced from one or more special layers of tissue within the sporange, the *archespore*, while the axial mass develops into a solid *columel*. The uppermost portion of the wall of the sporange forms a lid or *opercule*, which usually becomes detached from the lower portion, to which the term theca or *sporange* specially belongs, to allow the escape of the spores. The outermost layer of cells of the wall of the sporange is more or less completely differentiated into an epiderm, which is frequently penetrated by stomates. When the opercule is removed, the rim of the sporange is either quite smooth, when it is termed *gymnostomous*, or the edge is furnished with delicate hair-like appendages, constituting the *peristome*, arranged in a single row or frequently in two, when they are called respectively *teeth* and *cilia*, the former constituting the outer, the latter the inner row. The number of both teeth and cilia is always a multiple of four, or more correctly speaking, a 'power' of two. In the Hepaticæ the protoneme is either scantily developed or is altogether suppressed. The rest of the sexual generation consists either of a flat dichotomously branched thallus or thalloid stem, or of a slender stalk furnished with two or three rows of leaves. In the division into Foliose and Thalloid or Frondose forms, the Hepaticæ therefore present the transition from Cormophytes to Thallophytes. The mode of growth is always distinctly bilateral ; the thalloid forms cling

closely to the ground or to some other substratum ; and even in the foliose forms there is a well-marked tendency to the differentiation of an upper or dorsal and an under or ventral surface. The non-sexual generation or sporogone remains surrounded by the calypter until the spores are ripe ; the calypter is usually at length ruptured at the apex, and remains at the base of the sporogone as an open sheath, while the sporange projects above its apex to allow the escape of the spores. The mother-cells of the spores are either developed from the whole of the archespore, or, in most families, some of the cells develop into elaters ; there is no columel.

LITERATURE.

Hofmeister—On the Germination, Development, and Fructification of the Higher Cryptogamia, Ray. Soc., 1862.
Leclerc du Sablon., Ann. Sc. Nat., 1885, p. 126 ; and Bull. Soc. Bot. France, 1885, pp. 30 and 187.
Goebel—Die Muscineen, in Schenk's Handbuch der Botanik, vol. ii., 1882.

Class VII.—Musci.

The germinating spore of Mosses gives rise to a prothallium which is always in the typical families of a filamentous conferva-like character, and is hence known as the *protoneme*. On this is produced the leafy plant with differentiated stem and leaves by lateral budding. These together, therefore, constitute the sexual generation or oophyte, which terminates in the production of the sexual organs. From the fertilised oosphere proceeds the *sporogone*, which comprises in itself the non-sexual generation or sporophyte.

The protoneme first originates, in typical mosses (Bryaceæ), as a tubular bulging of the endospore or inner coat of the spore, divisions taking place in it in the transverse direction only. It has an unlimited power of apical growth, and often branches copiously, forming a dense felt of considerable size above or below the surface of the soil, in the former case densely filled with chlorophyll. The protoneme usually disappears altogether after the appearance on it of the leafy plant ; but in some cases, especially in the Phascaceæ, it remains vigorous even after the formation of the sporogone. In the Sphagnaceæ the protoneme consists of a flat plate of cells ; while in the Andreæaceæ cell-division begins to take place within the spore, the resulting prothallium consisting of a small cellular tissue. The buds which develop into the

leafy stems appear never or very seldom to arise at the apex of a primary filament of the protoneme, but always as lateral branches.

Mosses display a certain amount of differentiation of tissues. The apical cell of the stem is, except in Fissidens (Hedw.), a three-sided

FIG. 102.—*Catharinea* (*Atrichum*) *undulata* W. and M. (magnified). (After Schimper.)

pyramid. The primary meristem of the stem, situated beneath the growing point, which develops into the permanent tissue, usually becomes differentiated into an inner and a peripheral mass; the latter, although

not strongly defined, partaking of an epidermal character ; the bright red or yellow cell-walls are considerably thicker than those of the central thin-walled large-celled fundamental tissue. In some genera a further differentiation takes place of the axial portion of the central cylinder into a rudimentary 'vascular' bundle with thicker cell-walls ; similar rudimentary bundles being also formed in the pedicel of the sporogone. Both the partially lignified and the thin-walled cells have simple pits in their cell-walls ; these are found in all families of mosses, and are especially abundant in the midrib of the leaves. Some species of Sphagnum (L.) have rudimentary sieve-plates. The central bundle in the stem of Mnium, Polytrichum (L.), and other genera, has been shown by Haberlandt not to possess any of the strengthening functions of a true vascular bundle, but



FIG. 103.—*A*, germinating spore of *Funaria hygrometrica* L. (× 550); *s*, exospore ; *w*, rhizoid ; *v*, vacuole. *B*, portion of protoneme (× 90); *K*, rudiment of leafy axis ; *w*, rhizoid. (After Goebel.)

to be constructed for the purpose of the conduction of water. Its cells contain nothing but a watery fluid, without starch-grains, oil, or protoplasm. In genera which have no such central bundle, like Dicranum (Hedw.) and Leucobryum (Hpe.), the epidermal tissue of the stem and branches, with its perforated cells, forms a similar capillary apparatus. In the more highly developed mosses, Haberlandt notes the following distinct tissues :—(1) an epidermal tissue, sometimes developing trichomic structures ; (2) a mechanical system, consisting of elongated cells with thickened walls ; (3) an absorbing system, most strongly displayed in the rhizoids—also at the base of the sporange ; (4) an assimilating system, often composed of palisade-cells, in the leaves or in the sporange ; (5) a conducting system, consisting of the rudimentary 'vascular' bundles ;

(6) a reserve-system, usually represented by the aquiferous tissue; (7) a secreting system, developed typically in the sporange. No special secretory or excretory organs have been detected in mosses.

The *leaves* of mosses originate as broad papillose bulgings of a cell of the stem which becomes cut off by a septum. They are always of small size, sessile, and vary in shape from extremely narrow to broadly lanceolate or almost orbicular. The tissue of the greater part of the leaf usually consists of only a single layer of cells, all of which contain chlorophyll, except in the Sphagnaceæ and in Leucobryum, where the cells are of two different kinds, one large and empty, the other very small and chlorophyllous, thus giving the leaf a very light yellow-green colour. In most mosses the marginal cells of the leaf, and a row extending through the middle of the leaf from the base to the apex, are much smaller, and are disposed in several layers, though still thin-walled, thus constituting the rudiments of an epiderm and midrib. The midrib may even extend beyond the apex of the leaf as an awn or bristle. The leaves are usually crowded, especially in the neighbourhood of the sexual reproductive organs. Their phyllotaxis is spiral, or more rarely distichous. The branching of the stem of mosses is apparently neither dichotomous nor axillary; the number of lateral shoots is always much smaller than that of the leaves. When the primary shoot produces a so-called 'flower' at its apex, a lateral shoot situated beneath it not unfrequently displays a more vigorous growth of a monopodial character, and is then termed an *innovation*. *Prolification*, or the prolongation of a shoot by the continued growth of the bud within and above the male 'flower,' is a common phenomenon in Polytrichum. Nearly leafless shoots or *stolons* are sometimes formed beneath or on the surface of the ground, arising finally as erect leafy stems.

FIG. 104.—*Polytrichum commune* L.; proliferous male plant (slightly magnified).

In most mosses large numbers of *rhizoids* are formed on the under side of the stem, often clothing it completely with a reddish brown felt (see fig. 102). They differ from the protoneme in their tendency to grow downwards, and in not usually containing chlorophyll; but there is no sharp distinction between the two, each possessing the power of producing branches indistinguishable from those of the other.

The non-sexual propagation of mosses takes place in a variety of ways, the first step being, in nearly all cases, the production of a colourless filament of the nature of a protoneme. This protonemal branch, like the true protoneme which springs from a spore, may produce by lateral budding a number of leafy shoots. Protonemal branches of this kind are in some genera (such as Mnium, L., Barbula, Hedw., and Phascum, L.) produced in large numbers simply by turning over a tuft of moss on the soil. In this state they frequently hibernate, the portion above the soil disappearing entirely. In some mosses the leaves produce a protoneme simply by the growth of particular cells into segmented filaments; and this may even take place with detached leaves if kept moist. The seta or stalk of the sporogone has a special tendency to produce protonemes when in contact with damp soil. Rhizoids also may give rise to gemmæ or leaf-buds, whether above or beneath the surface of the soil. *Gemmæ* of a more complicated structure occur in a few species, as Aulacomnion androgynum (Schw.) (Bower, 'Journ. Linn. Soc.,' 1884, xx., p. 465), and Tetraphis pellucida (Hedw.), being stalked fusiform or lenticular multicellular bodies. In the latter species they are enveloped in an elegant cup or *cupule*, composed of a number of leaves, out of which they eventually fall, and then put forth protonemal filaments, which give rise first to a flat prothallus resembling that of a fern, and then to a leaf-bud. Gemmæ are produced in a variety of situations, as at the apex of a leaf and on the rhizoids.

FIG. 105.—*A*, young plant of *Barbula*, *m*; *h*, rhizoid, producing protonemes *p*, and underground gemma *k* (× 20). *B*, the same gemma (× 300). (After Goebel.)

The sexual organs of mosses are very commonly enveloped in closely crowded leaves which have undergone a certain amount of modification,

and the entire structure is sometimes—from analogy with the corresponding structure in Flowering Plants—called the *flower*. Such a flower may either be hermaphrodite, including both antherids and archegones, or unisexual, and the species may then be monœcious or dioecious. The female and the hermaphrodite flowers are not dissimilar in appearance, while that of the male flowers is altogether different.

FIG. 106.—*Funaria hygrometrica*; longitudinal section through male inflorescence; *a*, young antherid; *b*, nearly mature antherid; *c*, paraphyses; *d*, *e*, perigonal leaves (× 300). (After Goebel.)

In the hermaphrodite flowers the archegones usually occupy the central position, corresponding to the pistil in Flowering Plants, the antherids being arranged in an encircling spiral; while the whole is enveloped in a rosette of small leaves termed the *perichœte* or perianth. The entire structure resembles externally an elongated closed bud. Only a single archegone in each flower is usually fertilised or arrives at maturity. The female resemble the hermaphrodite flowers in every respect except

Fig. 107.—*Polytrichum commune*. *A*, *B*, mature plants with sporogone; *C*, male plant (nat. size).

that the antherids are suppressed. The male flowers present much greater variety in form and appearance; the male perianth or *perigone* is usually composed of broader, shorter, and thicker leaves, which sometimes sheath at the base, and are not unfrequently red. The flowers themselves are ovoid, globular, or discoid; the antherids usually stand in the axils of metamorphosed leaves. Both male and female flowers are provided with barren segmented filaments or *paraphyses*. In the male flowers the paraphyses are filiform, club-shaped, or spathulate, and terminate in several rows of cells; in the female flowers they are simple filiform bodies composed of a single row of cells. Their function appears to be to keep the archegones moist until they have been fertilised by the antherozoids.

The first antherid appears to be a terminal structure, being developed out of the apical cell of a branch. An hermaphrodite flower is probably derived from two independent shoots, the female shoot being formed immediately beneath the male organs. The mature *antherid* is a stalked club-shaped, or less often spherical sac, with a wall composed of only a single layer of cells. In the Sphagnaceæ it opens by longitudinal dehiscence; in the other orders by an apical slit through which the antherozoids, still enclosed in their mother-cells, are discharged as a thick mucilaginous mass, being imbedded in a jelly which is expelled in jets when the antherid bursts, but which is soluble in water. The *antherozoids* then escape from their mother-cell walls, and swim about as filiform bodies, furnished at the anterior end with two slender vibratile cilia, and containing a number of starchy granules. The male inflorescence of Polytrichum exhibits a remarkable tendency to prolification (see fig.

104). The *archegone* has somewhat the form of a very long-necked flask. The wall of the venter or ventral portion, which is ovoid and supported on a short stalk, consists of two layers of cells, while the elongated neck, which is often somewhat twisted, is composed of from four to six rows in a single layer. The interior of the *venter* consists of a single large cell, the *central cell*, which divides into two by a horizontal septum; the lower segment contains the *oosphere*, and the upper segment becomes the *ventral canal-cell*, while the neck is penetrated by a single axial row of

FIG. 108.—*A*, antherid of *Funaria hygrometrica*, discharging antherozoids, *a* (× 350). *B*, antherozoid of *Polytrichum*; *b*, still enclosed in mother-cell; *c*, free (× 800).



FIG. 109.—*Funaria hygrometrica*. *A*, longitudinal section through female inflorescence; *a*, archegones; *b*, perichætial leaves (× 100). *B*, archegone (× 500); *b*, venter and central cell; *h*, neck; *m*, opening of canal. *C*, opening of neck (more highly magnified), with stigmatic cells forced open.

cells which deliquesce into mucilage before impregnation. An open *canal* is thus left, through which the antherozoids penetrate to the oosphere; the terminal *stigmatic* or lid-cells of each row of the neck, constituting together the *stigma*, being forced apart by the exudation of the mucilage. The first archegones are formed from apical cells of shoots.

The impregnated oosphere, or *oosperm*, develops into an embryo, from which is derived the *sporogone* within the ventral portion of the archegone. After investing itself with a cell-wall, it divides by a number of longitudinal, radial, and transverse septa. At an early period in typical mosses the young elongated sporogone ruptures transversely the wall of the venter, the lower part of which forms a sheathing protection to its base, and is termed the *vagine*, while the upper part becomes elevated in the form of a cap or *calypter*. In the Sphagnaceæ the sporogone attains almost perfect development before the rupture of the archegone; in other mosses the various portions of which the sporange is composed are differentiated only at a later date. The sporange is at first filled with fluid contents, the greater part of which is the *archespore*,



FIG. 110.—*Funaria hygrometrica*. *A*, young plant with young sporogone. *B*, mature plant with mature sporogone; *s*, seta; *f*, sporange; *c*, calypter (natural size). *C*, longitudinal section of sporange (greatly magnified); *d*, opercule; *a*, annulus; *p*, peristome; *c*, columel; *s*, archespore; *h*, air-cavities. (After Goebel.)



FIG. 111.—Mouth of sporange of *Fontinalis antipyretica* L., with peristome: *ap*, teeth; *ip*, cilia (× 50). (After Schimper.)

developing into the mother-cells of the spores, from each of which are produced four spores by free-cell formation after preliminary indication of bipartition. The withered neck of the archegone, which has assumed a deep reddish brown colour, may often be recognised for some time surmounting the apex of the calypter. The mature sporogone consists of a pedicel or *seta*—which is usually of considerable length, the lower portion or *foot* being enclosed within the vagine, but is short in Sphagnum and some other genera—and the *sporange* or spore-capsule surmounted by the calypter, while the base of the seta is surrounded by the sheath-like *vagine*. The wall of the sporange is composed of several layers of cells, the outermost of which has a distinctly epidermal character, and is sometimes perforated by stomates with imperfect guard-cells.

While the greater part of the internal tissue is used up in the formation of the spores, the axial portion always remains unchanged in the form of a solid *columel*. There are no elaters. Leitgeb regards the sporogone of all mosses (including Sphagnaceæ) as consisting, in its earliest stage of development, of an inner mass of cells, the *endothecium*, distinctly separated from the peripheral mass, or *amphithecium*. In Sphagnaceæ the archespore is formed from the latter, in the other orders from the former portion. Among the typical mosses he again distinguishes three types, viz. :—(1) The *Archidium*-type : the spore-forming and sterile cells are intermingled in the endothecium ; the spore-sac is separated from the wall of the capsule by a bell-shaped cavity ; (2) the *Andreæa*-type : the endothecium is differentiated into the archespore and the columel, which does not penetrate the former ; the innermost layer of the amphithecium becomes the spore-sac, which is not separated from the wall of the capsule by any cavity ; (3) the *Bryum*-type : the endothecium is differentiated as in the last case, but the columel penetrates the spore-sac, which is separated from the wall of the capsule by a cylindrical cavity. In all true mosses the sporogone is developed by means of a two-edged apical cell. The ripe spores are roundish or cubical, with a thin, finely granulated yellowish, brownish, or purple cuticularised exospore, and an endospore of cellulose, and contain protoplasm, chlorophyll, and oil. The number of spores in a capsule varies from sixteen (Archidium, Brid.) to an immense quantity, and their size also varies inversely. Several cases of hybridism have been recorded in mosses.

Mosses are found in all climates, from the coldest to the hottest ; they are most abundant in temperate regions and in damp situations, clothing old walls, the trunks of trees, &c. A few grow in stagnant, and one genus (Fontinalis, L.) in running water. Some species are saprophytes. They are of scarcely any economical value, but are of great importance in nature in the formation of soil.

LITERATURE.

Bruch & Schimper—Bryologia Europæa, 1836–1865.
Schimper—Recherches anatom. et physiol. sur les Mousses, 1848.
Wilson—Bryologia Britannica, 1855.
Hofmeister—Pringsheim's Jahrb. wiss. Bot., 1863, p. 259.
Unger—Sitzber. Akad. Wiss. Wien, xliii., 1861, p. 497.
Lorentz—Moosstudien, 1864 ; Pringsheim's Jahrb. wiss. Bot., 1867, p. 363 ; Flora, 1867.
Berkeley—Handbook of British Mosses, 1863.
Leitgeb—Sitzber. Akad. Wiss. Wien, 1868, 1869.
Nägeli—Pflanzenphys. Untersuch., Heft i. p. 75.
Janczewski—(Archegonium) Bot. Zeit., 1872, pp. 377 *et seq.*
Stahl—Bot. Zeit., 1876, p. 689.
Kienitz-Gerloff—(Sporange) Bot. Zeit., 1878, pp. 33 and 49.

Braithwaite—The British Moss-Flora, 1880–1887.
L'Abbé Hy—Bull. Soc. Bot. France, 1880, p. 106; Ann. Sc. Nat., xviii., 1884, p. 105.
Goebel—Flora, 1882, p. 323.
Firtsch—Ber. Deutsch. Bot. Gesell., 1883, p. 83.
Satter—Ber. Deutsch. Bot. Gesell., 1884, p. 13.
Haberlandt—Ber. Deutsch. Bot. Gesell., 1883, p. 263; and Pringsheim's Jahrb. wiss. Bot., 1886, p. 359.
Magdeburg—Die Laubmooskapsel als Assimilations-Organ, 1886.
Vaizey—(Sporogone) Ann. of Bot., i., 1887, p. 73.
Limpricht—Die Laubmoose, in Rabenhorst's Crypt.-Flora Deutschland, 1885–1888.

The Musci are classified under four orders, as follows. The Sphagnaceæ exhibit much more important peculiarities than the other three orders, and are ranked by some writers of authority as a distinct class.

Order 1.—Bryaceæ.

This order includes the vast majority of the genera of mosses, and all the more familiar forms except the bog-mosses. The sporanges or 'fruits' form objects of great beauty in the autumn and winter, their usual period of maturity, fertilisation taking place in the spring or early summer. In some species the sporogone requires more than a year for its full development. The sporogone consists of a sporange which is always surmounted by a calypter, easily removed by the wind; beneath this is the opercule, which becomes detached, either alone or together with the *annulus*, a circular layer of hygrometric epidermal cells between the opercule and the edge of the capsule; the whole elevated on a longer or shorter stalk or *seta*, which is inserted at its lower end in the *vagine*. The portion of the seta concealed in the vagine is known as the *foot*, and acts as a kind of root, all the food-material needed for the development of the sporogone being absorbed through it. The central strand of tissue in the seta of the Polytrichaceæ consists, according to Vaizey, of two portions—a *leptophloem* or rudimentary phloem, in which the storing up and conduction of the food-material takes place; and a *leptoxylem* or rudimentary xylem, which serves for the conduction of the transpiration-current to the lower portion of the sporange furnished with stomates. In the Polytrichaceæ, in addition to the opercule, a horizontal layer of cells termed the *epiphragm* remains attached to the points of the teeth of which the peristome is composed, and covers the mouth of the sporange after the removal of the opercule. The sporange is penetrated by a complete axial columel. The spores are formed by free-cell formation in fours within *spore-mother-cells*, themselves derived from a single primordial layer, the *archespore*; the walls of the spore-mother-cells finally deliquesce, leaving the spores floating in a fluid

which for a time fills the *spore-sac*, composed of two or three layers of barren cells immediately enclosing the mass of spores. Between the spore-sac and the wall of the mature sporange is an annular *air-cavity*, traversed horizontally by rows of chlorophyllous cells, the *trabecules*. The opercule is simply a piece of the epiderm of the sporange. In most genera a portion of the wall of the sporange, situated near the base of the columel, consists of an assimilating system composed of spongy or palisade-parenchyme, containing chlorophyll, and marked by the presence of stomates. If the detachment of the opercule leaves the mouth of the sporange with a smooth edge, it is said to be *gymnostomous*, as in Pottia (Ehrh.). More often the mouth of the open sporange is furnished with hair- or tooth-like appendages, arranged in one or two rows, constituting the *peristome*. The single row of these appendages, or the outer row if there are two, are called *teeth*, the inner row *cilia*. In some genera the cilia are furnished with lateral processes uniting them with one another, or they are replaced by a lattice-work of longitudinal or transverse ridges termed the *endostome*. The inner and outer layers of both teeth and cilia differ from one another in their hygroscopic capacity; hence, as the moisture of the air varies, they bend inwards and outwards, or sometimes coil spirally round one another (Barbula, Hedw.). The peristome has a very beautiful appearance under the microscope, and its structure furnishes useful characters for the discrimination of the genera. In most genera the teeth and cilia are not composed of cells, but of pieces of thickened cell-wall which become detached from a



FIG. 112.—*A*, longitudinal section of sporange of *Polytrichum piliferum* Schreb. (×15). *B*, transverse section (×50. *w*, wall of sporange; *c*, columel; *p*, peristome; *ep*, epiphragm; *a*, annulus; *i*, air-spaces traversed by trabecules; *s*, spore-sac; *ap*, apophyse; *st*, seta. (After Lantzius-Beninga.)

layer of cells beneath the epiderm; but in Polytrichum the teeth are composed of bundles of thickened prosenchymatous cells. The exterior peristome may have two distinct forms. Either the teeth have a double outer and a single inner series of plates (*Diplolepidæ*), or the exterior series is simple (*Aplolepidæ*), and then the inner series is nearly always double. The Aplolepidæ never have a double peristome; and in the Diplolepidæ the inner peristome is occasionally wanting in particular families or genera. In much the larger number of genera (*Arthrodonteæ*) the teeth are septated by transverse walls; in a much smaller number

FIG. 113.—*Hypnum populeum* Sw. (natural size).

FIG. 114.—*Tetraphis pellucida* Hedw. *a* (slightly magnified), with open sporange; *b*, ditto with gemma; *c*, sporange with calypter (greatly magnified); *d*, open sporange, showing peristome.

FIG. 115.—*Bryum argenteum* L. (natural size).

FIG. 116.—*Splachnum ampullaceum* L. (natural size).

(*Nematodonteæ*) the transverse septa are wanting. The cilia, when present, are usually shorter and less developed than the teeth; they are also composed of two layers of plates, often marked on the surface by a beautiful network; their divisions correspond to those of the teeth. The number of both teeth and cilia is always a multiple of 4, the most common numbers being 8, 16, 32, and 64. The most perfect type of peristome is seen in the Encalypteæ, from which all the less perfect forms are, according to Philibert, derived by degradation. In addition to the presence of an epiphragm, the genus Polytrichum presents the peculiarity, in most species, of the seta being swollen beneath the sporange, forming an annular cushion known as the *apophyse* (see fig. 112).

In most Bryaceæ the tissue of the leaf is nearly homogeneous, with the exception of the margin and an elementary midrib composed of elongated prosenchymatous cells; but in Leucobryum (Hpe.) the small chloro-



FIG. 117. Sporange of *Polytrichum commune*, showing epiphragm. *A*, covered by calypter; *B*, with calypter removed; *C*, with opercule removed (magnified).

FIG. 118.—Sporange of *Hypnum populeum*, showing peristome (magnified).

phyllous cells interlace among large empty cells with circular orifices in their walls, as in Sphagnum.

The very numerous genera of Bryaceæ are further classified as under.

Acrocarpi.—Fructification produced at the extremity of the branches. *Illustrative genera*:—Weissia (Hedw.), Dicranum (Hedw.), Leucobryum (Hpe.), Pottia (Ehrh.), Tortula (Hedw.), Bartramia (Hedw.), Encalypta (Schreb.), Fissidens (Hedw.), Grimmia (Ehrh.), Orthotrichum (Hedw.), Zygodon (H. & T.), Tetraphis (Hedw.), Buxbaumia (Hall.), Polytrichum (Dill.), Aulacomnion (Schw.), Bryum (H. & T.), Mnium (B. & S.), Funaria (Schreb.), Splachnum (B. & S.), Tayloria (Hook.), Barbula (Hedw.), Ceratodon (Brid.).

FIG. 119. Peristome of *Atrichum undulatum* (magnified).

FIG. 120.—Peristome of *Cinclidium stygium* (magnified).

Pleurocarpi.—Fructification lateral, not at the extremity of the principal branches. *Illustrative genera*:—Hedwigia (Ehrh.), Fontinalis (L.), Hookeria (Sm.), Hypnum (Dill.), Leucodon (Schw.), Neckera (Hedw.).

LITERATURE (in addition to the papers already quoted).

Philibert—(Peristome) Rev. Bryol., 1884-1888.
Vaizey—(Polytrichaceæ) Journ. Linn. Soc., xxiv. (1888), p. 162.

Order 2.—Phascaceæ.

In the small order of Phascaceæ the roundish sporange dehisces neither by the detachment of an opercule nor by longitudinal slits, but decays to allow of the escape of the spores; the calypter is ruptured laterally without being raised up as a cap; the columel is sometimes wanting. According to Leitgeb, Archidium (Brid.) resembles the Hepaticæ more closely than the Bryaceæ in the processes which lead to the formation of the spores, especially in the differentiation of the archespore into spore-mother-cells which are irregularly interspersed among cells that remain sterile. The spore-mother-cells do not number more than from one to seven in each sporange; in each of them four spores are formed tetrahedrally.

Fig. 121.—*Ephemerum serratum* Hampe; mature plant with persistent protoneme (magnified). (After Luerssen.)

Fig. 122.—*Pleuridium subulatum* Rabenh.; sporange (magnified). (After Luerssen.

The Phascaceæ are cæspitose in their habit; the protoneme persists until the maturity of the sporogone. *Principal genera* :—Phascum (L.), Archidium (Brid.), Ephemerum (Hpe.), Pleuridium (Brid.).

Literature.

Leitgeb—Sitzber. Akad. Wiss. Wien, 1880, p. 447.
Müller—Pringsheim's Jahrb. wiss. Bot., 1867, p. 237.

Order 3.—Andreæaceæ.

The Andreæaceæ constitute a small order of mosses, comprising the single genus Andreæa (Ehrh.), characterised by the absence of an opercule to the sporange, which opens by four, or very rarely eight, longitudinal slits, not reaching either to the base or the apex of the capsule.

The calypter is elevated, as in the Bryaceæ, on the summit of the ripe sporogone in the form of a cap ; there is a short seta buried in the vagine, and the whole sporogone is elevated on a stalk or *pseudopode*, as in the Sphagnaceæ. At the base of the sporange is an enlarged apophyse. The structure of the sporange differs from that of the Bryaceæ in the columel not penetrating the archespore, and in the absence of a cavity between the spore-sac and the wall of the sporange. The contents of the spore divide, while still within the exospore, into four or more cells. As in Sphagnum, the oosphere is always enveloped in a hyaline mass of mucilage in which the antherozoids imbed themselves.

The Andreæaceæ are also cæspitose in their habit, and are natives of cold or mountainous regions.

Fig. 123.—*Andreæa alpestris* Schmp. (× 5).

Fig. 124.—*A. alpestris* Schmp.; dehiscent sporange and apophyse (magnified).

Literature.

Kühn—Entwickelungsgeschichte der Andreæaceen, 1870.
Waldner—Bot. Zeit., 1879, p. 595 ; and Entwick. d. Sporogone v. Andreæa, 1887.

Order 4.—Sphagnaceæ.

The bog-mosses form a large portion of the vegetation of bogs and swamps, and are characterised by the spongy structure of the whole plant, the light yellowish green colour of the leaves, and the bright red globular spore-capsules. The protoneme is much less developed than in typical mosses ; and when the spore germinates on dry ground a flat *prothallium* intervenes between it and the leafy stem. The stem branches abundantly, giving a cæspitose appearance to the whole plant ; and *innovations*, produced below the apex after the ripening of the fructification, become detached by the decay of the lower part of the stem, and carry on an independent existence. The leaves are lanceolate and apiculate, usually arranged in a $\frac{2}{5}$ phyllotaxis, larger than in other mosses, and of a peculiar structure of their own. As the leaf develops,

the single layer of cells of which it is composed becomes differentiated into cells of two distinct kinds. A comparatively small number, of a lozenge-shaped form, grow to a large size and lose the whole of their contents, while their walls are provided with spiral thickening-bands, and frequently display a large circular orifice opening from one cell into the next. A much larger number of cells remain permanently of a very small size, are very narrow in proportion to their length, and, being filled with protoplasm and chlorophyll, constitute the whole of the nutritive tissue of the leaf. These nutritive cells form a kind of network ramifying among the large empty cells ; but, as their total area is small compared with these latter, the entire leaf has, to the naked eye, a semi-transparent very light yellow-green appearance. The tissue of the stem consists of cells of three distinct kinds. In the centre is a cylinder of thin-walled elongated colourless parenchymatous cells ; this is enveloped in a layer of dotted prosenchymatous cells, the walls



FIG. 125.—*Sphagnum acutifolium* Ehrh. *A*, megaspore ; *B*, microspore ; *C*, protoneme, with *pr*, rudiments of young plants (magnified). (After Schimper.)



FIG. 126.—Flat prothallium of *S. acutifolium*, with young leafy stems (× 120). (After Schimper.)

of which are thicker and of a brown colour ; while outside all is an epidermal layer of large thin-walled empty cells, sometimes with spiral

thickenings and circular orifices, similar to those of the leaves. These serve as capillary tubes, through which the water of the bogs in which these mosses grow is raised, and the whole plant is in consequence always saturated with water like a sponge.



FIG. 127.—*A*, portion of surface of leaf of *S. acutifolium*. *cl*, small chlorophyllous cells; *f*, large empty cells; *l*, orifices in these cells. *B*, transverse section (magnified).



FIG. 128.—Transverse section of stem of *S. cymbifolium* Dill. *x*, inner cells with colourless wall; *r*, outer layer of cells; *ee*, peripheral layers of cells with orifices, *l* (× 900). (Aft r Luerssen.)

In their organs of reproduction, and especially in the structure of the sporogone, the Sphagnaceæ exhibit some divergences from the typical mosses. Some species are dicœcious, and the 'flowers' of the moncœcious species are never hermaphrodite, the male and female organs being always distributed on different branches. The male branches are distinguished by their densely crowded leaves, which are often of a bright red colour, giving a catkin-like appearance to the branch. On removing these the antherids are found near the middle of the branch. They are minute nearly globular or elliptical bodies, elevated on a slender stalk, and dehiscing by longitudinal fission into valves. The antherozoids are spiral bodies with many coils, and two large flagellate cilia at the posterior end. The archegones are formed towards the extremity of the female branches, are accompanied by paraphyses, and are enveloped by perichætial leaves. They resemble in all essential points those of other mosses. On impregnation the oosperm divides by a horizontal septum into two cells, the sporogone originating from the upper cell only. The nearly spherical usually bright red sporange differs from that of other mosses in being completely enclosed within the venter almost till maturity. It is, in most species, elevated on a long slender pedicel, the *pseudopode*, which must not be confounded morphologically with the seta of other mosses, being a prolongation of the axis below the vagine. The lower portion of the sporogone is widened out into a broad disc-like *foot*, resembling in appearance the apophyse of Polytrichaceæ, which is seated on the top of the pseudopode, and enclosed in the vagine. The calypter, when finally ruptured, is not elevated in the form of a cap, but remains attached as a

FIG. 129.—*S. acutifolium*. *a*, antheridial branches; *b*, leaves of primary stem; *ch*, perichætial leaves with sporogones (× 5). (After Schimper.)

frill to the base of the sporange, which dehisces by a transverse slit near the apex, detaching a strongly convex opercule. There is no peristome nor annulus. A portion of the contents of the sporange remains undifferentiated in the form of a low columel not reaching to the apex. The remainder is converted into spores, which differ from those of other mosses in being of two kinds, *megaspores* and *microspores* (see fig. 125). According to Warnstorf ('Hedwigia,' 1886, p. 89; and 'Verhandl. Bot.



FIG. 130.—*S. acutifolium*. *A*, male branch, with leaves removed to expose antherids, *a* (magnified). *B*, antherid (more highly magnified) dehiscing. *C*, antherozoid (still more magnified). (After Schimper.)



FIG. 131.—*A*, *S. acutifolium*, section of female inflorescence; *ar*, archegones; *ch*, perichætial leaves. *B*, longitudinal section of sporogone, *sg*; *ar*, archegone; *c*, calypter; *sg'*, foot; *v*, vagine; *ps*, pseudopode. *C*, *S. squarrosum* Pers.; *sg*, sporogone; *d*, opercule; *c*, ruptured calypter; *qs*, pseudopode; *ch*, perichætial leaves (magnified). (After Schimper.)

Verein,' Brandenburg, 1886, p. 181), these two kinds of spore are found either in the same or in different sporanges; the diameter of the former varies between 30 and 33 mm., that of the latter between 12 and 18 mm. The megaspore is by far the most common form, and its germination only has at present been observed. Warnstorf suggests that the two kinds are sexually differentiated, the megaspores giving rise to a female, the microspores to a male prothallium, as in the Heterosporous Vascular Cryptogams.

All the species of the single genus Sphagnum (L.) grow in bogs and swamps, often covering enormous tracts of ground, and entering largely into the composition of peat. The best authorities differ widely as to the number of species; the most divergent forms are distinguished by well-marked characters, but these merge into one another by a complete series of connecting links.

LITERATURE.

Schimper—Entwickelungsgeschichte der Torfmoose, 1858.
Waldner—Bot. Zeit., 1879, p. 595; and Entwickelung d. Sporogone v. Andreæa u. Sphagnum, 1887.
Braithwaite—Sphagnaceæ of Europe and North America, 1880.
Warnstorf—Die Europ. Torfmoose, 1881; and Flora, 1884.
Limpricht—Bot. Centralbl., x., 1882, p. 214.
Röll—Flora, 1886.

Class VIII.—Hepaticæ.

The Hepaticæ or Liverworts are small elegant plants, usually of a bright green colour, which are especially abundant on damp ground or rocks, or by the sides of streams; a few species are aquatic. Some of the genera bear a considerable external resemblance to lichens, others to mosses. Their vegetative structure is either an undifferentiated thallus, or consists of a distinctly differentiated stem and leaves, in both cases attached to the soil by rhizoids. The former are known as the *Thalloid* or *Frondose*, the latter as the *Foliose Hepaticæ*. The transition marks the passage from the upper of the two great divisions of the vegetable kingdom, the Cormophytes, to the lower division, or Thallophytes; but intermediate forms occur in the genera Fossombronia (Rad.) and Blasia (Mich.). Even the foliose forms have no true vascular tissue, and no true roots, the functions of which are performed by the rhizoids. Both sections have, except in Riella (Mont.) and Haplomitrium (N. ab E.), a distinct bilateral or dorsiventral structure; the free side which faces the light is differently organised from that which faces and often clings to the substratum, and which is not exposed to light. The mode of branching in the thalloid forms is dichotomous, and the growing region of the shoot commonly lies in an apical depression formed by the more rapid growth of the cells lying right and left of the apical cell, which has a form allied to wedge-shaped. The filiform stem of the foliose forms, on the other hand, ends in a bud with a more or less prominent cone of growth, and the apical cell is a three-sided pyramid. The leaves of the foliose forms always consist of only a single layer of cells, without even a rudimentary midrib; while the stem sometimes contains the first rudiments

of vascular bundles in the form of cambium-strings, and is furnished with a slightly differentiated epidermal layer. In the thalloid forms the thallus is composed of a more or less thick plate of tissue, which in one order, the Marchantiaceæ, possesses on the upper side a strongly developed epiderm, provided with stomates of very peculiar form, unlike anything that occurs elsewhere in the vegetable kingdom.

The first result of the germination of the spore is either a filiform protoneme, a flat plate of cells, or a mass of tissue ; but the differentiation of the protoneme from the sexual generation is not so well marked as in the Musci. The non-sexual propagation of the Hepaticæ takes place either by *innovation*, *i.e.* by the continual dying away of the stem behind, or by *gemmæ*, which exhibit a high degree of development. In the thalloid genera Marchantia (L.), Lunularia (Mich.), and Blasia, these gemmæ are found in peculiar outgrowths of the upper surface of the thallus known as *cupules*, which are cup-shaped in Marchantia (see fig. 150), crescent-shaped in Lunularia, flask-shaped in Blasia. From the base of these cupules there spring hair-like papillæ, the apical cells of which divide repeatedly in both directions, and constitute the gemmæ. In some of the foliose genera, *e.g.* Madotheca (Dum.), the gemmæ are formed out of cells belonging to the margin of the leaf, and simply detach themselves. Vöchting states that in Lunularia, and Marchantia also, isolated masses of cells possess the power of regeneration or development into new individuals, to whatever part of the thallus they may have belonged. Shoots resembling a normal thallus spring from the pedicel of the inflorescence of Marchantia polymorpha (L.) when lying prostrate on the soil.

FIG. 132.—*Jungermannia nemorosa* L. (× 10).

The locality of the sexual organs of reproduction, antherids and archegones, varies in the different orders. In one genus, Anthoceros (Mich.), they are endogenous, or originate in the tissue of the thallus itself ; in the remaining thalloid forms they are produced on the upper side of the thallus ; and in the Marchantiaceæ on special vertical outgrowths, some of which bear antherids on their upper, others archegones

on their under side ; these male and female *inflorescences*, as they are termed, may be either monœcious or dicœcious. In the foliose orders there is a great variety in their locality and mode of origin.



The antherid originates as a papilliform swelling of a superficial cell, from which it is marked off by a septum. When mature it is seated on a pedicel or stalk, and consists of an external layer of cells containing chlorophyll, which encloses the mother-cells of the antherozoids. The antherid dehisces by longitudinal fissures into valves, and the mother-cells themselves escape into the surrounding moisture, into which each discharges an antherozoid. The antherozoids are slender threads of protoplasm, with from one to three spiral coils, and are provided at the anterior end with two long and very slender cilia, by means of which they 'swarm' in the water with a rotating motion. The archegone also first makes its appearance as a papillose outgrowth of a superficial cell, which then becomes separated in the same manner. After this mother-cell has divided several times longitudinally, the central one of

FIG. 133.—*Gottschea appendiculata* N. ab E. (magnified).

the cells thus formed divides transversely into an upper *stigmatic* or *lid-cell* and a lower cell. Two layers are subsequently formed, the upper of which becomes the neck of the archegone, the lower its ventral

portion, or *venter*. The lowermost cell of this ventral portion, now known as the *central cell*, increases considerably in size, and divides by a transverse septum into a lower and larger portion, which encloses the *oosphere*, and an upper and smaller portion, the *ventral canal-cell*. In the meantime the upper layer of cells increases in length by the formation of a number of fresh cells, the *neck-canal-cells*. The ventral portion of the archegone becomes eventually enclosed in a wall, and by the deliquescence of the neck-canal-cells an open channel is formed down to the oosphere. In addition to the perichæte, the archegones are frequently surrounded by a circular wall, originating as an outgrowth of the thallus, and known as the *perigyne* or involucre.

The result of the impregnation of the oosphere by one or more antherozoids is the production of the embryo, from which is derived the *sporogone*, which alone constitutes the sporophyte, or non-sexual generation. It is formed entirely from the ventral portion of the archegone. Its external form and internal structure vary greatly in the different groups : as is also the case with the course of cell-divisions in its formation. Ultimately the wall of the spore-capsule becomes differentiated from the *archespore*, or layer of tissue which develops into the mother-cells of the *spores* and *elaters* when these latter organs are present. There is usually no solid axis or columel. The cells which develop into elaters cease early to divide transversely, and thus remain long, while the rest of the cells round themselves off, and become mother-cells of spores. The mature elaters (fig. 159) have in their wall an elongated single or double spiral band, the twisting and untwisting of which on the absorption and giving off of moisture helps to disseminate the spores. Leclerc du Sablon finds the sporogone of the typical Hepaticæ to be composed, at a very early stage, of sixty-four cells, each of which subsequently divides into four. These cells now elongate in the direction of the axis of the sporogone, and then become differentiated into two kinds. In the one kind the nucleus undergoes repeated bipartitions, and these give rise to the spore-mother-cells ; in the other kind the nucleus does not divide, and the protoplasm forms spiral granulations ; these become the elaters. Rarely (as in Riella) they are replaced by barren cells filled with food-material for the nutrition of the growing spores. The two kinds of cell are equal in number, each alternating with the other. The degree of complexity of the sporogone in the different orders of Hepaticæ corresponds in the main to the degree of development of the vegetative organs. In the Jungermanniaceæ it bursts longitudinally into four valves, and the walls are composed of two layers of cells furnished with 'ornaments,' or elevated markings of various patterns ; in the Anthoceroteæ it splits longitudinally into two valves ;

in the Ricciaceæ and Marchantiaceæ it bursts irregularly, and the wall is composed of a single layer of cells without ornaments, or nearly so. The spores also vary considerably in the different orders. In many Jungermanniaceæ the spore has only a single cuticularised membrane, which is entirely used up in the formation of the germinating filament. In most genera the wall is composed of two distinct separable layers, the *exospore* and *endospore*; while in Sphærocarpus (Mich.), Corsinia (Radd.), and some others there is a third outer layer, often beautifully sculptured, which is derived from the membrane of the special mother-cells of the spores. This layer is called by Leitgeb the *perinium*. Warnstorf (Verhandl. Bot. Ver. Brandenburg, 1886, p. 181) finds in Blyttia (Endl.) two kinds of spore, larger and smaller, which he believes to produce female and male plants respectively. In Sphærocarpus the spores are combined into tetrads. When the spore germinates, the endospore breaks through both exospore and perinium, when the latter is present, and protrudes as the first rhizoid.

LITERATURE.

Bischoff—Nov. Act. Acad. Leop. Car., 1835.
Gottsche—*Ibid.*, 1838.
Gottsche, Lindenberg u. Esenbeck—Synopsis Hepaticarum, 1844.
Kny—Pringsheim's Jahrb. wiss. Bot., 1865, p. 64.
Leitgeb—Bot. Zeit., 1871, p. 557, and 1872, p. 33; Mittheil. naturw. Ver. Steiermark, 1872; Unters. über die Lebermoose, 1874–1881; and Ber. Deutsch. Bot. Gesell., 1883, p. 246.
Janczewski—(Archegone) Bot. Zeit., 1872, p. 372 *et seq.*
Carrington—British Hepaticæ, 1874.
Kenitz-Gerloff—Bot. Zeit., 1875, pp. 777 *et seq.*
Vöchting—Pringsheim's Jahrb. wiss. Bot., 1885, p. 367.
Satter—Sitzber. Akad. Wiss. Wien, 1882.
Leclerc du Sablon—(Antherozoids) Comptes rendus, cvi., 1888, p. 876.

Liverworts are distributed throughout the entire globe, growing mostly in moist situations. Many tropical species are epiphytic on the leaves of Flowering Plants or ferns. They are of no economic importance. They are classified under five orders, of which the first includes both foliose and thalloid, the remaining four almost entirely thalloid, forms.

ORDER 1.—JUNGERMANNIACEÆ.

In this, much the largest order of the class, are included genera with every variety of vegetative development, from an undifferentiated thallus to a slender filiform stem, with sessile leaves seated either in two rows on the upper side, or in three rows, two of them on the upper, and the third, the *amphigasters*, smaller and adpressed to the under side. The thalloid

forms have, except in Haplomitrium, a bilateral structure resembling that of the Marchantiaceæ ; rhizoids and rudimentary foliar structures are formed on the under, the sexual organs on the upper side. In the foliose forms the leaves, always very small, are frequently bisected or bilobed, the lower lobe or *auricle* being the smaller one, and amplexicaul or concave. Goebel states that in Java many of the Jungermanniaceæ are epiphytic, and that in these the auricle is frequently hollowed out into the shape of a pouch or pitcher, serving as a receptacle for water (fig. 140). In some species of Physiotium (N. ab E.) this receptacle is prolonged into the so-called 'tubular organ.' The leaves of the foliose

FIG. 134.—*Calypogeia Trichomanis* Cord. (magnified).

FIG. 135.—*Fossombronia pusilla* N. ab E., male plant. *a*, natural size ; *b*, magnified.

species consist of a single layer of cells without even the rudiments of vascular bundles. There are species which form a connecting link between the foliose and the thalloid Jungermanniaceæ. The mode of branching varies greatly, but growth always takes place by means of a three-sided pyramidal apical cell.

As respects the sexual organs, some species are monœcious, others dicœcious. In the foliose genera they are usually formed at the apex of the primary shoots or of special small fertile branches, which have often an endogenous origin on the ventral side. These constitute the *acrogynous* section of the order, which includes all the foliose genera except Haplomitrium. In the thalloid genera, or *anacrogynous* section, they

appear on the dorsal surface of the shoot, at some distance from the apex; while in the acrogynous forms they are formed in close proximity to the apical cell. In Radula (Dum.) the entire female inflorescence, composed of a number of archegones enclosed in a perigyne, is developed from the apical cell of a shoot, and from its youngest three segments. Neither archegones nor antherids are elevated on receptacles, as in the Marchantiaceæ. The antherids usually occur

FIG. 136.—*Pellia epiphylla* Cord., male plant. *a*, natural size; *b*, magnified.



FIG. 137.—*Radula complanata* Dum. Plant with closed and open sporange (× 2).

FIG. 138.—*Jungermannia barbata* Schreb. Under side of leaves with ciliated amphigasters (magnified).

singly or in groups in the axils of the leaves. In Pellia (Radd.) the antherids are imbedded in the thallus, the archegones appearing in large numbers at the apex of the shoot. In the Geocalyceæ (*e.g.* Calypogeia, Radd.) the female branches are so hollowed out that the archegones are



FIG. 139.—I. Under side of stem of *Frullania Tamarisci* Dum., with true leaves and amphigasters (magnified). II. Leaf of *F. dilatata* (more magnified).

FIG. 140.—Auricle of *Frullania*, sp. (magnified). (After Goebel.)

sunk in a deep pitcher-shaped hollow or tube, within which the sporogone is subsequently formed. In other genera they are concealed by the nearest leaves. The modified leaves which thus enclose a group of archegones, or of both archegones and antherids, constitute the *perichæte*,

each archegone being, in addition, usually surrounded by a distinct membranous envelope, the perianth or *perigyne*.

In the formation of the sporogone, the fertilised oosphere first divides by a wall at right angles to the axis of the archegone. Only the upper of the two cells thus formed—that is, the one that faces the neck of the archegone—undergoes further divisions; it becomes the apical cell of the sporogone, and sometimes again divides transversely once or twice before a longitudinal wall makes its appearance in it; the two cells thus formed finally divide into four apical cells arranged as octants of a hemisphere. The basal portion of the growing archegone swells out and penetrates down into the tissue of the stem, forming the *vagine*.

FIG. 141.—Sexual organs of *Radula complanata*. *ar*, archegone; *an*, antherid; *b*, leaf. (After Hofmeister.)



FIG. 142.—*Jungermannia bicuspidata* L. Longitudinal section of immature sporogone, *sg*; *ar*, calypter; *ar'*, unfertilised archegones; *p*, base of perigyne; *st*, stem; *b*, leaf. (After Hofmeister.)

After frequent divisions have taken place, the wall of the spore-capsule becomes differentiated from the inner tissue, out of which are developed the spores and elaters. There is no columel. By rapid extension of the hitherto short pedicel, the calypter is ruptured at the apex, and the globular sporogone, containing the already ripe spores, becomes elevated. The inner of the two layers of which the wall of the sporogone is composed has become absorbed before the ripening of the spores; the single layer of cells which still remains is ruptured at the apex, and splits into four (rarely more) longitudinal valves, which separate suddenly in the form of a star, carrying with them at the same time the elaters, and thus bringing about the dispersion of the spores. The mature elaters are

long fusiform thin-walled cells, marked internally by from one to three brown spiral bands, but more complicated in structure in the foliose than in the thalloid genera.

Illustrative genera.—*Foliose*: Radula (Dum.), Jungermannia (L.), Lejeunia (G. & L.), Frullania (Radd.), Madotheca (Dum.), Mastigobryum (N. ab E.), Calypogeia (Radd.), Lepidozia (Dum.), Plagiochila (Dum.), Geocalyx (N. ab E.), Chiloscyphos (Cord.), Gymnomitrium (N. ab E.), Lophocolea (Dum.). *Thalloid*: Metzgeria (Cord.), Aneura (Dum.), Fossombronia (Radd.), Pellia (Radd.), Blasia (Mich.).

LITERATURE.

Leitgeb—Bot. Zeit., 1871, p 556; and Abhandl. Bot. Ver. Brandenburg, 1880, p. 58.
Gottsche—Abhandl. Gesell. Naturf. Hamburg, 1880, p. 39.
Goebel—(Epiphytic Species) Ann. Jard. Bot. Buitenzorg, vi., 1887, p. 21.

ORDER 2.—MONOCLEACEÆ.

This small order appears to occupy an intermediate position between the Jungermanniaceæ and the Anthoceroteæ. The vegetative structure is either thalloid or foliose. The elongated sporange dehisces longitudinally, and contains elaters, but has no columel.

Principal genus:—Monoclea (Hook.).

FIG. 143.—*Monoclea Forsteri* Hook. (magnified).

FIG. 144.—*Anthoceros lævis* L., male plant (natural size).

FIG. 145.—*A. lævis*. *a*, dehiscent sporange (× 2).

ORDER 3.—ANTHOCEROTEÆ.

The vegetative structure consists of a flat ribbon-like thallus, the irregular dichotomous ramifications of which form a circular disc composed of one or more layers of cells, each cell containing only a

single chlorophyll-corpuscle. The antherids and archegones arise endogenously on the upper side of the thallus, apparently without any definite arrangement, and are not protected by a perigyne. The mature sporogone is an elongated dehiscent two-valved pod, provided with stomates, which forces its way through a mass of tissue overarching the archegone, and is known as the *involucre*. Within the sporange is, in most genera, a solid axial *columel*; the wall consists of four or five layers of cells; the rest of the contents, developed from the archespore, becoming the mother-cells of the spores and elaters. Except in some species of Anthoceros (L.) the elaters are of simpler structure than in the other Hepaticæ, having no spiral bands. Anthoceros possesses peculiar cavities on the under side of the thallus, opening by slits or fissures, which are regarded by some authors as stomates, by others as mucilage-receptacles. Filaments of Nostoc, which have found their way into these cavities through the slits, cause peculiar changes in them, and have been mistaken for endogenous gemmæ. The genus Anthoceros is of much interest from the fact that the sporophyte-generation shows a greater vegetative energy than is usually the case with Muscineæ; growth continues at the base of the sporange, and new spores are formed there after those at the apical portion are already mature.

Principal genus:—Anthoceros (L.).

LITERATURE.

Leitgeb—Die Anthoceroteen, 1879.

ORDER 4.—RICCIACEÆ.

The Ricciaceæ are regarded by Leitgeb as forming a connecting link between the Jungermanniaceæ and the Marchantiaceæ; but in some respects they are simpler in their structure than either of these orders. The thallus is usually flat, and branches dichotomously; it floats on water or roots in the soil. In Riella (Mont.) it is submerged and erect, and has the appearance of a ring forming a continuous spiral round an axial stem. It is always destitute of stomates, but is provided with internal air-cavities, and with rudimentary foliar organs among the rhizoids. The antherids and archegones are not, as in the Anthoceroteæ, endogenous, but are developed from young superficial cells of the upper surface, which grow into papillæ and become overarched, in the course of their development, by the surrounding tissue. Both antherids and archegones are enclosed in an *involucre* formed in this way; the antherids are sessile, the involucre sometimes constituting an elevated neck above them. In Riccia (L.) the archegones are ultimately buried in the

thallus; while in Oxymitra (Bisch.) they are raised above the surface. The sporange is a thin-walled spherical capsule, occasionally produced under water, entirely filled with spores without true elaters or columel, and, with its calypter, depressed in the thallus. It is much less differentiated in its structure than in the other orders. In all the genera except Riccia and Oxymitra the elaters are represented by sterile cells among the spore-mother-cells. The sporange bursts irregularly when ripe, but the spores are only set free by the decay of the surrounding tissue of the thallus. The spores of Sphærocarpus (Mich.) and Corsinia (Radd.) have a beautifully sculptured extine. Riella is altogether diœcious, and perfects its fructification beneath the water.

FIG. 146.—*Riella helicophyila* Mont. (magnified)

FIG. 147.—*Sphærocarpus terrestris* Sm. Frond and archegone (magnified).



FIG. 148.—*Riccia glauca* L. *A*, section of apical region of frond. *ar*, archegone; *c*, oosphere (× 50). *B*, immature sporogone, *sg*; *ar*, neck of archegone (× 300). (After Hofmeister.)

Principal genera:—Riccia (L.), Duriæa (Bor.), Oxymitra (Bisch.), Riella (Mont.), Sphærocarpus (Mich.), Corsinia (Radd.).

LITERATURE.

Kny—Pringsheim's Jahrb. wiss. Bot., 1866, p. 364.
Leitgeb—Die Riccieen, 1879.

Order 5.—Marchantiaceæ.

The thallus is flat and ribbon-shaped, and usually branches dichotomously from two apical cells; it is frequently furnished with a well-



Fig. 149.—*Marchantia polymorpha* L. Male plant (natural size).

Fig. 151.—*M. polymorpha.* Female inflorescence (magnified).

Fig. 150.—*M. polymorpha.* Male inflorescence and cupule (magnified).

Fig. 152.—*Fegatella conica* Cord. Male plant (natural size).

marked midrib, and is coriaceous in texture. It is composed of three distinct layers of cells, viz.:—(1) the air-chamber-layer to which the

stomates belong; (2) a close tissue containing but little chlorophyll, and with the cell-walls pitted or reticulately thickened, without intercellular spaces but sometimes containing mucilage-receptacles; and (3) a ventral



FIG. 153.—*A*, transverse section through middle portion of thallus of *M. polymorpha* (× 30); *B*, through marginal portion (more highly magnified). *p*, colourless layer without intercellular spaces *o*, epiderm of upper side; *chl*, chlorophyllous layer; *sp*, stomate; *s*, partition-walls between air-chambers; *u*, lower epiderm; *h*, rhizoids; *b*, leaf-like lamellæ. (After Goebel.)



FIG. 154.—Portion of young receptacle of *M. polymorpha*. *A*, vertical section; *o*, epiderm; *sp*, stomate; *S*, partition-wall separating air-chamber from chlorophyllous cells; *g*, mucilage-cell. *B*, *C*, young stomate; *po*, pore. (After Goebel.)

epidermal layer, from which spring rhizoids and leaf-like lamellæ. The mucilage-passages are especially developed in Fegatella (Radd.) and

Preissia (Cord.), and the thallus of the latter genus has also rudimentary vascular bundles. The stomates which penetrate the epidermal layer of the upper surface of the thallus into the air-chamber-layer are of a struc-



FIG. 155.—*A*, *B*, *C*, young shoot of *M. polymorpha* (slightly magnified) with cupules; *vv*, apical region. *D*, portion of epiderm (more highly magnified). *sp*, stomate.



FIG. 156.—Female inflorescence of *M. polymorpha* seen from the under side. *sr*, radiating branches; *f*, sporogone. (After Goebel.)

ture quite peculiar to this order. Each stomate is formed, according to Leitgeb, by the simple separation of four or more superficial cells, and the subsequent segmentation of these in a direction parallel to the surface.



FIG. 157.—*A*, male plant of *M. polymorpha*. *B*, longitudinal section through inflorescence, *hu* *o*, *o*, openings to antheridial cavities, *a*. *C*, nearly ripe antherid. *D*, two antherozoids (× 800). (After Goebel.)

They are situated in the centre of plates of a rhombic form, consisting of portions of the epidermal layer which overarch large air-cavities. From the base and sides of these air-cavities spring chlorophyllous cells

in rows directed upwards, but not actually reaching the epidermal layer of cells through which the stomates penetrate; while beneath them is the non-chlorophyllous layer, consisting of cells longest in the horizontal direction without intercellular spaces. Each stomate has a number of guard-cells formed by radial cell-divisions. The details in the structure of the stomates differ in the different genera. Leitgeb describes them as of two kinds, simple and canaliculate. The former are epidermal pores situated immediately above the air-chambers; the latter, which occur in Marchantia and Preissia, have the appearance of canals excavated in the surface of the thallus. Some of the rhizoids of Marchantia are characterised by singular internal thickenings to the cell-wall. The peculiar non-sexual organs of propagation of Marchantia, Lunularia (Mich.), and other genera, known as *cupules*, have already been described (figs. 150, 155). A peculiar non-sexual mode of propagation by means of gemmæ occurs in Fegatella (Radd.).



FIG. 158.—Development of archegone of *M. polymorpha* (× 300). *I—V*, before, *VI—VIII*, after fertilisation. *e*, central cell with oosphere; *f*, young embryo; *sl*, lowest cell of axile row; *pp*, perigyne. *IX*, immature sporogone in venter of archegone (× 30); *a*, neck of archegone; *st*, stalk of sporange which contains young spores and elaters. (After Goebel.)

The sexual reproductive organs of the Marchantiaceæ are, in most of the genera, borne on erect branches of the thallus of a peculiar umbrella-like form, which have been variously termed *receptacles*, *discs*, and *inflorescences*. They may be male, female, or bisexual; and, when unisexual, the species may be monœcious or dioœcious. In Fegatella the male inflorescences are oval discs sessile upon the thallus (fig. 152). The inflorescence is generally regarded

as a transformed thalloid axis. The antherids spring from superficial cells of these branches which are depressed in hollows on the upper surface of the disc, and become overarched by the surrounding tissue. With the exception of one section, the Targionieæ, in which they occur at the apex of ordinary shoots, the archegones are borne on the under surface of the female discs, which are always stalked, while the male discs may be either stalked or sessile. The archegones are variously surrounded by *involucres* or *perigynes*. Leitgeb describes the sexual organs as being at first distributed over the surface of the thallus, and becoming subsequently collected into groups or inflorescences, which have at first a dorsal position, but become constantly pushed towards the apex. The mature sporange is usually shortly stalked, and contains *elaters*, which radiate from the centre towards the circumference. It has no central columel. It either dehisces at the apex with numerous teeth, or is four-lobed, or the upper portion becomes detached by an annular fissure as an opercule. The elaters are well developed, and are furnished with several spiral bands, but do not usually appear to take any part in the expulsion of the spores from the sporange.



FIG. 159.—*A*, piece of elater of *M. polymorpha* (magnified). *A'*, a portion more highly magnified. *B*, pitted cell of thallus. *C*, *D*, rhizoids with internal thickenings.

The thallus of many Marchantiaceæ displays remarkable hygroscopic properties, which have their seat in the 'mechanical' layer, *i.e.* the layer of closely packed cells containing but little chlorophyll, which underlies the air-containing assimilating layer. On desiccation this layer contracts greatly, so that the epidermal layer with its stomates is completely protected from further evaporation by the recurved ventral surface covered with brown or violet scales. In this condition the dried-up thallus may retain its vitality for a very long period. The cells of the mechanical layer are frequently occupied by colonies of Nostoc.

Illustrative genera :—Marchantia (L.), Targionia (L.), Fegatella (Radd.), Reboulia (Radd.), Fimbriaria (N. ab E.), Dumortiera (N. ab E.), Plagiochasma (L. & L.), Preissia (Cord.), Lunularia (Mich.).

LITERATURE.

Mirbel—Mém. Acad. Sc., xii., 1835.
Strasburger—Pringsheim's Jahrb. wiss. Bot., 1870, p. 409.
Vogt—Bot. Zeit., 1879, pp. 729 and 745.
Goebel—Arb. Bot. Inst. Würzburg, 1880, p. 529.
Leitgeb—Sitzber. Akad. Wiss. Wien, 1880, pp. 40 and 123; and Die Marchantieen, 1881.
Prescher—(Mucilage-receptacles) Sitzber. Akad. Wiss. Wien, 1882.
Mattirolo—(Hygroscopic Properties) Malpighia, ii. 1888, p. 181.

FOSSIL MUSCINEÆ.

No remains have been found earlier than the Tertiary formations which appear to belong to Muscineæ. Here and in the Quaternary beds remains or impressions occur which have been referred to various families of Musci and Hepaticæ, including leaves of a single species of Sphagnum and a single moss-capsule. The leaves of Jungermanniaceæ are not uncommonly found enclosed in amber.

THIRD SUBDIVISION AND CLASS IX.

CHARACEÆ.

The true position of this small group in a natural system of classification has been a subject of much controversy. By some writers of high authority it is regarded as occupying the highest place among green Algæ. On the other hand, although without any lignification of their tissue, the Characeæ display, in the structure of their vegetative organs, a distinctly higher type of structure than the Thallophytes, in the distinct differentiation of the plant into a primary axis or stem, and secondary axes or branches; but the branches are similar in structure to the primary stem. They are, in fact, Cormophytes rather than Thallophytes; and it seems best to retain them as a distinct subdivision intermediate between the Muscineæ and the highest Algæ.

The plant is acrogenous, growing by means of an *apical cell* contained in an apical bud; the main stem has indefinite apical growth, the branches increasing by definite apical growth. The branches and the organs of sexual reproduction grow in the axils of other lateral organs of more simple structure, which are usually termed *leaves*; those that subtend the reproductive organs being by some writers described as *bracts* or bracteoles. In all the Characeæ these appendicular organs spring in whorls from well-defined *nodes* of the primary stem, imparting the peculiar habit to the plants by which they are distinguished from nearly all other Cryptogams. Each internode consists, in the Nitelleæ, of a single very large cell extending along its whole length, and many times longer than broad. In the majority of the Chareæ this internodal cell is invested by a layer of similar elongated cells of much smaller diameter arranged spirally round it, collectively known as the *cortex*, and giving the stem the appearance of being spirally striated. Each node consists, in the corticated species, of a single layer or plate of small cells from which the cortex is derived. From the nodes spring the whorls of branches and their subtending leaves. The branches are altogether similar in structure to the primary axis. The leaves have also, in the Chareæ, a simple cortical layer, with the exception of the apex, where

the large terminal cell is exposed. In addition to the leaves there spring, from the basal nodes of some species of Chara (L.), other leaf-like structures known as *stipules*, one, two, or three in connection with each leaf. The stipular cells are always undivided by septa, and arise as papillæ on the cortical cells.

FIG. 160.—*Chara fragilis* Desv. (natural size).

The cortex of the stem and branches is developed out of the nodal plate of cells; the upward and downward prolongations from the nodes usually meeting about the middle of each internode, where they dovetail into one another. These cortical internodal cells do not, however, like the axial cells, remain entire; they divide, both transversely and longitudinally, into three parallel rows of cells, the central row of each series being somewhat elevated into a ridge. The mode and extent of development of the cortical cells vary according to the species. The number of leaves in a whorl is usually from four to ten. At the lower part of the main stem the internodes are shorter, and from the nodes spring *rhizoids* or rooting filaments which serve to fix the plant in the soil, consisting of long hyaline nearly undivided tubes, which grow obliquely downwards, and lengthen only at their apex. The rhizoids are always trichomic, springing from superficial cells.

FIG. 161.—Fertile branch of *C. hispida* L. (magnified).

The nearly hemispherical apical cell of the terminal bud of the stem first divides by a transverse wall into a new apical cell and a disc-shaped segment-cell. Each segment then again divides by a wall parallel to the first; the lowest of these does not again divide, but develops into the axial internodal cell, while the upper one undergoes vertical division, and becomes a node. Each successive whorl on the main stem alternates with those immediately above and below it, so that the oldest leaves of a

whorl, which subtend the branches, are arranged in a spiral line running round the stem; but this is not the case with the branches or secondary axes, where the members of contiguous whorls are superposed.

The Characeæ exhibit in an especially clear and beautiful manner the phenomenon of *cyclosis*, or rotation of the protoplasm (see fig. 163). The best objects for observation are the large internodal cells of Nitella (Ag.), the apical cells in the leaves of Chara, or some of those belonging to the reproductive organs, especially to the 'manubria.' The cell first of all develops vacuoles in its protoplasm, which coalesce into a single



FIG. 162.—Longitudinal section through bud of *C. fragilis*, showing apical cell, *t*, and segments, *g*, *b*. *A*, cells empty. *B*, with cell-contents, granular protoplasm, chlorophyll-grains, and vacuoles. *C*, with cell-contents contracted by iodine (× 500). (After Sachs.)

large sap-cavity. The outermost thin parietal layer of protoplasm, in which are imbedded most of the grains of chlorophyll, remains motionless; within this motionless lining is a thick layer of protoplasm, in which a regular current gradually sets up, up one side of the cell and down the other; the boundary between the two currents being marked by hyaline bands entirely destitute of chlorophyll, the *neutral zones*, in which no movement is visible. The direction of the rotating movement in each cell stands in a definite relation to that of all the other cells of the plant. From time to time the movement ceases, and then begins again in the opposite direction. Before the rotation commences the cell-nucleus has

usually broken up into a number of fragments. The current is most rapid next to the stationary parietal layer, and becomes gradually slower towards the interior. As the cell grows the rotating protoplasm becomes differentiated into a watery and a less watery denser portion, the former having the appearance of a hyaline cell-sap, in which the latter floats in the form of larger or smaller roundish lumps. Since these denser bodies are passively swept along by the clear rotating protoplasm, the appearance is presented as if the cell-sap caused the rotation. Together with the denser lumps of protoplasm of less regular form, there are also a number of globular masses carried along in the current, which are covered with delicate protoplasmic spines or cilia; their nature and function are involved in obscurity.

Owing to the large size of the cells and the distinct differentiation of the nucleus, the internodal cells of the main axis of Chara and Nitella, as well as the apical cells of the leaves, have been largely used for following the complicated processes connected with cell-division and the division of the nucleus. Schmitz describes the process as one of 'direct division of the nucleus,' Treub and Strasburger as one of 'fragmentation;' Johow differs in some respects from all previous observers; Cagnieul (Bull. Soc. Bot. France, 1884, p. 211) finds the process especially easy to follow in the mother-cells of the antherozoids. Schaarschmidt (Bot. Centralblatt, vol. xxii., 1885, p. 1) describes peculiar cell-wall thickenings and grains of 'cellulin' in Chara hispida (L.).

The Characeæ do not produce spores, *i.e.* single non-sexual propagative cells; but are multiplied non-sexually in three different ways, the nodes being always the place of origin of the propagative cells. (1) Chiefly in Lychnothamnus stelliger (A.Br.), but also in C. hispida, C. aspera (Willd.), and Lamprothamnus alopecuroides (A.Br.), structures called *bulbils* or 'amylum-stars' are formed, agglomerations of cells developed round the larger internodal cells at the level of the nodes; they are of beautiful regularity, and are densely filled with starch and other food-materials. On germinating they appear to produce at first other bulbils, and from these a new plant. (2) Chara fragilis and other species produce, on old hibernating or on cut nodes, in the axils of the leaves, peculiar branches known as *gymnopodal shoots*, which differ from the ordinary branches in the partial or entire absence of the cortex in the lowest internode and in the first whorl of leaves. The cortical branches which descend from the first node become detached, bend upwards, and propagate themselves. (3) Also on C. fragilis, Pringsheim describes the occurrence of 'pro-embryonic,' or more properly of *prothalloid branches*. These also spring from the nodes of the main axis, but differ essentially from the ordinary branches, presenting a similar structure to the pro-

thallium or 'pro-embryo' which proceeds from the germinating oosperm. No mode of vegetative propagation is known in the other genera.

The sexual reproductive organs of the Characeæ, the male *antherids* and the female *archegones*, are visible to the naked eye as minute orange-red globes and elliptical green bodies springing from the nodes in the axils of leaves or bracts. The antherids are globular bodies, of a bright red colour when mature, from ½ to 1 mm. in diameter, morphologically the terminal cell of a leaf or lateral leaflet. The moderately thick wall of the antherid is made up of eight flat disc-shaped cells called *shields*, four of which, situated round the distal pole of the ball, are triangular, while the four situated round the base are four-sided. On their inner face there lies a layer of chlorophyll-grains, which eventually turn red, while the outer face is clear and transparent; the walls of these cells are folded inwards at the edge where they meet. From the centre of the inner face of each shield a cylindrical cell, termed a handle or *manubrium*, projects inwards nearly to the centre of the globe. The antherid is supported on a short flask-shaped *pedicel-cell*, which also projects into the interior between the four lower four-sided shields. At the free end of each of the eight manubria is a roundish hyaline-cell, the head-cell or *capitulum*. These twenty-five cells—viz. the eight shields, eight manubria, eight capitula, and the pedicel-cell—constitute the



FIG. 163.—*Nitella flexilis* Ag. *A*, nearly ripe antherid subtended by two bracts showing direction of protoplasm-currents, and neutral zone, *i*. *B*, manubrium, with capitulum, secondary capitula, and whip-like filaments. *C*—*F*, antheridial filaments, showing formation of antherozoids. *G*, antherozoids (*C*—*G* × 550). (After Sachs.)

framework of the antherid. Each capitulum bears six smaller cells, or *secondary capitula*; and from each of these grow four long whip-shaped filaments, bent into a number of coils and filling up the interior of the globe. The manubrium, capitulum, secondary capitula, and whip-shaped filaments, bear a resemblance to a many-thonged whip. The number of these filaments in an antherid amounts to nearly 200, and each filament is divided by transverse septa into from 100 to 200 small disc-shaped cells. The protoplasm in each of these antheridial cells becomes gradually transformed into an *antherozoid* strongly resembling the corresponding organ in Muscineæ rather than in Thallo-



FIG. 164.—*A*, portion of branch of *C. fragilis*; *a*, antherid; *S*, archegone; *c*, crown; *β′*, *β″*, bracts (× 50). *B*, a young antherid; *SK*, young archegone (× 350). (After Sachs.)

phytes. It is a slender thread of protoplasm coiled spirally like a corkscrew, somewhat thickened at the posterior end, and bearing at its pointed anterior end two long fine cilia. The number of antherozoids in an antherid is, as will be seen, from 20,000 to 40,000. When ripe, the eight shields fall apart, and the antherozoids escape from their mother-cells, and move about rapidly in the water by means of their vibratile cilia. This appears generally to take place in the morning, the antherozoids swarming about for some hours till the evening.

The Characeæ are either monœcious or diœcious. In the former case the male and female organs are formed in close juxtaposition on the same node, the archegone being somewhat below the antherid in Nitella, above it or by its side in Chara. The archegones, like the antherids, are metamorphosed leaves. When ready for fertilisation, the archegone has a longer or shorter ovoid form, and is borne on a short *pedicel-cell*. In the interior is an axial row of cells enveloped by five tubes, which are at first straight, but are afterwards coiled spirally round the axial row. The lowest portion of each of these tubes is an elongated unsegmented cell; while at the upper part one or two very small cells are segmented off. In Nitella each of the terminal cells again divides into two by a vertical septum. The five terminal cells of Chara and the ten terminal cells of Nitella are not twisted, and form together the *crown*. When the archegone is ready for impregnation these crown-cells separate from one another, forming the *neck*, and leaving an open passage down to the axial row. This *apical cavity* is, however, very nearly closed below by a *diaphragm* formed by the projecting inwards of the five neck-cells, through which there is only a very narrow opening for the entrance of the antherozoids. The apical cell of the axial row is much larger than the rest, and is the female or *germ-cell*, corresponding to the central cell in the archegone of the higher Cryptogams. It is filled with protoplasm, oil-drops, and starch-grains; its apical portion, the *apical papilla*, or receptive spot, containing only hyaline protoplasm. Between the apical cell and the pedicel-cell of the archegone, there is in Chara only a single cell, in Nitella a group of cells, the 'Wendungszellen.' Before fertilisation the crown is a compact structure covering the apical cavity; but when the archegone is ready for impregnation a small aperture is formed in its



FIG. 165.—*A*, fertile branch of *Nitella flexilis* (natural size); *i*, internode; *b*, branches. *B*, upper portion of fertile leaf, *b*; *K*, node; *nb*, bracts; *S*, young archegone. *C*, older leaf with two bracts; *a*, antherid; *S*, spermocarp. *D*, half-mature spermocarp (highly magnified). (After Sachs.)

centre, through which the antherozoids force their way, and finally enter the apical cell by the deliquescence of the upper portion of its cell-wall, and coalesce with the apical papilla. The whole contents of the apical cell may be regarded as the oosphere.

Impregnation causes at first very little external change in the structure of the female organ. The protoplasm of the oosphere, now invested with a cell-wall and transformed into an *oosperm*, gives place to starchy or oily matter; the walls of the enveloping tubes which lie next it increase in thickness and hardness, and the oosperm thus becomes invested in a hard black shell or *pericarp*. The structure thus formed, the so-called 'fruit' or *spermocarp* of the Characeæ ultimately becomes detached, falls into the mud at the bottom of the water, and there germinates in the next spring.

When the spermocarp germinates, the oosperm first divides into



FIG. 166.—*A–D*, stages in development of archegone of *N. flexilis*. *b*, apex of fertile leaf; *x*, 'Wendungszellen;' *K*, crown (× 300). (After Sachs.)

FIG. 167.—Calcareous spermocarp of *C. hispida* (magnified).

three cells, a large basal and two apical cells, the former apparently serving the purpose of supplying with nutriment the young plant which proceeds from the latter; and these three cells may be said together to constitute the *embryo*. From one of the two apical cells proceeds a long hyaline unseptated filament, commonly called the *primary root*, by means of which the young plant is attached to the soil. The other of the two apical cells develops into a hypha-like filament, consisting at first of a single row of cells with limited apical growth, and called by some writers the 'pro-embryo,' or more correctly the *prothallium*. In this prothallium are developed two *primary nodes* at considerable distance from one another, and separated by a very long internode. From the lower of these two primary nodes there springs a whorl of *rhizoids*, which soon

usurp the functions of the primary root. The upper of the two nodes is still at some distance from the apex of the prothallium, this apical portion above the upper node consisting of a few much shorter cells. From this upper node is developed the new plant. It is divided by longitudinal septa into two inner and six or eight peripheral cells. The peripheral cells ultimately become rudimentary leaves, which do not, however, form a true whorl. In the midst of them appears a bud, or growing point, developed from one of the inner cells, from which springs the new stem, in a direction nearly at right angles to that of the prothallium. At present the formation of the prothallium has been observed only in the genus Chara.

FIG. 168.—Germination of *C. fragilis*. *sp*, spermocarp; *w'*, first root; *i*, first internode of prothallium; *d*, first node; *w''*, rhizoids; *g*, second elongated internode of prothallium; *g*, second node with first whorl of leaves; *pl*, apical portion of prothallium (× 4). (After Pringsheim.)

A remarkable instance of *parthenogenesis* has been recorded in Chara crinita (Wallr.). The species is dicœcious, and male plants are extremely rare. On the female plants the oospheres develop into oosperms without apparently any possibility of their having been impregnated; and the spermocarps thus formed germinate in the ordinary way.

The Characeæ consist of only a comparatively small number of species, but some of them very abundant, growing submerged in deep or in shallow, in stagnant or in running, or occasionally in brackish water. Several species are grown with great facility in fresh-water aquaria, where they multiply very rapidly. The presence of certain species may be detected by the fœtid odour of sulphuretted hydrogen given off when decaying. Phipson (Compt. Rend., lxxxiv., 1879, pp. 316, 1078) attributes this odour to the presence of a special substance which he calls *characin*. The typical genus Chara is distinguished by its power of extracting calcium carbonate from the water in which it grows, the whole plant becoming thus covered with a calcareous incrustation, which frequently renders it difficult to make out the structure. Hence the family has acquired the popular names of 'brittleworts' and 'stoneworts.' Nitella translucens (Ag.) sometimes forms enormous mat-like masses at the bottom of ponds.

The systematic position of the Characeæ has been a matter of much

controversy. In habit and in general appearance they resemble the Algæ, among which they are placed by the majority of writers. But in some important points of structure they differ so widely from all known families of Algæ, that a true estimate of their relationships appears to require their location in a distinct subdivision by themselves. With the exception of the Fucaceæ and the Conjugatæ, the Characeæ stand alone among the larger groups of Cryptogams in the entire absence of true spores. Seeing that the oosperm germinates directly in the soil, the embryo which results from its first divisions developing directly into the new plant, there is no 'alternation of generations' in any accurate sense of the term. From those classes where a true alternation of generations attains its fullest development, the Muscineæ and Vascular Cryptogams, the Characeæ differ in the complete suppression of the sporophyte-generation; while Phanerogams (at all events those Angiosperms which are destitute of endosperm) deviate, on the other hand, in the suppression of the oophyte-generation. In the investment of the oosperm with a lignified pericarp directly without any previous breaking up into carpospores, the Characeæ again differ essentially from all classes of Algæ.

The Characeæ are divided into two orders, viz.—

1. CHAREÆ.—Stem and branches usually corticated and calcareous; leaves usually with one or two stipules at their base; antherids usually solitary on each node; crown always five-celled; pericarp often calcareous. *Genera*: Chara (L.), Lamprothamnus (A. Br.), Lychnothamnus (Leon.).

2. NITELLEÆ.—Stem and branches not corticated nor calcareous; leaves without stipules; archegones often clustered; crown always ten-celled; pericarp not calcareous. *Genera*: Nitella (Ag.), Tolypella (A. Br.).

LITERATURE.

Göppert and Cohn—Bot. Zeit., 1849, pp. 665 *et seq.*
Braun—Monber. Berlin Akad. Wiss., 1852, p. 220; and 1853, p. 45; and (Parthenogenesis) Abhandl. Berl. Akad. Wiss., 1856, p. 337.
Thuret—Ann. Sc. Nat., xvi., 1851, p. 18.
Montagne—Ann. Sc. Nat., xviii., 1852, p. 65.
Nägeli—Beiträge zur wiss. Bot., ii., 1860, p. 61.
Pringsheim—Jahrb. wiss. Bot., 1863, p. 294.
Braun—Conspectus systematicus Characearum europæarum, 1867; and Abhandl. Berlin Akad. Wiss., 1882.
De Bary—Monber. Berlin Akad. Wiss., 1871, p. 227; and Bot. Zeit., 1875, pp. 377 *et seq.*
Bennett—Journ. of Bot., 1878, p. 202.
Groves—Journ. of Bot., 1880, p. 97.

Müller—Bull. Soc. Bot. Genève, 1881.
Johow—Bot. Zeit., 1881, pp. 729 *et seq.*
Nordstedt—Hedwigia, 1888, p. 181.
Allen—Characeæ of America, 1888.

FOSSIL CHARACEÆ.

In various strata, commencing with the Cretaceous, remains known as *gyroliths* are found, sometimes in great abundance, which appear to be the petrified pericarp of the spermocarp of Characeæ. Upwards of forty species have been described, some of them closely resembling existing forms.

FOURTH SUBDIVISION.

ALGÆ.

The degree of affinity between the small group of Characeæ and the very large group of Algæ is, as has already been mentioned, a point on which the best authorities are not agreed. But, in passing from one to the other, we finally cross the line which separates the Cormophytes from the Thallophytes. From this point we have to do exclusively with plants whose vegetative organs are in no sense differentiated into axial and appendicular, and which further contain no true vessels and no woody tissue. On their lower limit there is no sharp line of demarcation between the Algæ and the chlorophyllous Protophyta, but the consideration of Algæ as a group by themselves, distinct from Fungi, we regard not only as convenient, but as also most in accordance with their probable affinities.

Within the limits above mentioned, the degree of complexity in the structure of Algæ is very various, and the different types will be best described under the separate families. In their vegetative structure we may recognise three types: the elaboration in the development of a *single cell*, the loose association of cells into a family or *cœnobe*, and the close aggregation of cells into a *filament* or a *thallus*. To the latter belong all the higher families, and in some of these we see indications of the various kinds of tissue found in vascular plants. The higher forms, consisting of a well-developed thallus of large size, in which the cells are associated with one another in all three directions, are almost exclusively marine, and include the whole of the organisms popularly known as *seaweeds*. In the larger forms the plant is attached to the substratum—a rock, stone, or other large alga—by a root-like organ of attachment known as the *disc*. The attachment is, however, always superficial, and the organ takes no part in the absorption of nourishment for the plant. The organ may result from the repeated division of a single cell, or it may be more complicated, being formed out of the termination of the downward growth of cortical rows of cells. In nearly all fresh-water Algæ the single cell, the cœnobe, or the filament is en-

closed in a more or less strongly developed gelatinous sheath. The greater number of families exhibit both a sexual and a non-sexual mode of reproduction, though in some cases one or the other mode has not yet been detected. In the great majority of families the non-sexual propagating bodies are motile cells or *zoospores*, minute masses of protoplasm formed singly from the whole contents of a cell by rejuvenescence, or more often in large numbers by free-cell formation, destitute of a true cellulose membrane, but containing protoplasm and a contractile vacuole, and provided with two or sometimes a larger number (rarely only one) of vibratile cilia, by means of which they move about actively for a time, then come to rest, excrete a cellulose membrane, and develop into a new plant. In one class only, the Florideæ, are the zoospores replaced by non-motile *tetraspores*; in the Conjugatæ and Fucaceæ they are altogether wanting. The simplest form of sexual reproduction is that of *conjugation*, or the coalescence of two comparatively undifferentiated masses of protoplasm. These masses of protoplasm may be either the contents of stationary cells, which are nearly or quite alike, as in the Conjugatæ, or they may be motile ciliated bodies indistinguishable from zoospores—*zoogametes*—or they may be distinguished from the true zoospores by their smaller size. From the conjugation of zoogametes there is a gradual transition through intermediate stages to a true sexual process, the impregnation of a stationary *oosphere* by a motile *antherozoid*, usually much smaller than the oosphere, the result being the production of an *oosperm* by the encysting of the oosphere in a coat of cellulose. In the higher families the oospheres and antherozoids are formed in special cells or organs, known as *oogones* and *antherids* respectively. In the Florideæ the process displays very great complication; the structure in which the oosphere is formed is known as the *carpogone*; the fertilised oosphere is the *carposperm*, which often breaks up into *carpospores*. In this class also the antherozoids are replaced by motionless protoplasmic bodies known as *pollinoids*. Multiplication by the simple *fission* of individuals, by the detachment of *gemmæ*, or buds, and by the encysting of special cells or masses of cells into *cysts*, also occur. In the green Algæ (Confervoideæ heterogamæ and isogamæ and Conjugatæ) single non-motile cells which become detached for the purpose of propagation are termed by Wille *akinetes* when they are formed without rejuvenescence, *aplanospores* when formed by rejuvenescence. The former occur in Trentepohlia (Mart.), Conferva (L.), and Ulothrix (Ktz.), as well as in the Nostocaceæ and Rivulariaceæ, the latter in the Confervaceæ. The two kinds pass into one another, and akinetes into vegetative cells, by insensible gradations.

Any classification of Algæ which attempts to follow the lines of affinity—in other words, any natural system of classification—must be based on a consideration of both the vegetative and the reproductive organs. All the families of Algæ appear undoubtedly to have sprung from the PROTOCOCCOIDEÆ, and their further development has taken place in three directions—the perfection and differentiation of the individual cell, the association of cells into cœnobes, and cell-division. The production of cœnobes may be supposed to start from such forms as Botryococcus among *Protococcaceæ*; the first step in the development of the CŒNOBIÆ being the *Sorastreæ*, including Sorastrum, Cœlastrum, and Selenastrum, motile colonies of non-ciliated cells, with no known production of zoospores. The series attains a much higher development in the *Pandorineæ*, including Pandorina, Gonium, and Stephanosphæra, where reproduction is effected by the conjugation of zoogametes. Simple organisms like Chlamydomonas and Chlamydococcus, consisting merely of conjugating zoogametes, are possibly retrogressions from the higher forms, though they may also be stages in a direct ascent from Protococcus. Eudorina, with a rudimentary differentiation of antherozoids and oospheres, unquestionably indicates the line of development of *Volvox*, in which this differentiation is more strongly developed. In Volvox we have the culmination of the attempt of nature to evolve higher organisms out of cœnobes. *Hydrodictyon* is probably an aberrant member of this group, and the *Pediastreæ* are more or less nearly related to them.

From the *Eremobiæ* the fuller development of the individual cell has advanced a further stage in the MULTINUCLEATÆ, composed of the *Siphonocladaceæ* and *Siphoneæ*, and characterised by each individual consisting of an enormously developed cell, often ramifying greatly and attaining gigantic dimensions, and possessing several, often a very considerable number, of nuclei. In the Siphonocladaceæ the only known mode of reproduction is by the conjugation of zoogametes; and Botrydium displays a distinct affinity with Botrydina among the Eremobiæ. The Siphoneæ or Cœloblastæ, represented by Vaucheria, are a higher development of the same series, in which true sexual organs, oogones and antherids, are formed in addition to non-sexual zoospores; and in this genus culminates the striving after a higher development in the elaboration of a single cell.

A rudimentary cell-division is exhibited in the *Nostochineæ* among Protophytes, but accompanied by other conditions which prevented its full success there. Where cell-division originated in the Protococcoideæ is not clear, probably in the Eremobiæ; we find it already fully developed in the CONFERVOIDEÆ ISOGAMÆ, the members of which consist of a

single unbranched or branched filament of cells, the only known modes of multiplication being the conjugation of zoogametes and the direct germination of larger zoospores. In the lowest two classes, the *Chroolepideæ* and *Ulotrichaceæ*, embracing a very small number of genera, the filament is usually unbranched; in the two higher, the *Confervaceæ* and *Pithophoraceæ*, further vegetative activity is displayed in the copious branching; and in the former we have an indication of affinity with the Multinucleatæ in an occasional plurality of nuclei.

The exact course of evolution from the isogamous Confervoideæ is obscure, but it would appear to have taken place in three distinct lines. The first of these, which evidently came to an abrupt conclusion, is the Conjugatæ, consisting of the *Zygnemaceæ*, *Mesocarpeæ*, and *Desmidieæ*, a well-marked and sharply differentiated group with no near affinities. The first two orders, consisting of unbranched filamentous forms, are probably derived directly from the Confervoideæ, although the change in the mode of reproduction is very abrupt. The production of zoospores is entirely suppressed, and they are reproduced solely by the conjugation of cells belonging to the same or to different individuals. The Desmidieæ must then be regarded as a group adapted, by a certain amount of retrogression in both vegetative and reproductive characters, to life in shallow water; and derived, through such filamentous genera as Desmidium and Hyalotheca, from Zygnemaceæ with lateral conjugation. By some writers the Diatomaceæ are associated with the Desmidieæ; our reasons for placing them among the Protophyta will be given hereafter. The mode of reproduction by conjugation attains its climax in the Mesocarpeæ.

The second line of descent is that of the brown seaweeds. In the Phæosporeæ we have every shade of transition in the mode of reproduction from isogamous to heterogamous. The typical Phæosporeæ, such as Punctaria and Ectocarpus, are characterised by the possession of two kinds of zoosporange, unilocular and multilocular. The zoospores produced in these two kinds of zoosporange present no difference in size or form; but those from the unilocular sporanges appear in all cases to germinate directly, while those produced in the multilocular sporanges are sometimes zoogametes with sexual functions. In some families one or the other kind of zoosporange is suppressed. In the *Ectocarpaceæ* and some other genera we have a mode of reproduction closely resembling that in the isogamous Confervoideæ, except in the greater differentiation of the cells which become zoosporanges, a conjugation of zoogametes which are to all appearance exactly alike, though a slight differentiation is exhibited in the fact of one of them coming to rest and partially losing its cilia before conjugation takes place. In the *Cutleriaceæ* the differ-

entiation is more complete ; the male and female swarm-cells are produced either on the same or on different individuals ; the female are much larger than the male, and come perfectly to rest, entirely losing their cilia before being impregnated by the former. In the *Dictyotaceæ* the differentiation is carried still further, and the female reproductive bodies are from the first motionless oospheres not provided with cilia. Several families of Phæosporeæ exhibit reduction or degradation of the vegetative structure ; and among these we are disposed to place the small group of *Syngeneticæ*, consisting of but two genera, Hydrurus and Chromophyton, which resemble one another in but few points except the possession of a brown endochrome. In the former genus the propagative bodies are reduced to non-ciliated masses of brown protoplasm, which germinate directly without impregnation ; in the latter the vegetative structure is almost entirely suppressed ; the propagative bodies are uniciliated masses of protoplasm of two kinds, but without any observed process of conjugation. The step from the Dictyotaceæ to the FUCACEÆ is an easy one. In the highest type of brown seaweeds, such as Fucus or Durvillæa, with their typical heterogamous or 'oogamous' reproduction, consisting in the impregnation of a comparatively large oosphere by a number of minute antherozoids, we have the highest attainment of this series of development.

The third line of descent from isogamous Confervoideæ is a much more direct one, to the CONFERVOIDEÆ HETEROGAMÆ, including the three orders *Sphæropleaceæ*, *Œdogoniaceæ*, and *Coleochætaceæ*. In the first of these, which comprises only a single species, we have a distinct differentiation of the male and female reproductive cells, the latter having now become permanently quiescent ; but still a strong reminiscence of the Confervaceæ in the multinucleated cells. The Œdogoniaceæ exhibit a distinct advance in vegetative structure, and still more in the cells which contain the male and female reproductive bodies being, for the first time in this series, differentiated into antherids and oogones respectively. Between the Œdogoniaceæ and the Coleochætaceæ we have an evident connecting link in Bulbochæte ; but the typical genus Coleochæte presents a singular reduction of the non-reproductive portion of the thallus from a filament to a single plate of cells. The mode of sexual reproduction has now attained a much higher degree of complexity. The oogone is surmounted by a tubular appendage, the trichogyne, through which the motile antherozoids find their way to the oosphere in order to impregnate it. The fertilised carpogone, as it is now called, then becomes invested by a cortical layer of cells, forming the complex body known as the sporocarp.

The Coleochætaceæ lead up directly to the highest type of structure attained by Algæ, the FLORIDEÆ or red seaweeds, a well-defined and

natural group, though exhibiting remarkable variety in the degree of development of the sexual organs. So striking is the resemblance in the mode of impregnation in the most highly developed genera of Florideæ, such as Callithamnion, Dudresnaya, or Corallina, to that in Coleochæte, that it is scarcely possible to doubt the direct descent of one from the other ; the chief difference is in the replacement of motile antherozoids by pollinoids which have no active power of motion. The process of fertilisation is the most intricate which occurs among Thallophytes, and presents a remarkable forecast, so to speak, of the mode afterwards elaborated in Flowering Plants, though only after a very long interval, comprising the entire evolution of the Muscineæ and Vascular Cryptogams. With the loss of motility of the male reproductive cells is also correlated a corresponding loss of motility of the non-sexual reproductive cells or tetraspores. In the higher families of the Florideæ we have also the highest development of the organs of assimilation found among Thallophytes. If the view is correct that the higher Florideæ are derived directly from the Coleochætaceæ, it follows that we must regard all the less highly developed families of this group as retrogressions from the parent type ; and this view appears to offer the most probable explanation of the true position of some aberrant forms. In the *Helminthocladiaceæ* and *Squamariaceæ* the degeneration is exhibited solely in the less perfect development of their thallus or vegetative structure. In the *Lemaneaceæ* this is accompanied also by a simpler structure of the sexual organs. But here, as well as in Batrachospermum, we have the first rudimentary appearance of a phenomenon resembling that known as 'alternation of generations,' which plays so important a part in the Vascular Cryptogams, and which may possibly indicate the genesis of the Muscineæ. In the *Porphyraceæ* we find a reduction of the thallus to a simple filament or plate of cells, accompanied by only a rudimentary development of both carpogones and trichogynes, and a limited reversion to motility in the tetraspores. Regarding the Porphyraceæ as exhibiting retrogression from the more complicated Florideæ, rather than as the lowest member of an ascending series, it is difficult to resist the conclusion that the *Ulvaceæ* are derived from the Porphyraceæ by further retrogression, displayed in the entire suppression of antherids and oogones, and a reversion to the primitive conjugation of zoogametes. In their vegetative structure the Ulvaceæ differ widely from the isogamous Confervoideæ, with which they are usually associated, while the close resemblance between Ulva and Porphyra is obvious. In the Florideæ the Algæ attain their highest type of development.

By far the larger number of Algæ grow entirely immersed in water, running or stagnant, fresh, brackish, or salt ; some float on the surface without any attachment ; others are found on moist soil, among moss,

&c. The whole of the marine vegetation of the globe, with the exception of a very small number of species of Flowering Plants, belongs either to the Algæ or to the chlorophyllous Protophytes. They vary in size from the microscopic Desmidieæ and Pediastreæ to that of shrubs or trees in the case especially of some genera of Laminariaceæ and Fucaceæ ; and in these classes, as well as in the Florideæ, we find a rudimentary differentiation, not only of tissues, but of organs, which leads the way to the much more complete development in the higher classes of the vegetable kingdom. Fresh-water Algæ are, with very few exceptions (species of Bangia, Hildenbrandtia, Lithoderma, Hydrurus, &c.), green. Among marine Algæ there are many genera of green seaweeds, belonging chiefly to the families Confervaceæ, Siphonocladaceæ, and Ulvaceæ ; but these mostly grow in shallow water. As regards all organisms growing in deep seas, it appears to be essential to them that the green colour of the chorophyll should be masked by a coloured pigment, red in the case of the Florideæ, brown in those of the Phæosporeæ and Fucaceæ ; the nature of these pigments will be discussed under the separate families. A few of the smaller species, belonging to the Coleochætaceæ, Chordariaceæ, and Squamariaceæ, grow attached to stones, larger Algæ, or other marine objects, as flat discs, gelatinous cushions, or calcareous incrustations, and the deposition of lime takes place to a much larger extent in the corallines. The proportion of ash to the organic constituents is much larger in marine than in land or fresh-water plants, seaweeds having the power of extracting from the salt water large quantities of the soluble salts contained in it. The larger species of Fucaceæ and Laminariaceæ are largely used in the north of Europe for manuring the land and for foddering cattle ; and in former times the manufacture from their ashes of kelp and barilla was an important industry. They are also an important commercial source of iodine. From the quantity of gelatine contained in their thallus, some species of Ulvaceæ, Porphyraceæ, Fucaceæ, and Laminariaceæ are also occasionally used as articles of food or for medicinal purposes.

Literature.

Greville—Algæ, in Scottish Cryptogamic Flora, 1823–28 ; Algæ Britannicæ, 1830.

Kützing—Phycologia generalis, 1843 ; Tabulæ Phycologicæ, 1845–69 ; and Species Algarum, 1849.

Harvey—Phycologia Britannica, 1846–51 ; Nereis Australis, 1847–49 ; British Marine Algæ, 1849; Nereis Boreali-Americana, 1851–58; Phycologia Australica, 1858–63.

Hassall—British Fresh-water Algæ, 1845.

Nägeli—Die neuern Algensysteme, 1847.

Agardh—Species, Genera, et Ordines Algarum, 1848–80 ; Till Algernes Systematik, 1872–87.

Thuret—Antheridies des Cryptogames, 1851 ; Zoospores des Algues, 1851 ; and Etudes Phycologiques, 1878.

Landsborough – Popular Hist. of British Seaweeds, 1851.
Pringsheim – Ueber Befruchtung u. Keimung der Algen, 1855.
Gray—British Seaweeds, 1867.
Wood—Fresh-water Algæ of North America, 1873.
Bornet & Thuret—Notes Algologiques, 1876-80.
Falkenberg—Die Algen, in Schenk's Handbuch der Botanik, vol. ii., 1881.
Hauck – Die Meeresalgen, in Rabenhorst's Kryptogamen-Flora Deutschland, 1883-85.
Schmitz – Die Chromatophoren der Algen, 1882 ; and Journ. Micr. Soc., 1883, p. 405.
Cooke—British Fresh-water Algæ, 1884.
Gay—Bull. Soc. Bot. France, 1886, Sess. Extraord., p. 21.
Bennett – Journ. Linn. Soc., xxiv., 1887, p. 49.
Wolle – Fresh-water Algæ of the United States, 1887.
Wille—(Resting-spores) Pringsheim's Jahrb. wiss. Bot., 1887, p. 492.
Stroemfelt—(Attachment-disc) Bot. Centralbl., xxxiii., 1888, pp. 381 & 395.

Class X.—Florideæ.

This large family—known also as Rhodosporeæ and Rhodospermeæ—exhibits the highest type among Algæ in the mode of sexual reproduction, and also possibly in the development of the vegetative organs. It consists almost entirely of seaweeds, including all the red and purple kinds. A few species only, belonging to the genera Hildenbrandtia (Nard.), Batrachospermum (Bory), Lemanea (Bory), Bangia (Lyng.), and a few others, grow in fresh water. Some of these are green, but the great majority of the Florideæ are of a bright red colour, varying with purple, brown, yellowish, or dirty white. The 'fronds' do not attain nearly the size of those of the Fucaceæ and Laminariaceæ, but they are often of delicate texture and finely divided, rendering them the most beautiful of our seaweeds.

The thallus varies within very wide limits in its degree of development. In a few genera, such as Callithamnion (Lyng.), it consists of distinct filaments of cells which are almost always branched ; in others, as Porphyra (Ag.), Hildenbrandtia, and Cruoria (Fr.), of a flat plate of cells, composed of only a single or of several layers ; in the fresh-water genus Batrachospermum, of an axis with beautifully regular whorls of branches ; while in most seaweeds it constitutes a massive parenchyme, or the filaments are held together by a more or less dense gelatinous envelope. Growth takes place, in the majority of cases, by means of a single apical cell, but this is often followed by a strong intercalary growth. The apical cell is not three-sided, as in Vascular Cryptogams, but is either wedge-shaped, dividing by walls which incline alternately to the right and left, or it divides by nearly parallel walls. In some forms, however, especially the prostrate Melobesiaceæ and Squamariaceæ, and

in the Nemalieæ, the apical cell appears to be replaced by a group of equivalent cells. Wille distinguishes six types of Florideæ as respects their mode of growth; in four of these groups growth takes place from a single apical cell; in two from an apical mass of cells, with peripheral growth. In the so-called *polysiphonous* seaweeds, such as Polysiphonia (Grev.), a transverse section of the 'frond' shows a large central elongated cell, surrounded by four or more smaller cells, which are also elongated in the longitudinal direction, and which are known as *siphons*. These *pericentral tubes* are often connected with one another and with the axial cell by threads of protoplasm. When the pericentral tubes are wanting the frond is *monosiphonous*. In some genera belonging to the Ceramiaceæ either the single axial tube or both axial and pericentral tubes are surrounded by a *pseudo-cortex* formed of a dense agglomeration of secondary branches originating at the nodes and closely adpressed to the primary branch. These are always the result of further division of the pericentral tubes, the apical cell remaining undivided. In other genera a similar cortical tissue is composed of moniliform rows of cells arranged at right angles to the axis.

FIG. 169.—*Gigartina mamillosa* Ag., with cystocarps (natural size). (After Luerssen.)

Although there is in the Florideæ no distinct differentiation of the

tissue into epidermal, assimilating, and conducting systems, still there are, in the higher forms, cells which are especially concerned with assimilation, and which may be either isodiametrical, or elongated in either direction. Such assimilating tracts are classed by Wille under three heads, viz.:—(1) those which act also for purposes of conducting; (2) those which are altogether distinct from the conducting cells; and (3) those where, in addition to an assimilating, there are also primary and secondary conducting cells. In some species the 'frond' assumes the appearance of a stalked leaf, as in Hydrolapathum (Rupr.) and Delesseria (Grev.), often of the most beautiful form, and presenting even a rudimentary venation. The genera Melobesia (Lmx.), Hildenbrandtia, Cruoria, and some others, consist of small algæ, mostly marine, with crustaceous or gelatinous thallus, growing flat on stones or larger algæ, often of lichen-like appearance. In their mode of growth some Florideæ display bilateral symmetry, and the branching may be either monopodial or sympodial. In Polysiphonia, Spyridia



FIG. 170.—*Polysiphonia opaca* Zan. *a*, with cystocarps; *b*, with tetrasporanges (natural size); *c*, branch with tetrasporanges; *d*, branch with cystocarp (× 100). (After Kützing.)

(Harv.), and some others, the 'phyllotaxis' is spiral. The genera Rytiphlœa (Ag.), Vidalia (Lmx.), Amansia (Lmx.), and Polyzonia (Suhr) are distinguished by the endogenous formation of the normal lateral shoots. In Pollexfenia (Harv.) and allied genera, Falkenberg records a remarkable congenital union of all the branches of the thallus which lie in the same plane.

The red colouring matter of the Florideæ has been investigated by Rosanoff, Cramer, Askenasy, Sorby, and others. Cramer first extracted it, from Bornetia secundiflora (Ag.) and Callithamnion caudatum (Ag.), in the form of regular crystals, partly hexagonal, partly octohedral, and gave it the name *rhodospermin*. Rosanoff extracted it by cold fresh water, and found it carmine-red in transmitted, reddish yellow, or rarely green, in reflected light. It is insoluble in alcohol. The chlorophyll-grains also exhibit fluorescence when left behind, if the pigment, the *phyco-erythrin* of Askenasy and Sorby, has escaped from them in consequence of injury to the plant. The spectrum of the chlorophyll is nearly identical with that in Flowering Plants. The compound pigment of the red Algæ is termed by Schütt *rhodophyll*, the term phyco-erythrin being limited to the portion soluble in water, while the portion soluble in alcohol he calls *Florideæ-green*. Cohn found, in Bornetia (Thur.), colourless crystalloids of an albuminous substance coloured a beautiful red by the same pigment; and Klein has found similar crystalloids in several Florideæ. The chromatophores contain starch-grains, which differ both from the ordinary grains of Flowering Plants and from those of the brown Algæ in being coloured brown or red by iodine. Schmitz (Sitzber. Niederrhein. Gesell., 1880) has detected a number of nuclei in the vegetative cells of many Florideæ, but not in the reproductive cells. Hick (Proc. Brit. Ass., 1883; 'Nature,' vol. xxix., 1884, p. 581), Massee (*l.c.*), and Moore (Journ. Linn. Soc. Bot., vol. xxi., 1886, p. 595) find continuity of protoplasm from cell to cell very beautifully displayed in Callithamnion (Lyng.), Ptilota (Ag.), Polysiphonia, and several other genera, not only in the vegetative cells, but also in the tetrasporanges. In Corallina (L.), Melobesia, Liagora



FIG. 171. — *Hydrolapathum sanguineum* Stackh. *a*, two young fronds with two cystocarps; *b*, *c*, prolifications from the stem (natural size). (After Kützing.)

(Lmx.), and a few other genera, the 'frond' becomes densely incrusted by a deposit of calcium carbonate, giving to the so-called 'corallines' the external form and appearance of miniature corals.

The ordinary non-sexual propagative organs of the Florideæ are bright red motionless spores, commonly formed in fours in the mother-cell, and hence known as *tetraspores* (the sphærospores of Agardh), and the cell in which they are produced as a *tetrasporange*. The four spores are sometimes arranged in a row, when they are called *zonate* ; more often as quadrants of a sphere, when they are *cruciate* : rarely there are only one or two, or occasionally eight. In the Ulvaceæ, Lemaneaceæ,



FIG. 172.—*Crouania attenuata* Ag. *a*, branch (× 40) ; *b*, apex of branch (× 100) ; *c*, lower portion of branch with tetrasporanges (× 100). (After Kützing.)

and in some Nemalieæ they are altogether wanting. The tetraspores (see fig. 231) may be formed in the six following ways :—(1) The whole contents of the sporange become a single spore ; (2) the contents divide into two equal parts by a transverse wall ; (3) they divide into four quadrants by two successive bipartitions ; (4) they divide into four tetrahedra by simultaneous quadripartition ; (5) they divide into four by three parallel transverse walls ; (6) the contents divide into more than four spores. On germination the tetraspores may give birth either to sexual or to non-sexual individuals. In the monosiphonous Florideæ the tetrasporanges are usually formed at the expense of the ultimate branchlets. In other forms they are most commonly found scattered

near the margin of the 'frond,' sometimes (Rhodymenia bifida, Ktz.) imbedded in the thallus, and then often grouped into *sori* (Nitophyllum, Grev.); or, in the Corallinaceæ, enclosed in special *conceptacles*. In other genera (Phyllophora, Grev., &c.) they are developed in *nematheces*, wart-like elevations of the surface, where they are accompanied by barren hyphæ or *paraphyses*. In others again they are borne on metamorphosed pod-like branches known as *stichids*, as in Dasya (Ag.), Plocamium (Lmx.), &c. Only in the Porphyraceæ are the tetraspores endowed with a slow



FIG. 173.—*Nitophyllum punctatum* Harv. *a*, piece of frond with tetrasporanges (natural size); *b*, piece of the thallus with tetrasporanges (× 100); *c*, section through frond showing cystocarp (× 100. (After Kützing.)

amœboid motion. Zoospores are altogether unknown in the class, except in the Ulvaceæ; but other modes of non-sexual propagation occur in a few cases. In some genera of Ceramiaceæ special organs occur, known as *seirospores*. Melobesia is characterised by the production of *gemmæ*. In Monospora (Sol.) stalked gemmæ or *propagules* are produced at the forks of the branchlets, and readily become detached, apparently replacing the sexual organs, which are unknown in the genus. Lemanea (Bory) increases by budding. In Hydrolapathum peculiar bud-like prolifications are produced on the stem (see fig. 171).

FIG. 174.—*Dasya elegans* Ag. *a*, piece of frond with tetrasporanges; *b*, piece with cystocarp (natural size); *c*, branch with stichids (× 100); *d*, cystocarp (× 25). (After Kützing.)

A true understanding of the sometimes complicated process of sexual reproduction in the Florideæ has been much obscured by the numerous terms employed by the older writers for identical organs, and by incorrect notions as to their functions. The true sexual organs, *antherids* and *procarps*, are nearly always formed on individuals which do not produce tetraspores ; and the sexual individuals may be monœcious or diœcious ; the latter is the most common condition. If the Ulvaceæ are rightly included under Florideæ, we have here a wide departure from



FIG. 175.—Stages in the development of the reproductive organs of *Nemalion multifidum* Ag. (magnified). (*Spermat.* = pollinoids.)

the normal type, sexual reproduction taking place by the conjugation of motile swarm-cells.

The *antherid* consists, in its simplest form (Porphyraceæ), of a solitary cell at the end of a long segmented branch ; and in this case it gives birth to a single pollinoid ; in other forms the antheridial cells occupy a similar position to the tetraspores, being formed in groups at the expense of the ultimate branchlets. They are also sometimes pro-

duced, like the tetraspores, in wart-like protuberances or *nemathèces* on the surface of the thallus, and interspersed with paraphyses, sometimes (Gracilaria, Grev.) in depressions which are overarched by the surrounding tissue ; or, in the Corallinaceæ, in special *conceptacles*. When the thallus is otherwise unilamellar, as in Porphyra (Ag.), the spots where the antherids are formed divide by walls parallel to the surface. The fertilising bodies or *pollinoids* are naked masses of protoplasm, of a spherical or elongated form, sometimes with a beak-like appendage, and are discharged in succession one after another. They are carried along passively by the water, and are distinguished from the antherozoids of other Cryptogams by the absence of cilia, and, in most cases, of any spontaneous power of motion. Wright (Trans. Irish Acad., 1879, p. 27)

FIG. 176.—*Spermothamnion hermaphroditum* (magnified). *A*, branch with procarp (*t f g i*) and antherid (*an*) before fertilisation ; *B*, after fertilisation, the cystocarp developing ; *t*, trichogyne ; *c*, trichophore ; *g*, carpogenous cells. (After Nägeli.)

states that in Griffithsia (Ag.) the pollinoids have an obscure amœboid motion, as they have also in the Porphyraceæ ; according to Dodel-Port their access to the trichogyne is greatly facilitated by the currents made in the water by Vorticellæ and other Infusoria ; and there can be little doubt that fishes which feed on seaweeds are an important agent in promoting their fertilisation. The pollinoids and the tetraspores appear to be homologous in their origin.

The female organ before fertilisation—corresponding functionally to the pistil of Flowering Plants—is termed the *procarp*. In its simplest form (Porphyraceæ and Nemalieæ) it consists of a single cell with a lateral hair-like prolongation, the *trichogyne*. But in all the higher forms the procarp is composed of one or more fertile cells constituting the *carpogone*, and one or more infertile cells which make up the *tricho-*

phore, the function of which is to convey the fertilising substance from the trichogyne to the carpogone. The procarp is usually formed on the youngest parts of the plant, and often originates from the terminal cell of a lateral branch. Occasionally each carpogone has two trichogynes and two trichophores; or, again, each trichophore may be connected with two carpogones. The trichogyne often becomes eventually coiled



FIG. 177.—*Lejolisia mediterranea* Born. *a*, filament with tetrasporange; *b*, plant with cystocarps and antherids; *c*, empty cystocarp (× 150). (After Bornet.)

spirally at its base. It does not open to admit the entrance of the pollinoids; in the act of impregnation these bodies attach themselves to a spot near the apex of the trichogyne, and at the same time clothe themselves with a cell-wall. At the point of contact the cell-wall of both trichogyne and pollinoid is absorbed; the contents of the latter pass through the trichogyne and the trichophore to the carpogone, impregnat-

ing its contents, and the trichogyne then disappears. The carpogone now divides by a horizontal wall into two cells ; the upper one of these is functionless, and ultimately disappears ; the lower one contains the impregnated oosphere or *carposperm*. The carposperm does not, however, in any case possess the power of germinating directly. In the Porphyraceæ it breaks up into eight portions, the *carpospores*, which germinate after moving about with a slow amœboid motion. In all the other orders the contents of the carpogone undergo, after impregnation, more complicated divisions, and become differentiated into a sterile and a fertile portion, the *placenta* and the *nucleus*. The placenta may consist of one or more cells, and frequently occupies the larger portion of the carpogone ; or it may be reduced to very small dimensions. The nucleus is the mass of carpospores, and may be made up of a number of *secondary nuclei*. The mass of carpospores is sometimes, as in Callithamnion, completely exposed except for a gelatinous membrane by which it is surrounded ; but much more often there is gradually formed round the nucleus after impregnation, not only a layer of mucilage, but also a more or less hard solid layer, the *pericarp* ; and the whole structure thus constituted is then known as the sporocarp or *cystocarp*.

FIG. 172. – *Gracilaria compressa* Ag. Branch with cystocarps (natural size). (After Hauck.)

From this cystocarp the carpospores escape, when ripe, either by the decay of the pericarp ('coccidium' of the older systematists), or through an opening at its apex, the *carpostome* ('ceramidium' of older writers). In some genera (Polysiphonia, Lejolisia, Born., Bonnemaisonia, Ag.) the cystocarp is completely exposed, conspicuous, and sometimes stalked ; but it is usually, as in Gracilaria,

more or less imbedded in the thallus, often in special fertile branches ; and its situation is then usually indicated by an external wart-like swelling. In Polyides (Ag.), the Squamariaceæ, and other forms with a perfectly flat frond, the cystocarps are enclosed in *nematheces*.

In a considerable number of Florideæ the formation of the cystocarp is a more complicated process than that already described, the process of impregnation consisting of two distinct stages—(1) the fertilisation of the trichogyne by the pollinoids ; and (2) the fertilisation by the impregnated trichophore-cells of the carpospores, which may be at some considerable distance from the trichophore and trichogyne, even on a



Fig. 179.—*Dudresnaya coccinea* Crouan. *I*, young trichophore. *II*, young trichogyne ; *f*, young fertilising tubes. *III*, impregnated trichophore with pollinoids on the coiled trichogyne, *f* ; the fertilising tube, *f'*, impregnating successively the carpogones, *VII*, *VI*, and *V*. *IV*, carpogone before fertilisation ; *c*, carpogenous cell. *VIII*, masses of carpospores.

different branch. This is effected by means of long simple or branched tubes, the *fertilising-tubes*, or 'ooblastema-filaments' of Schmitz. The following is the process as it takes place in Dudresnaya (Born.). The trichophore consists of a row of cells which, before fertilisation, put out short branches, which subsequently develop into long tubes. No carpogone is found in the immediate neighbourhood of the trichophore ; but at some distance are a number of short segmented filaments, the terminal cells of which are considerably larger than all the rest. These cells are carpogones. The fertilising-tubes make their way between the filaments or hyphæ of which the thallus is composed, come into contact

with the carpogones, and convey to them the fertilising principle from the trichophore. The result is that each carpogone develops into a cystocarp containing carpospores. A single fertilising-tube may in this way impregnate a number of carpogones.

Schmitz describes the process of secondary impregnation in the more highly developed Florideæ as consisting in the fertile cells (carpogones) entering into communication, through orifices in their cell-walls, with certain special sterile cells rich in protoplasm, the *auxiliary cells*, to which the fertilising material is brought from the trichophore by the ooblastema-filaments. The details of this conjugation between the auxiliary cells and the carpogones are subject to great variation in different genera. In some cases the protoplasmic contents of the two cells coalesce completely, while in others their nuclei still remain distinct after conjugation. The carposperm, or cell resulting from this conjugation, then grows rapidly, and peripheral cells divide off from it, leaving a large central cell which alone remains sterile, all the peripheral cells developing into carpospores. In the Corallinaceæ the ooblastema-filament enters into conjugation successively with several neighbouring auxiliary cells. In a larger number of genera the process is as follows :— A short branch of the carpogone, usually consisting of three or four cells, becomes attached laterally to a branch of the thallus, and curves in such a way that the carpogone-cell is closely applied to the nearest auxiliary cell, or reaches it by means of a short protuberance from one or both of the conjugating cells. The entire oosphere, or at all events its nucleus, then passes over into the auxiliary cell. In the Gigartinaceæ the auxiliary cell itself becomes the central cell of the cystocarp.

Literature.

Nägeli u. Cramer—Pflanzenphysiol. Untersuch., 1855, 1857.
Pringsheim—Monber. Berl. Akad. Wiss., 1855, p. 133 (Quart. Journ. Microsc. Sc., 1856, p. 124).
Rosanoff—(Rhodospermin) Mém. Soc. Sc. Nat. Cherbourg, 1856; Ann. Sc. Nat., iv., 1865, p. 320; and Compt. Rend., lxii., 1866, p. 831.
Van Tieghem—Compt. Rend., lxi., 1865, p. 804.
Solms-Laubach—Bot. Zeit., 1867, p. 161.
Askenasy—(Rhodospermin) *Ibid.*, p. 233.
Sorby—(Rhodospermin) Journ. Microsc. Soc., 1871, p. 124.
Klein—Flora, 1871, p. 161; and 1880, p. 65.
Agardh—Epicrisis Syst. Florid., 1876; and Florid. Morphol. (with atlas), 1879.
Bornet and Thuret—Notes Algologiques, fasc. i. and ii., 1876, 1880; and Etudes Phycol., 1878.
Falkenberg—Nachricht. Gesell. Wiss. Göttingen, 1879 and 1880.
Ambronn—Bot. Zeit., 1880, p. 61; and Sitzber. Bot. Verein Brandenburg, 1880, p. 74.

Schwendener—Monber. Preuss. Akad. Wiss., 1880, p. 327.
Berthold—Pringsheim's Jahrb. wiss. Bot., 1882, p. 569.
Schmitz—Sitzber. Akad. Wiss. Berlin, 1883, p. 215.
Ardissone—Phycol. mediterranea, Part i., Florideæ, 1883.
Buffham—Journ. Quek. Micr. Club, 1884, p. 337.
Massee—Journ. Microsc. Soc., 1884, pp. 198 *et seq.*; and 1886, p. 561.
Wille—(Tissue-systems) Bot. Sälsk. Stockholm (see Bot. Centralblatt, xxi., 1885, pp. 282, 315; xxiii., 1885, p. 330; and xxvi., 1886, p. 86); and Nov. Act. Leopold-Carol. Akad., lii., 1888, p. 51.
Schütt—(Phyco-erythrin) Ber. Deutsch. Bot. Gesell., 1888, p. 36.

A complete classification of the very numerous types of structure belonging to the Florideæ would carry us beyond our present limits; and the principles of such a classification are by no means agreed on by the best authorities, many details connected with the process of fertilisation being still obscure. Agardh, in his 'Epicrisis,' divides the family into six groups, dependent on the structure and mode of development of the cystocarp, and into twenty-two orders. We shall therefore confine ourselves to an account of some groups or special forms, the structure of which has been specially studied, commencing with the most highly differentiated orders.

In the Ceramiaceæ, which are exclusively marine, are included a considerable number of the more delicate red seaweeds of our own and other coasts, included in the genera Callithamnion (Lyng.), Griffithsia (Ag.), Ptilota (Ag.), Crouania (Ag.), Ceramium (Lyng.), and others. The thallus is either monosiphonous and uncorticated, or more or less corticated. In Crouania (fig. 172) the branches are beautifully whorled. The procarp frequently consists of a carpogone and two trichogynes. The cystocarps are formed externally on the branches or at their base, and are frequently closely surrounded by them as by an envelope. With rare exceptions the cystocarp consists of a roundish or lobed nucleus, enclosed in a colourless gelatinous membrane, without any pericarp, and composed of a larger or smaller number of closely packed carpospores. The tetrasporanges are usually external, and the mode of division of their contents varies greatly. The complicated process of fertilisation in Dudresnaya has already been described. In addition to the tetraspores, two other special kinds of non-sexual organs of propagation occur in the order. Some species of Ceramium are characterised by the presence of *favellæ*, dense agglomerations of spores, resembling the cystocarps, but produced at the ends of branches, quite exposed except for a thin colourless membrane. They appear to be homologous to the multipartite tetrasporanges. Callithamnion seirospermum (Griff.) (Seirospora Griffithsiana, Harv.), C. versicolor (Drap.), and some other species, produce *seirospores*,

FIG. 180.—Branch of *Callithamnion seirospermum* Griff., with seirospores (× 200). (After Kützing.)

FIG. 181.—Branch of *Ceramium strictum* Grev., with favellæ (× 40). (After Kützing.)



FIG. 182.—*Melobesia membranacea* Lmx. *a*, vertical section through female conceptacle; *b*, vertical section through conceptacle with tetrasporanges; *c*, vertical section through male conceptacle (× 350); *d*, pollinoids (× 1300). (After Rosanoff.)

branched rows of roundish or oval spores resulting from the division of terminal cells of particular branches, or produced on the main branches.

LITERATURE.

Cramer—Pflanzenphysiol. Untersuch., 1857 and 1863.
Nägeli—Sitzber. Münch. Akad. Wiss., 1861, p. 297.
Pringsheim—Abhandl. Berlin. Akad. Wiss., 1862, p. 1.

The CORALLINACEÆ (Corallina, L., Melobesia, Lmx., Lithothamnion, Phil., Amphiroa, Lmx., &c.) are distinguished from other marine



FIG. 183.—*Corallina officinalis* L. *a*, longitudinal section through conceptacle with tetrasporanges; *b*, longitudinal section through cystocarp (× 100). (After Bornet.)

algæ by their calcareous habit. Most of the species are natives of warmer seas. In Corallina the thallus is at first soft and flexible, but it soon becomes very hard and brittle from the deposition of calcium carbonate. The red colour and branching habit give this genus a remarkable resemblance to small corals, as in C. officinalis (L.), the common 'coralline' or 'nullipore' of our southern coasts. Many species of Melobesia (fig. 182), Lithothamnion, Lithophyllum (Phil.), and other genera, grow as lichen-like incrustations or in the form of small flat

discs attached to rocks, or on the leaves of Zostera, or on other seaweeds. The sexual reproductive organs and the non-sexual organs of propagation are alike formed in small cavities or *conceptacles*, which are either entirely imbedded in the thallus, or more often form external wart-like or ovoid swellings. The female conceptacle opens at the apex by an *ostiole* (fig. 182, *a*); the very short sporiferous filaments, the terminal cells of which



FIG. 184.—*Corallina rubens* L. *a*, branch with three cystocarps and a male conceptacle; conceptacles of *Melobesia Thuretii* Born. are attached to the upper part of the branch (× 20); *b*, longitudinal section through a male conceptacle deprived of its calcareous incrustation (× 160); *c*, pollinoids (× 400). (After Bornet.)

become the carpospores, spring from the base of the cavity, and are accompanied by paraphyses. The male conceptacles are of similar structure; the pollinoids (fig. 182, *d*) possess one or two short appendages. The non-sexual present a general resemblance to the sexual conceptacles; the tetrasporanges spring from their base and sides, and are accompanied by paraphyses; the contents of the sporange not un-

frequently divide into only two tetraspores. The conceptacles are not unfrequently surmounted by singular horn-like processes. Melobesia produces peculiar branching septated *gemmæ*.

LITERATURE.

Rosanoff—Mém. Soc. Sc. Nat. Cherbourg, 1866.
Areschoug—Observ. Phycol., iv., 1875.
Solms-Laubach—Die Corallinenalgen d. Golfes Neapel, 1881.

The greater number of our red seaweeds belong to the following orders, established by Agardh, the boundaries of which are not in all

FIG. 185.—*Chondrus crispus* Stackh. *a*, with cystocarps (natural size); *b*, uppermost portion of frond with tetrasporanges (natural size); *c*, section through the frond and a portion of the cystocarp (× 100). (After Kützing.)

cases well defined, and in regard to many of which much yet remains to be discovered as to their mode of reproduction, viz.:—CRYPTONEMIACEÆ (Nemastoma, Ag., Grateloupia, Ag., Halymenia, Ag., Dumontia, Lmx., Cryptonemia, Ag., &c.), GIGARTINACEÆ (Chondrus, Grev., Gigartina, Lmx., Kallymenia, Ag., &c.), RHODYMENIACEÆ (Rhodymenia, Grev., Chylocladia, Grev., Plocamium, Lmx., Hydrolapathum, Rupr., &c.), DELESSERIACEÆ (Nitophyllum, Grev., Delesseria, Lmx.), SPHÆROCOCCACEÆ (Sphærococcus, Stackh., Gracilaria, Grev., &c.), HYPNÆACEÆ

(Hypnæa, Lmx.), GELIDIACEÆ (Gelidium, Lmx., &c.), SPONGIOCARPEÆ (Polyides, Ag.), LOMENTARIACEÆ (Lomentaria, Gaill.), RHODOMELACEÆ (Laurencia, Lmx., Chondria, Ag., Rhodomela, Ag., Polysiphonia, Ag., Rytiphlæa, Ag., Amansia, Lmx., Vidalia, Ag., Dasya, Ag., Pollexfenia, Harv., &c.), SPYRIDIACEÆ (Spyridia, Harv.), and WRANGELIACEÆ (Spermothamnion, Aresch., Bornetia, Thur., Lejolisia, Born., Wrangelia, Ag., &c.). A peculiar mode of fertilisation is described in the case of Gracilaria confervoides (Grev.). The procarp is composed of six or seven cells, and its position is indicated by an external swelling on the 'frond;' the pericarp is formed before fertilisation. The cells both of the procarp and of the placenta fuse together by the disappearance of their cell-walls; and the fused cells of the procarp and of the placenta are placed in communication with one another by protoplasmic protrusions or *diverticula*, proceeding from the fused cells of the procarp and passing through their swollen walls. The cells which form the free surface of the placenta now produce radiating rows of spores formed basipetally, while from the fused procarpial cells other diverticula arise which also form spores at their free ends independently of the placental cells. It would appear as if by this means a portion of the protoplasm of the pollinoid were conveyed through the trichogyne and the procarpial cells, which play the part of the auxiliary cells of Dudresnaya, to the placental cells. In Polyides the cystocarps are congregated in external amorphous wart-like excrescences formed of vertical confervoid filaments, of which the cartilaginous 'frond' is almost entirely composed, set in firm jelly.



FIG. 186.—*Rhodymenia Palmetta* Grev. *a*, with tetrasporanges (natural size); *b*, section through apex of frond with tetrasporanges (× 100). (After Kützing.)

Plocamium coccineum (Huds.) is one of the most abundant of our red seaweeds. Rhodymenia palmata (Grev.), a very common red seaweed,

is eaten in Scotland under the name of 'dulse.' Chondrus crispus (L.) is known, when dried, as 'carrageen' or 'Irish moss.' Gracilaria lichenoides (L.), which yields copious gelatine, known as 'Fucus amylaceus,' is largely used as an article of food in Ceylon and on other tropical coasts. Japanese isinglass is a product of species of Gelidium. The



FIG. 187.—Vertical median section of swelling on frond of *Gracilaria confervoides* Grev. *p*, *p*, cells of procarp ; *pc*, placental cells ; *s*, *s*, *s*, spores (× 400). (After Johnson.)

substance known to microscopists as 'agar-agar' is yielded also by Gracilaria lichenoides, Eucheuma spinosa (Ag.), and other seaweeds. Several other species are employed in different parts of the world as glues and varnishes. Marchesettia spongioides (Hauck) (Areschougiaceæ), from Singapore, has a remarkable resemblance to a sponge.

LITERATURE.

Berthold—(Cryptonemiaceæ) Fauna u. Flora Golfes Neapel, 1884.
Johnson—(Gracilaria) Annals of Botany, i. 1888, p. 213.

The SQUAMARIACEÆ (Hildenbrandtia, Nard., Cruoria, Fries, Peyssonnelia, Dcne., &c.) are a small group of small marine, or rarely fresh-water Algæ, growing on stones or on the shells of molluscs or crustacea, or attached to larger algæ. The 'frond' is expanded flat or hemispherical, gelatinous, membranaceous, or crustaceous, with lichen-like habit, composed of a single layer of cells, or more often of short densely

packed vertical filaments. The species of Hildenbrandtia form rose-coloured incrustations on rocks, stones, and shells in salt or (H. rivularis, Ag.) running fresh water. The tetrasporanges are either terminal cells of special filaments, enclosed in a gelatinous coating and rising vertically from the flat thallus ; or they are formed in nematheces, in external wart-like protuberances, or in depressions in the surface of the thallus. The cystocarps are also either formed in nematheces, or are external,



Fig. 188.—*Hildenbrandtia prototypus* Nardo. Vertical section through thallus, showing three conceptacles with cystocarps (× 300). (After Kützing.)

springing from moniliform fertile filaments. It is stated that there are sometimes two kinds of carpogone, one provided with a trichogyne, the other not. After a carpogone of the first kind has been fertilised through its trichogyne, it puts out a fertilising-tube or 'ooblastema-filament,' which impregnates a carpogone of the second kind. In Hildenbrandtia a large number of antherids are developed from a single cell of the thallus.

Literature.

Schmitz—Sitzber. Niederrhein. Gesell., 1879.
Borzì—Rivista Scientifica, 1880.
Petit—Bull. Soc. Bot. France, 1880, p. 194.
Wollny—Hedwigia, 1886, pp. 1 and 125.

The Helminthocladiaceæ (including Nemalieæ, Batrachospermeæ, and Chætangiaceæ) comprise a number of marine (Helminthocladia, Ag., Nemalion, Ag., Liagora, Lmx., Galaxaura, Lmx., &c.) and fresh-water (Chantransia, Fries, Batrachospermum, Roth, Thorea, Bory) forms, the relationship of which to one another is uncertain, and the family is not likely to be one that will be ultimately retained. The fresh-water species are mostly of small size, but of great beauty from the elegant symmetry and arrangement of the branches. The 'frond' of Liagora and Galaxaura is calcareously incrusted, like that of a coralline ; that of

some other genera is very soft and gelatinous: the whole is usually enclosed in a gelatinous envelope. The thallus is of filamentous structure, and either simple or branched, the secondary axes often arranged in whorls. The main axis usually consists of a central row of cylindrical cells surrounded by a *pseudo-cortex* composed of smaller cells, either placed in rows at right angles to the axis, or parallel to it. It may, however, consist of a single uncorticated row of cells. Batrachospermum is a genus of small green fresh-water Algæ of great beauty from the symmetry of their branching. The stem grows by means of a cup-shaped



FIG. 189.—*Batrachospermum moniliforme* Roth. *a*, portion of plant (× 30); *b*, portion with cystocarps, *cccc* (× 150). (From nature and after Cooke.)

apical cell, which divides by septa ; the resulting cells do not divide further, but elongate and swell somewhat at each end into a bone-shaped form, producing the whorled branches. From the basal cells of these branches secondary branches grow vertically downwards over the main axis, producing the pseudo-cortex. According to Sirodot, absorption takes place only in special thin-walled cells ; the *resting-cells*, with thicker walls, often display continuity of protoplasm. In Nemalion and Batrachospermum the procarp is unicellular, and bears at its apex the long thin-walled trichogyne. The antherids are scattered in groups at the end of peculiar ovoid cells. After fertilisation the carpogone divides by longitudinal walls into a multicellular *glomerule* enclosed in a gelatinous envelope ; the terminal cells of the branches of the glomerule produce the spores. The *cystocarps* are external in the axils of the branches in Chantransia and

FIG. 190.—*Chantransia virgatula* Thur. with tetrasporanges (× 100). (After Kützing.)

FIG. 191.—*C. corymbosa* Thur. *I*, unicellular procarp, *b*, with pollinoid attached. *II–V*, stages in the branching of the procarp after impregnation (× 400). *VI*, cystocarp (× 250). (After Thuret.)

Batrachospermum, more or less imbedded in the thallus in the other genera of the order. In several of the genera tetraspores are unknown.

A very singular genetic connection exists between the genera Chantransia and Batrachospermum, it being possible to transform the former into the latter by changing its conditions of life. The germinating carpospores of Batrachospermum put out a kind of protoneme, which is the Chantransia of Fries, the non-sexual generation of Batrachospermum; this can propagate itself by simple budding from generation to generation, producing, as a rule, as its organs of propagation, only non-sexual tetraspores. Chantransia grows especially in dark situations under water, and, when transferred to the light, undergoes a metamorphosis. There springs up from the Chantransia protoneme a branch which is in every respect a Batrachospermum, and which bears sexual organs only, and no tetraspores. On one species only of Chantransia, C. corymbifera (Thur.), are sexual organs known. Although this phenomenon is sometimes spoken of as an example of 'alternation of generations,' it is not identical with the process known under that name in the higher Cryptogams, being rather a difference in the mode of development dependent on a change in the vital conditions.

LITERATURE.

Solms-Laubach—Bot. Zeit., 1867, p. 161.
Sirodot—Compt. Rend., lxxvi., 1873, pp. 1216, 1335; lxxix., 1874, p. 1366; xci., 1880, p. 862; xcii., 1881, p. 993; Bull. Soc. Bot. France, 1875, p. 128; and Les Batrachospermées, 1884.
Arcangeli—Nuov. Giorn. Bot. Ital., 1882, p. 155.
Massee—Journ. R. Microsc. Soc., 1886, p. 561.

The LEMANEACEÆ are a small group of fresh-water Algæ, comprising the genera Lemanea (Bory) and Sacheria (Sir.), growing in rapidly running water, as beneath mill-wheels. The thallus is filiform and cartilaginous, of a dull grey or greenish colour, and consists of a single row of tubular cells, or of an axial row surrounded by rows of smaller cortical cells; it displays swellings or projections at regular intervals. It grows by means of an apical cell, from which segments are cut off at right angles to the direction of growth. By transverse septation each segment divides into a central cell surrounded by peripheral cells; the central cell becomes a member of the central axis, the peripheral cells members of the cortical tubes. The 'frond' increases by budding at the free surface of the rooting system, finally producing cæspitose tufts. No non-sexual spores are known. The only other mode of reproduction is sexual. The antherids are short cylindrical cells produced externally in the neigh-

bourhood of the swellings, with a more or less regular annular arrangement. The trichogyne is a long transparent cylindrical tube, simple or branched, produced within the tubular thallus. The procarp is unicellular, and of a very simple structure. The carpogone puts out before impregnation a number of segmented filaments resembling paraphyses, and enclosed in a transparent jelly. After the oosphere has been im-



FIG. 192.—*A*, *Lemanea nodosa* Ktz. (natural size). *B*, section of tube with female organs; *t*, trichogyne, at the base of which sporiferous filaments (*s*) are beginning to be formed (× 200). *C*, transverse section of male organs; *a*, antherids (× 400). (After Sirodot.)

pregnated, it puts out an 'ooblastema-filament,' and the trichogyne disappears; at the extremity of this filament are produced the carpospores. On germinating the carpospore puts out a protonemal filament somewhat resembling that of mosses, on which the fertile axes are produced as lateral branches. This is regarded by some as a rudi-

mentary alternation of generations ; and Peter asserts (Bot. Verein München, Feb. 28, 1887) that the sexual form of Lemanea fluviatilis (Ag.) may develop out of heteromorphic branches of a Chantransia.

LITERATURE.

Wartmann—Beitr. zur Anat. und Entwick. der Lemanea, 1854.
Sirodot—Ann. Sc. Nat., xvi., 1872, p. 5; 1873, p. 241; and Compt. Rend., lxxix., 1874, p. 1366.
Ketel—Anat. Untersuch. über Lemanea, 1887.

The PORPHYRACEÆ or Bangiaceæ are marine or fresh-water Algæ belonging to the two genera Porphyra (Ag.) and Bangia (Lyngb.). They are regarded by some authors as much lower in structure than the red

FIG. 193.—*a*, *b*, *Bangia ceramicola* Chauv. Portions of barren and fertile filaments (× 330). (After Thuret.)



FIG. 194.—*B. fusco-purpurea* Lyngb. Filament in different stages of development (× 200). (After Kützing.)

seaweeds, but are best placed at present, from their mode of sexual reproduction, as one of the lowest families of Florideæ. The thallus is either filiform (Bangia), or is a thin transparent plate (Porphyra) com-

posed, in its vegetative portion, of a single layer of cells ; in both cases coloured by phycoerythrin. The tetrasporanges and the male and female organs appear to be homologous to one another, and not to be sharply differentiated. The tetraspores are motile for about forty-eight hours after their escape from the tetrasporange ; by some writers they are described as being endowed with an amœboid change of form. The trichogyne is quite rudimentary ; the pollinoids attach themselves singly or in numbers to the fertile portion of the thallus where the oogones or rudimentary carpogones occur. While in this position they are invested by a thin cell-wall of cellulose, and then put out a slender thread of protoplasm which pierces the cell-wall of the oogone, nearly the whole of the protoplasm of the pollinoid passing into this organ. According to Berthold, the contents of the oogone break up, after impregnation, into eight carpospores, the 'octospores' of Janczewski, which move about, on escaping, in an amœboid manner, putting out and withdrawing protoplasmic protrusions, then come to rest and germinate. Porphyra vulgaris (L.), not uncommon on the coasts of Western Europe, is eaten under the name 'purple laver.'



FIG. 195. Tetraspores of *Bangia fusco-purpurea* Lyng., showing amœboid changes of form (magnified). (After Reinke.)

LITERATURE.

Janczewski—Ann. Sc. Nat., 1873, p. 241.
Reinke—Pringsheim's Jahrb. wiss. Bot., 1878, p. 274.
Goebel—Bot. Zeit., 1878, p. 199.
Berthold—Mittheil. Zool. Stat. Neapel, 1880 and 1882.

The position of the ULVACEÆ is still uncertain. The group includes a small number of genera—Ulva (L.), Enteromorpha (Lk.), Phycoseris (Ktz.), Prasiola (Ag.), and Monostroma (Thur.)—of fresh-water or more often of marine or brackish Algæ, of a bright green colour, consisting of a flat usually ribbon-shaped plate, composed of either one (Monostroma) or two (Ulva) layers of cells ; less often (Enteromorpha) having the form of a tube. The cells are sometimes arranged symmetrically in groups of four (Prasiola). The male and female reproductive organs, which are rudimentary in the Porphyraceæ, are entirely suppressed in the Ulvaceæ, and we find a reversion to a much simpler mode of repro-

duction in the conjugation of equivalent swarm-cells. Ulva produces two kinds of swarm-spore—*megazoospores* with four cilia, and micro-



Fig. 196.—*Ulva*. *a*, portion of frond showing cells which produce the swarm-spores ; *b*, portion of frond with empty cells ; *c*, megazoospores (magnified).

Fig. 197.—*Enteromorpha intestinalis* Lk. (natural size). (After Hauck.)

zoospores or *zoogametes* with two cilia ; and these two kinds are produced either on the same or on different individuals. The megazoospores germinate directly. Conjugation of the smaller swarm-cells has been observed in Ulva, Enteromorpha, and Monostroma, but the coalescence takes place only gradually ; some time after conjugation four cilia and two pigment-spots are still to be detected. In Ulva the *zygosperm* thus formed divides repeatedly after becoming attached to some solid substance ; the cells thus formed separate, and a small colony of multicellular individuals is produced, each of which develops into a new 'frond,' dividing first into a filament and then into a plate. In Monostroma a non-sexual mode of propagation has been observed. At



Fig. 198.—Stages in the conjugation of the zoogametes of *Monostroma bullosum* Thur. (magnified).

certain spots the cells divide in a direction parallel to the plane of the thallus, and a small projection is thus formed which becomes detached and develops into a new 'frond.' Geddes (Trans. R. Soc. Edinburgh, 1881, p. 555) describes also a process of gemmation in Enteromorpha. The ordinary mode of reproduction in the Ulvaceæ so closely resembles that in the Confervaceæ that the two families are generally considered as nearly allied: but the difference in the structure of the thallus is so great that it is difficult to believe in any near affinity between them. On the other hand, the structure of the thallus in Ulva and Porphyra is almost identical; and it is at least as probable that the Ulvaceæ are derived from the Porphyraceæ by further retrogressive metamorphosis displayed in the complete suppression of the antherids and carpogones, and the reversion of the tetraspores into motile swarm-spores of two kinds, the smaller of which are zoogametes. On the other hand, Monostroma may be allied to Tetraspora among the Protococcaceæ. Several species of Enteromorpha form a large portion of the green vegetation in salt ditches or on muddy sea-shores, or on rocks between high and low water mark; Prasiola grows on bare rocks or stones, or on salt soil; the substance known as 'green laver' consists of several species of Ulva.

LITERATURE.

Jessen—Prasiolæ Monographia, 1848.
Thuret—Mém. Acad. Sc. Nat. Cherbourg, 1854, p. 9.
Wittrock—Monogr. Monostroma, 1866.
Janczewski et Rostafinski—Mém. Acad. Sc. Nat. Cherbourg, 1874, p. 369.
Areschoug—Bot. Notiser, 1876.
Thuret et Bornet—Etud. Phycol., 1878.
Reinke—Pringsheim's Jahrb. wiss. Bot., 1878, p. 531.
Borzì—Studi Algologici, 1883.

Class XI.—Confervoideæ Heterogamæ.

The small group of green fresh-water Algæ comprised in this class form a connecting link between the lower Florideæ and the Isogamous Confervoideæ, from which latter they are distinguished by the sharp differentiation of their male and female organs of reproduction. In the highest order, the COLEOCHÆTACEÆ, the mode of reproduction bears a striking analogy to that in the Nemalieæ, being effected through the agency of a tubular trichogyne; but the male elements are no longer immotile pollinoids, but motile swarming antherozoids, naked biciliated or multiciliated protoplasmic bodies, but much more closely resembling the corresponding structures in the lower Algæ than the coiled anthero-

zoids of the Muscineæ and Vascular Cryptogams. The vegetative thallus is here either composed of branching filaments, or is reduced to a flat plate of cells. In the two lower orders, the Œdogoniaceæ and the Sphæropleaceæ, it consists of an unbranched filament of uninucleated or multinucleated cells; and in them the impregnation of the oosphere by the motile antherozoids is brought about directly, without the intervention of a trichogyne. Non-sexual propagation by means of swarming biciliated zoospores, formed within zoosporanges, and bearing a close resemblance to the antherozoids, occurs in the Coleochætaceæ and Sphæropleacæ; in the Œdogoniaceæ the zoospores are much larger bodies, bearing a tuft of cilia. The class displays a rudimentary alternation of generations.

Order 1.—Coleochætaceæ.

The typical genus of this small order, *Coleochæte* (Bréb.), comprises several species of small fresh-water Algæ forming minute discs or cushions attached to submerged plants, from $\frac{1}{10}$ to $\frac{1}{4}$ inch in diameter, consisting, in the simpler forms, of a single layer of cells, often arranged in rays proceeding from a common centre. Some of these cells are furnished with colourless bristle-like protuberances fixed into narrow sheaths. All the cells of which the thallus is composed usually lie in one plane, but their degree of union with one another varies. In C. scutata (Bréb.) they are closely united into a compact disc, which continues to grow by peripheral increase, the marginal cells dividing by radiating and tangential walls. In C. soluta (Pringsh.) the thallus consists of a number of branches, which ramify dichotomously, and lie side by side, more or less closely crowded, in one plane. In other species, as C. pulvinata (A. Br.) and divergens (Pringsh.), the branches do not ramify in one plane only, but develop also segmented ascending branches, which form, in the latter species, together with the original disc, a nearly hemispherical cushion. The entire thallus is always enveloped in mucilage.

Non-sexual propagation takes place in Coleochæte by means of biciliated *zoospores*, produced either in all the cells of the thallus or only in the terminal cells of the branches. The entire protoplasmic contents of the mother-cell or *zoosporange* are used up in the production of the zoospores, which escape from them either at the side or at the back. Sexual reproduction is effected by the impregnation of an *oosphere* formed within an *oogone* by motile *antherozoids*, through the agency of a *trichogyne*. The oogone is, in C. pulvinata, always constituted out of the terminal cell of a branch, which swells up and at the same time elongates at its extremity into a narrow hair-like trichogyne, which then

opens at its apex and exudes a drop of colourless mucilage. The oogone still contains chlorophyll, and its protoplasmic contents contract into a green oosphere. At the same time flask-shaped protuberances grow out from adjoining cells, and these, becoming cut off from the parent-cells by septa, are the *antherids*. The entire contents of each antherid escape as a single ovoid antherozoid furnished with two very long and slender cilia. Other species are diœcious, the antherids being produced on different individuals from the oogones. The antherozoids probably pass into the trichogyne through its open apex, and thence into the oogone; but the act of impregnation has not actually been observed hitherto.

The first result of the impregnation of the oosphere is its investment

FIG. 193. *Coleochæte pulvinata* A. Br. *A*, portion of fertile thallus (× 350): *an*, antherid; *og*, oogone; *h*, hyaline hair; *z*, antherozoid. *B*, ripe oogone with its pericarp, *r* (× 280). *C*, formation of carpospores within the spermocarp (× 280). *D*, zoospores (× 280). (After Pringsheim.)

by a cell-wall of cellulose, and a considerable increase in its size. The fertilised oogone, with the exception of the trichogyne, then becomes surrounded by a *pericarp*, or cortical layer of cells; the oogone and pericarp together constitute the *spermocarp* enclosing the fertilised oosphere or *oosperm*. The spermocarp subsequently becomes further invested by a *cortex* of closely applied branches resulting from the continued development of cells at the base of the spermocarp. After the complete development of this organ, which takes place between May and July, the vegetative cells of the thallus disappear, and its walls assume a dark-brown colour. The spermocarp remains dormant through the winter:

the cortical layer is then thrown off, and the oosperm divides into several cells or *carpospores*. The germinating carpospore does not give rise to a new thallus, but to a zoospore, which gives birth to several successive non-sexual generations propagated by zoospores, until the cycle of generations is completed by the production of a sexual individual.

Mycoidea parasitica Cunn. (Trans. Linn. Soc., vol. i., 1879, p. 301) is probably nearly allied to Coleochæte, which it resembles in the nature of its thallus and in its mode of reproduction. It is endophytic in the cells of the leaves of Camellia in tropical India, inflicting great injury on the trees. Ward (Trans. Linn. Soc., ii., 1884, p. 87) contends that Mycoidea is in reality an epiphyllous lichen. Möbius (Ber. Deutsch. Bot. Gesell., 1888, p. 242) regards Mycoidea, and the nearly allied Chætopeltis (Berth.), as more probably belonging to Chætophoraceæ.

Literature.

Brébisson—Ann. Sc. Nat., i., 1844, p. 25.
Pringsheim—Jahrb. wiss. Bot., 1860, p. 1.
Kny—Ber. Deutsch. Bot. Gesell., 1884, p. 93.

Order 2.—Œdogoniaceæ.

This small order, as at present constituted, comprises only two genera—Œdogonium (Lk.) and Bulbochæte (Ag.).

Œdogonium includes several species, abundant in streams, ponds, and tanks. They are readily distinguished by the fact that they never branch, by the cells being of small diameter and considerable length, filled with a homogeneous dark-green protoplasm with a parietal nucleus, and by the peculiar appearance of annular striæ near one end of some of the cells. These striæ result from the appearance known as 'intercalary surface-growth.' Below the septum is formed an annular deposit or cushion of cellulose; at this place the cell-wall splits, as if by a circular cut, into two pieces, which separate from one another, but remain united by a zone of the cell-wall formed by an extension of the cushion. This process is constantly repeated over a short space of the cell-wall immediately beneath a septum, each slit being a little further from the septum than those that preceded it; so that these pieces, forming small projections, give to the upper end of the cell the appearance of consisting of *caps* placed one over the other; while its lower end appears as if enclosed in a long sheath consisting of the portion of the cell-wall below the caps. This lower portion of the cell is always cut off by a septum from the upper cap-bearing portion. The filaments are fixed at their base by a *rhizoid* to solid bodies or submerged plants.

Non-sexual propagation takes place in Œdogonium by means of *zoospores*, the formation of which affords a typical example of the process

first described by A. Braun as the 'rejuvenescence' of a cell, i.e. the transformation of the entire protoplasm of a vegetative cell into a 'primordial cell,' which subsequently invests itself with a new cell-wall, and forms the starting-point of the life of a new individual. In some one cell of a filament, either the terminal or some other, sometimes even in the single cell of which a young filament is composed, the protoplasm contracts into a globular body which ultimately becomes free by the rupture of the cell-wall, by a transverse slit, into two very unequal halves. When this takes place in the terminal cell of a filament or the single cell of a young individual, the upper smaller portion of the cell-wall is lifted up like a lid, or even completely thrown off like a cap. The zoospore thus formed, which in some species is one of the largest and most striking known, has a nucleus, a red 'pigment-spot,' and an anterior hyaline region, to which is attached a tuft of cilia, visible even before its escape from its mother-cell. At the period of escape it is still enveloped in a transparent membrane, which, however, it soon breaks through, and then moves about in the water with great velocity for perhaps half an hour, displaying at this period a number of vacuoles. On coming to rest, the zoospore becomes attached by its anterior hyaline end, loses its cilia, invests itself with a cell-wall, puts out a rhizoid from the point of attachment, and develops into a filament with transverse septa. From the position occupied by the zoospore in the mother-cell, the direction of growth of the new individual must be at right angles to that of the parent-filament. Many of the plants which spring from zoospores are non-sexual, producing nothing but zoospores.



FIG. 200.—Portion of filament of *Œdogonium*. *w* in *A*, the cushion of cellulose which has lengthened to the piece of cell-wall, *w'* in *B*; *c*, cell-caps (magnified).

Œdogonium is also reproduced by *resting-spores* (Wille, Bot. Gesell. Stockholm, Sept. 26, 1883; see Bot. Centralbl., xvi., 1883, p. 215).

The sexual reproduction of Œdogonium still shows a high degree of differentiation of the male and female elements. The *antherozoids* are very similar in form to the zoospores, but much smaller, and they are provided with a similar tuft of cilia. The *antherids* are cells belonging to ordinary filaments, but shorter and not containing so much chlorophyll

as the rest, lying either singly among the ordinary vegetative cells, or sometimes in groups of as many as twelve. In most species each antheridial cell divides either horizontally or vertically into two 'special mother-cells,' each of which gives birth to an antherozoid. The *oogones* are developed either in the same filament as the antherids or not, some species being monœcious, others dicœcious. They are also frequently in groups of from three to six. Their development always takes place out of



FIG. 201.—I. *A*, filament of *Œdogonium ciliatum* Hass.; *n*, zoosporange; *og*, oogone with 'dwarf male,' *m*. *B*, oogone at the moment of impregnation; *o*, oosphere; *z*, antherozoid; *m*, 'dwarf male.' *C*, *Œ. gemelliparum* Hass.; piece of filament in which zoospores, *z*, are being formed, from which the 'dwarf males' are produced (× 250). II. Zoospores; *A*, still within the zoosporange; *B*, in the act of escaping; *C*, free zoospore. (After Pringsheim.)

the upper half of the lower portion of a cell provided with cell-caps at its upper end, which has just divided, and which, directly after the division, swells up into a spherical or ovoid form. Immediately before impregnation the protoplasm contracts into an *oosphere*, containing in one portion densely crowded chlorophyll-grains, and, at the spot opposite to the part of the wall of the oogone which is to open, a hyaline 'receptive spot.' The oogone opens in several ways. Sometimes an oval orifice is

formed at the side, through which the colourless portion of the protoplasm protrudes in the form of a papilla which takes up the antherozoids. In other cases the oogone splits in the same way as the zoosporanges, throwing back a kind of lid ; through the lateral crevice exudes some colourless mucilage in the form of a beak-like canal, through which the antherozoids enter, and coalesce with the hyaline portion of the oosphere. Immediately after impregnation the *oosperm* invests itself with a cell-wall, and assumes a brown colour, still remaining within the oogone, which separates from the other cells of the filament, and falls to the ground. where the oosperm passes a period of rest before germination as a *hypnosperm.*

FIG. 202.—*Bulbochæte setigera* Ag. *B*, unicellular antheridial plant. *A*, *C*, young bicellular plants. *D*, mature plant with oogone, *o*, and 'dwarf male,' *dm* (× 400). (After Cooke.)

In some species the mode of fertilisation is more complicated. Peculiar zoospores known as *androspores* are produced non-sexually in special cells of the parent-plant, similar to those which give birth to the antherozoids, only that there is in their case no preliminary formation of 'special mother-cells.' These androspores, which closely resemble the antherozoids in form and size, fix themselves after swarming to a definite spot on the female plant, on or near an oogone, producing very small male plants, which are known as 'dwarf males' or *micrandres.* Each of these consists of two or three cells, the uppermost of which is an antherid. This gives birth to one or more antherozoids, which escape

by the lifting up of a lid, and which impregnate the oospheres in the usual way.

When the hypnosperm germinates after a lengthened period of rest, it does not immediately develop into a new plant, but breaks up into several zoospores, usually four ; these give birth to several generations of non-sexual plants, until the cycle is completed by the production of antherids and oogones. The sexual plants, however, especially the female ones, produce zoospores as well.

The species of *Bulbochæte* (classed by some writers with Coleochæte) are minute plants growing in fresh water, and differing from Œdogonium in having branched filaments ; the terminal cells of the branches ending in long hyaline bristles, which are swollen at the base. The modes of reproduction correspond closely to those in Œdogonium. Wittrock includes also Coleochæte and Sphæroplea among the Œdogoniaceæ.

LITERATURE.

Braun—Verj. in der Natur, 1851 (Ray Soc. Bot. and Phys. Mem., 1853).
De Bary—Ueber Œdogonium u. Bulbochæte, 1854.
Pringsheim—Jahrb. wiss. Bot., 1858, p. 1.
Carter—Ann. and Mag. Nat. Hist., 1858, p. 29.
Juranyi—Pringsheim's Jahrb. wiss. Bot., 1873, p. 1.
Wittrock—Prodr. Monogr. Œdogoniearum, 1874.
Wille—Pringsheim's Jahrb. wiss. Bot., 1887, pp. 443 and 454.

ORDER 3.—SPHÆROPLEACEÆ.

This order comprises at present the single species *Sphæroplea annulina* Ag., the simplest of the class, found occasionally on flooded fields. The filaments are cylindrical and unbranched, and are composed of cells which vary in their comparative length and breadth to an extraordinary degree ; sometimes the length will hardly exceed the breadth, while in other cases it may be as much as ninety times as great. The transverse cell-walls are of great thickness, their surface is irregularly wavy, and they swell out here and there into great 'beams' and excrescences of cellulose from both the lateral and longitudinal walls. During the development of these septa, orifices are formed in them for the passage of the antherozoids. The cells contain a large number of chromatophores and starch-grains, as well as, when mature, a considerable number of small nuclei.

The sexual reproductive elements are *oospheres* and *antherozoids*, formed in different cells of the same filament, which may therefore be regarded as rudimentary *oogones* and *antherids*. A filament may consist of

only two cells, and then one becomes an oogone, the other an antherid. When the number of cells is greater, the oogones and antherids sometimes alternate with one another, but this is not always the case. The contents of an oogone break up into several spherical oospheres, each of which is characterised by a hyaline speck or 'receptive spot.' The antherozoids are produced in extraordinarily large numbers by the breaking up of the contents of an antherid which had previously assumed a brownish red colour. They are furnished with two long slender vibratile cilia, and enter the oogones through the orifices in the transverse walls already mentioned; in their passage they go through remarkable

FIG. 203.—Portion of filament of *Sphæroplea annulina* Ag.; upper cell containing oospheres and antherids, lower cell an impregnated oosperm (× 500). (After Cohn.)



FIG. 204.—*S. annulina*. *A*, young unicellular plant (× 900); *B*, portion of mature filament, showing thick transverse wall and two nuclei, *n* (× 800). (After Rauwenhoff.)

changes of form. The fertilised oospore, or *oosperm*, clothes itself with a thick cuticularised warty membrane, and its contents turn a brick-red colour. It usually hibernates within the oogone in the form of a *hypnosperm*; in the spring its contents break up into three or four zoospores, each of which develops into a slender thread consisting at first of a single fusiform cell which displays no distinction of base and apex, each extremity being elongated into a flagelliform point. The oosphere may also break up into zoospores without previous impregnation.

Probably nearly allied to Sphæroplea, but of somewhat uncertain position, is *Cylindrocapsa* (Reinsch) (Cienkowski, Mél. Biol. Acad. St. Pétersbourg, 1876, p. 534), the mode of reproduction of which is

heterogamous. In other respects it indicates affinity with Ulothrix, and is surrounded by a remarkably thick lamellated gelatinous envelope. It is made by some writers the type of a distinct family, the CYLINDROCAPSACEÆ.

LITERATURE.

Cohn—Ann. Sc. Nat., v., 1856, p. 187.
Heinricher—Ber. Deutsch. Bot. Gesell., 1883, p. 433.
Rauwenhoff—Rev. Internat. Sc. Biol., 1883, p. 176; and Arch. Néerl. Sc. Exact. et Nat., 1887, p. 91.

Class XII.—Fucaceæ.

This family—adopting the limits first proposed by Thuret—consists of a small number of genera of large olive-brown seaweeds distinguished by their mode of sexual reproduction, and by the entire absence, throughout the class, of zoospores, or indeed of any kind of non-sexual spore.

The thallus or 'frond' is often several feet in length, cylindrical or flattened, or, in Himanthalia (Lyng.), cup-shaped, of a cartilaginous texture, and is attached to the sea bottom by a branched *rhizoid* or attachment-disc. This organ is altogether superficial, and has no function in the absorption of food-material, like the root of higher plants. It is formed entirely of filaments originating from the stipe or stem. In some species detached branches have the power of maintaining their existence, and even multiplying for an indefinite period, floating on the surface. Nearly all the species are perennial. Although the thallus does not display the same amount of external differentiation into 'stipe' and 'frond' as some of the Laminariaceæ, the differentiation of internal tissues is quite as great. In the centre of the thallus is a *medullary* system composed of elongated cells, and surrounded by a *cortical* system of shorter nearly isodiametrical cells; there is only a very rudimentary development of epiderm. The thallus increases in thickness by the radial division of the outermost rows of cells or *hyphæ* of which the cortex is composed. Growth in length is entirely apical, taking place, according to the most recent observations, by the segmentation of a single well-marked four-sided *apical cell*, which may be seated, as in Fucus furcatus (Ag.), at the base of a depression at the apex of the frond. Grabendörfer states (Bot. Zeit., 1885, pp. 609 *et seq.*) that in Durvillæa (Bory) there is no apical growing point. The thallus of the Fucaceæ always branches dichotomously and monopodially, the branches lying, when not disturbed, in a single plane. In the more highly



FIG. 205.—*Himanthalia lorea* Lyng. (natural size). (After Hauck.)

FIG. 206.—*Halidrys siliquosa* Lyng. (natural size). (After Hauck.)

developed genera, such as Sargassum (Ag.) and Durvillæa, there is a more or less advanced differentiation of *lamina* or leaf from *stipe* or stem; and the 'leaves' are even arranged spirally and furnished with a rudimentary midrib.

The cell-wall often consists of two distinct layers, an inner firm compact but thin layer, and an outer gelatinous one which swells greatly in fresh water, filling up the intercellular spaces, and causing the slimy character which the Fucaceæ commonly assume after they have lain for some time in fresh water. The cells contain chlorophyll; but the green colour is in all cases entirely masked by a pigment of a brown or olive colour, which can be extracted from the dead plant by cold fresh water. The nature of this pigment has been investigated by Millardet, Rosanoff, Sorby, Schütt, and others. According to Millardet, an olive-green alcoholic extract may be obtained from quickly dried and powdered specimens. If this is then shaken up with double its volume of benzine and allowed to settle, the upper layer of benzine is coloured green by having taken up the chlorophyll, while the lower alcoholic layer is yellow, and contains *phycoxanthin* in solution. Thin sections of the thallus, after complete extraction by alcohol, still yield a reddish brown substance which in fresh cells adheres to the chlorophyll-grains, but can be dissolved out with cold water, especially if the frond be first reduced to powder. To this reddish brown substance, the spectrum of which has one absorption band between E and F, Millardet gives the name *phycophæin*. Schütt states that the spectrum of phycophæin has no characteristic absorption bands, but a regular increase of absorption from the red towards the blue end. He proposes to limit the term phycophæin to the portion soluble in water, and phycoxanthin to that soluble in alcohol, while the entire compound pigment of the Fucaceæ and Phæosporeæ he would call *phæophyll*. Sorby applies the term *fucoxanthin* to the principal colouring matter of the olive-green seaweeds (Fucaceæ and Laminariaceæ). It is soluble in bisulphide of carbon, imparting to it a beautiful amber colour; its spectrum shows two obscure absorption-bands in the yellow. Hick (Journ. of Bot., 1885, pp. 97 and 354) has detected continuity of protoplasm in several species of Fucus, both in the cortical layers and in the central medullary tissue. The intercommunication of the protoplasmic contents of contiguous cells is also effected, as in the Laminariaceæ (see p. 244), through structures of the nature of sieve-plates.

In many of the Fucaceæ *air-bladders* are formed in the frond by the hollowing out of large cavities in the interior of the tissue, which serve to float the frond in the water, and to aid in the process of fertilisation. These are especially noticeable in the common 'bladder-wrack'



FIG. 207.—*a*, *Fucus vesiculosus* L. ; *b*, *F. serratus* L., with their attachment-discs united in growth (natural size). (After Kützing.)

of our coasts, Fucus vesiculosus (L.), and in other species of the genus. In the 'gulfweed' of the 'Sargasso Sea' (Sargassum bacciferum, Ag.) these bladders are spherical, and are elevated on pedicels above the surface of the frond, giving them the appearance of berries (fig. 211). In Halidrys (Grev.) they are ovoid, segmented, and pod-like in appearance (fig. 206).

The only reproductive organs of the Fucaceæ are sexual, *antherids* and *oogones*. Both kinds are formed in globular cavities known as *conceptacles*, which are either distributed uniformly over the thallus, or are congregated in particular portions of it, which are then known as *receptacles*. These are always either the terminal portions of branches, or are in the parts provided with air-bladders, so as to be elevated above



FIG. 208.—Section of male conceptacle of *F. vesiculosus*, clothed with branched hyphæ bearing the antherids; *o*, ostiole (magnified).

the surface of the water for the purpose of fertilisation. In Fucus (L.) the receptacles constitute the warty extremities of the branches; in Himanthalia (Lyng.) the whole of the long whip-shaped stem which proceeds from the cup-shaped thallus is a receptacle (fig. 205); in Sargassum (Ag.) they occupy distinct fertile branches. In some species there are separate *male* and *female conceptacles*, and then they are always dicœcious; in other species antherids and oogones are contained in the same conceptacle, the female organs occupying the lower, the male organs the upper part of the cavity. There are also sometimes cavities of precisely similar structure, but producing no sexual organs, which may be degraded or aborted conceptacles. Both the fertile and barren conceptacles are clothed internally by a dense weft of loose hyphæ, which

are a prolongation of those of which the thallus is composed, and frequently project, through the mouth or *ostiole* of the conceptacle, into the surrounding water. When infertile these hyphæ are known as paranemes or *paraphyses*. In the male conceptacles they are usually branched, unbranched in the female. Both the barren and fertile conceptacles are always first formed in the neighbourhood of the growing point, the cavity originating from the absorption of a row of cells at right angles to the surface.

The *antherids* are produced on lateral branches of the hyphæ in the male or in the upper part of bisexual conceptacles. Each consists of an ovoid thin-walled or sometimes double-walled cell, the abundant protoplasm of which breaks up into a number (usually sixty-four) of



FIG. 209.—Section of female conceptacle of *F. vesiculosus*, clothed with unbranched hyphæ bearing the oogones; *o*, ostiole (magnified).

minute *antherozoids*, pointed at one end, with a pair of cilia of unequal length placed laterally below the beak-like apex, and contains an orange-red pigment spot and a nucleus. The olive-brown *oogones* are developed from unbranched hyphæ in the female, or in the lower part of bisexual conceptacles. These fertile hyphæ are at first unicellular, and are bounded at the base by a septum; the single cell subsequently divides into a basal *pedicel-cell*, and an upper portion, which swells into a spherical or ellipsoidal form, the oogone, filled with protoplasm coloured brown by phycophæin, and always provided with a wall composed of two layers. Either the whole of the contents of the oogone contract into a single *oosphere*, or it divides into two, four, or eight oospheres, each with its own nucleus. Impregnation always takes place outside

the conceptacle. The outer layer of the double wall of the oogone bursts, the inner layer still continuing to enclose the oospheres in a thin bladder-like membrane. In this form they escape from the conceptacle through the ostiole into the surrounding water, where the remaining membrane is also absorbed. In the meantime the antherids have become detached, the inner layer of the double cell-wall having burst through the outer layer, and collect in large numbers before the ostiole of the female or of the bisexual conceptacles, forming orange-red masses which are often caught by the paraphyses which hang out from the ostiole, or are left on the shore at low tide. On the return of the tide, or after they have remained for a time entangled in the paraphyses, the inner membrane of the antherid also becomes absorbed, and the antherozoids escape at the same time that the oospheres become released from their

FIG. 210.—*F. vesiculosus*. *A*, branched hypha bearing antherids (× 160). *B*, antherozoids (× 330). *I*, oogone, *Og*, containing eight oospheres; *p*, unbranched hyphæ. *II*, oospheres preparing to escape; *a*, outer, *i*, inner layer of cell-wall of oogone. *III*, oosphere surrounded by antherozoids. *IV*, *V*, stages in germination of oosperm (× 160). (After Thuret.)

enveloping membrane. The antherozoids frequently collect round the oospheres in such numbers that the motion of their cilia imparts to the comparatively very large passive oosphere a rolling movement which lasts for about half an hour, when they become absorbed into it and impregnate it. The oospheres are receptive over their whole surface; and, although it has been calculated that the bulk of an oosphere is equal to that of from 30,000 to 60,000 antherozoids, an oosphere can apparently be fertilised by a single antherozoid. In this family the mode of repro-

duction consisting in the impregnation of a passive oosphere by motile antherozoids attains its highest development among Algæ. The antherozoids retain their motility and vitality for from one to three days. The oospheres will show signs of a rudimentary germination even when unfertilised, but in that case the germ soon perishes. Thuret succeeded in obtaining a hybrid Fucus by impregnating the oospheres of F. vesiculosus (L.) by the antherozoids of F. serratus (L.).

A short time after impregnation the *oosperm* invests itself with a cell-wall, fixes itself to some other body, and begins to germinate without any intervening period of rest. The first transverse division of the young germinating filament is followed by others in various directions, so that a solid mass of pseudo-parenchyme is at length formed, fixed to the bottom by a root-like rhizoid.

FIG. 211.—The gulfweed, *Sargassum bacciferum* Ag. (natural size).

The Fucaceæ constitute a small and well-marked family of seaweeds, united by some systematists with the Phæosporeæ, or at least with the Laminariaceæ, to make up the Fucoideæ of Agardh, or the Melanospermeæ of Harvey. They are, however, well distinguished by their mode of reproduction. The family is represented in Britain by the genera Halidrys (Grev.), Cystosira (Ag.), Pycnophycus (Ktz.), Fucus (L.), Ascophylla (Stackh.), and Himanthalia (Lyng.), and includes also the exotic genera Sargassum (Ag.), Pelvetia (Dcne.), Durvillæa (Bory), Splachnidium (Grev.), and a few others. Although the number of native British species described by Harvey is only thirteen, some of these occur in such vast quantities that the Fucaceæ

cover a larger amount of surface of tidal rocks than all the other seaweeds together. Among these may be especially mentioned the familiar bladder-wrack, Fucus vesiculosus, so abundant on all our coasts. The well-known gulfweed, Sargassum bacciferum, distinguished by its berry-like air-bladders, a native of warmer seas, is sometimes thrown up on our shores, where it is carried by the gulf-stream. It very rarely fructifies ; detached pieces, buoyed up by the air-bladders, being able to retain their vitality for an indefinite length of time. An enormous floating mass of this seaweed, consisting entirely of detached pieces, is said to cover an area of 200,000 square miles in the Atlantic, about lat. 20–25° N. and long. 40° W., where it has maintained itself with but little shifting since the time of Columbus, affording a home and breeding-place for countless numbers of marine animals. The family is, however, chiefly European ; a large proportion of the species live only in shallow water, being exposed at every ebb-tide, or only at neap-tides, when fertilisation takes place. The distinctions between the different genera are made to rest on the disposition of the air-bladders and conceptacles, and on the more or less distinct differentiation of the leaf-like organs. The frond of Splachnidium is partially gelatinous. The structure of that of Durvillæa, one of the largest of seaweeds, is very beautiful, being permeated by very large and regular cavities resembling a honeycomb. Together with the Laminariaceæ, our native Fucaceæ are largely used in the manufacture of kelp, though not to the same extent as formerly, and as a source of iodine ; they are also employed by farmers as a manure for their fields. On the coast of Chili the poorer classes use a species of Durvillæa for food, and a soup is made from it which is mucilaginous and sweet.

LITERATURE.

Agardh—Species, Genera, et Ordines Fucoidearum, 1848.
Thuret—Ann. Sc. Nat., 1854, p. 195.
Pringsheim—Monber. Berlin. Akad. Wiss., 1855, p. 133.
Rosanoff—(Pigment) Mém. Soc. Sc. Nat. Cherbourg, 1867, p. 145.
Millardet—(Pigment) Comptes Rendus, lxviii., 1869, p. 462.
Kraus et Millardet—(Pigment) Mém. Soc. Sc. Nat. Cherbourg, 1870, p. 23.
Kny—Bot. Zeit., 1872, p. 699; and 1875, p. 450.
Sorby—(Pigment) Proc. Roy. Soc., 1873, pp. 455 *et seq.*
Reinke—Jahrb. wiss. Bot., 1876, p. 399 ; and Bot. Zeit., 1877, p. 651.
Rostafinski—Beitr. z. Kenntniss d. Tange, 1876.
Thuret & Bornet—Etudes phycologiques, 1878.
Kuntze—(Sargassum) Engler's Bot. Jahrbuch, 1880, p. 191.
Bower—(Conceptacle) Quart. Journ. Microsc. Sc., 1880, p. 36.
Berthold—Die Cystoseiren (Fauna u. Flora Golfes Neapel), 1883.
Hanstein—Sitzber. Phys.-Med. Gesell. Würzburg, 1884, p. 104 ; and Arbeit. Bot. Inst. Würzburg, 1885, p. 289.

Dodel-Port—(Cystosira) Biolog. Fragmente, pt. i., 1885.
Behrens—(Fertilisation) Ber. Deutsch. Bot. Gesell., 1886, p. 92.
Schütt—(Phycophæin) *ibid.*, 1887, p. 259.
Woodworth—(Apical Cell) Ann. of Bot., i., 1888, p. 203.

Class XIII.—Phæosporeæ.

The Phæosporeæ or Phæozoosporeæ form, together with the Fucaceæ, the whole of the olive and brown seaweeds of the globe, formerly grouped together under the names Fucoideæ, Melanosporeæ, or Melanospermeæ ; but of many the history of development is at present but imperfectly known ; and when this is ascertained more fully, they may possibly be separated into groups having but little affinity with one another. A number of the Phæosporeæ are epiphytic, and a few parasitic on other seaweeds ; a very few grow in fresh water.

The ordinary mode of multiplication of the Phæosporeæ is, so far as is known at present, non-sexually by means of *zoospores*, which occur in all the orders except the most aberrant groups—the Dictyotaceæ, where they are replaced by motionless spores, and the Syngeneticæ. In the Sphacelariaceæ there is another mode of non-sexual propagation by means of gemmæ or *propagules*. Each zoospore has a large red pigment-spot and two cilia, a longer one pointing forwards and a shorter one directed backwards. They differ from those of the green Algæ, such as the Confervoideæ, in the lateral insertion of the cilia at the base of the colourless apex. They are produced in *zoosporanges*, which are either external, when they are usually the terminal cells of short branches, or are imbedded in the thallus, in which case they are frequently collected into definite groups or *sori*, and are interspersed with barren filaments or hyphæ, known as paranemes or *paraphyses*. These are often swollen and club-shaped at their apex ; the zoosporanges sometimes spring as lateral branches from similar filaments. The zoosporanges are of two kinds, *unilocular* and *multilocular* (the 'oosporanges' and 'trichosporanges' respectively of Thuret). The former are comparatively large, nearly spherical, ovoid, or pear-shaped, and their contents break up directly into a large number of zoospores which escape through a terminal or lateral opening. The latter kind have somewhat the appearance of jointed hairs, and are segmented in the transverse direction only ; or less often are more like the unilocular zoosporanges in form, but are divided internally by both transverse and longitudinal septa. Each cell gives birth to a single zoospore ; and these either escape each separately

from its own mother-cell, or an opening is formed at the apex of the sporange through which all the zoospores escape after dissolution of the septa. The zoospores are in all cases imbedded in mucilage ; no differ-



FIG. 212.—*Giraudia sphacelarioides* D. and S. *a*, upper portion of thallus (× 250) ; *b*, lower portion with multilocular sporanges (× 250). (After Areschoug.) *c*, portion of filament with unilocular sporanges (× 600). (After Hauck.)

ence is observable in size or form between those produced in the two kinds of sporange, but those from the unilocular sporanges appear in all cases to germinate directly, while those from the multilocular sporanges are sometimes *zoogametes* with sexual functions. The two kinds of sporange may be borne on the same or on different individuals ; in the former case they are occasionally developed at different times. In certain orders or groups one or the other kind is altogether wanting.

The various modes of sexual reproduction known in the Phæosporeæ present a most interesting gradual transition from the conjugation of equivalent motile zoogametes to the impregnation of a quiescent oosphere by motile antherozoids. In Ectocarpus (Lyng.), Giraudia (D. and S.), and Scytosiphon (Ag.) conjugation takes place between swarm-cells from the multilocular sporanges, which are to all appearance exactly alike ; but a slight sexual differentiation is exhibited in the fact of one of them coming to rest and partially losing its cilia before conjugation takes place. In Cutleria (Grev.) and Zanardinia (Nard.) the differentiation is more complete. The male and female swarm-cells are produced either on the same or on different individuals ; the latter are much larger than the former, and come perfectly to rest, entirely losing their cilia before being impregnated by the former. In Dictyota (Lmx.) the differentiation is carried still further, and the female reproductive bodies are from the first motionless masses of protoplasm not provided with cilia. In Dictyosiphon (Grev.) (Punctariaceæ) a different kind of conjugation has been observed.

The degree and mode of development of the thallus differ very widely within the class. A few species of Ectocarpaceæ, belonging to the genera Streblonema (Derb.) and Ectocarpus (Lyng.), are microscopic. Some of the Mesogloeaceæ and Ralfsiaceæ are small seaweeds epiphytic on those of larger growth, with a flat radiating thallus reminding one of Coleochæte. In some of the Ectocarpaceæ the thallus consists of simple branched or unbranched filaments resembling those of the Confervaceæ. In the Sphacelariaceæ each branch is composed of a row of larger central surrounded by a layer of smaller cortical cells, all originating from a large uncorticated apical cell. In the Cutleriaceæ filaments of cells become separated from the margin of the thallus, the basal portions of which are coalescent into a solid tissue, the increase in breadth of which is due to the branching of the filaments. The Laminariaceæ include, in the genera Alaria (Grev.), Laminaria (Lmx.), Macrocystis (Ag.), and others, the most gigantic of marine organisms, in which the thallus or 'frond' is to a certain extent differentiated externally into *rhizoid* or organ of attachment, *stipe* or stem, and

leaves. We have here also an approach to the internal differentiation of tissues which occurs in the higher plants, though this is not so strongly displayed as in the Fucaceæ. In Macrocystis, however, the lamina of the frond may be divided into an epidermal layer, cortical parenchyme, and medullary tissue. Sieve-hyphæ occur in all the genera, and in Macrocystis and Nereocystis (Post.) true sieve-tubes with sieve-plates and deposit of callus. Through the perforations in the sieve-plates Hick (Journ. Bot., 1885, p. 356) and Wille have detected the passage of strings of protoplasm connecting the cells with one another. The protoplasts of the cortex in Laminaria digitata (Lmx.) are described by Hick as rhizopod-like bodies spreading in such a way that the cells of each layer are brought into connection both with one another and with those of adjacent layers. The cells of the Phæosporeæ contain a carbohydrate closely resembling starch, but differing in not being coloured blue by iodine, and an olive-brown pigment soluble in cold fresh water identical with the *phycophæin* of the Fucaceæ. The tissues both



Fig. 213.—*Macrocystis pyrifera* Ag. (greatly reduced).

of the Laminariaceæ and of other large marine Algæ belonging to other groups display remarkable elasticity or other properties to enable them to resist the traction of the waves. In the larger species the frond is buoyed up by air-bladders.

Janczewski describes the occurrence in the class of three distinct modes of growth, viz.— (1) The thallus and all its ramifications terminate in a generative apical cell which divides in a direction parallel to its base, and thus gives birth to a series of segments. This occurs in the Sphacelariaceæ and in Dictyosiphon, but is the least common mode. (2) By peripheral growth, i.e. the marginal cells of the thallus are the youngest, and are more or less united into a generative peripheral zone (Myrionema, Grev., Leathesia, Gray, Ralfsia, Berk.). (3) By intercalary growth. This is much the most common mode, and there are, again, three modifications of it, viz. —1. The thallus terminates, when young, in one or more hairs, the common growing point of the thallus and of the hairs being situated at their point of junction (Ectocarpus, Desmarestia, Lmx., Carpomitra, Ktz., Cutleria, Sporochnus, Ag.). 2. The thallus is differentiated into three 'organs'—frond, stipe, and rhizoids; the growing point from which the stipe and frond originate is common to these two organs, while the rhizoids increase by apical growth (Laminariaceæ). 3. The absolutely undivided thallus is regenerated from the growing point situated at the base of the frond (Scytosiphon, Chorda, Punctaria, Grev., Asperococcus, Lmx.).

FIG. 214.—*Lessonia fuscescens* Bory (greatly reduced).

LITERATURE.

Magnus—Festschr. Gesell. naturf. Freunde, Berlin, 1873.
Areschoug—Bot. Notis., 1873.
Gobi—Bot. Zeit., 1877, p. 425.
Reinke—*Ibid.*, p. 441.
Thuret & Bornet—Etudes Phycologiques, 1878.



Fig. 215. – *Laminaria saccharina* Lmx., with rhizoids. *s*, portion of frond which produces zoospores (reduced ½). (After Reinke.)

Falkenberg—Mittheil. Zool. Stat. Neapel, 1878, p. 531.
Wille—Bot. Sällsk. Stockholm, Nov. 19, 1884 (see Bot. Centralbl., xxi., 1885, pp. 282 *et seq.*).

In so many of the Phæosporeæ the life-history is at present but imperfectly known, and different authors differ so widely as to the best characters to be employed in classification, that no attempt is here made to arrange into orders all the known forms. A description is given only of the best-marked groups.

The Laminariaceæ include many of the largest of the brown seaweeds of both warmer and colder seas; in the southern hemisphere they form dense submarine forests of gigantic size, frequently making even deep water impassable for boats, and forming a home for myriads of marine animals; the individual 'fronds' sometimes attaining a length of several hundred feet. The thallus is coriaceous, is not articulated, and is usually attached to the sea-bottom by *rhizoids* or root-like organs of attachment, or less often by a discoid expansion, from which springs a tough cylindrical *stipe* or stem, the tissue of which is more or less differentiated into a medullary portion, an internal and an external cortical portion, and an epidermal portion, the cells of which are coloured brown by phycophæin. It increases in length by intercalary growth at the junction of the stipe and *lamina*. Although most of the larger species are perennial, Areschoug states that even the largest species of Nereocystis (Post.) are annual. In others the stem increases in girth from year to year, attaining sometimes the thickness of a man's thigh. In Chorda filum (Stackh.), one of the commonest of our seaweeds, the entire thallus is cylindrical and whip-like, as much as forty feet in length,

FIG. 217.—Longitudinal section of *Macrocystis pyrifera*, showing sieve-tubes, *l*, and sieve-plates, *s*, with callus (× 300). (After Oliver.)



FIG. 216.—*Chorda filum* Stackh. *a*, uppermost and lowermost portion of frond (natural size); *b*, transverse section, showing differentiation into cortical and epidermal layers (× 200). (After Hauck.)

and is septated by transverse divisions; it is constantly dying off at the apex, the growing point lying at its base immediately above the rhizoids. More often the upper part of the thallus is differentiated into branched annual 'leaves' of cartilaginous texture, usually flat, but sometimes tubular, and often ribbed. Lessonia (Bory) grows erect to a great height, and resembles a branching tree with pendent leaves two or three feet long (fig. 214). In Thalassiophyllum (Post.), Agarum (Grev.), and other genera, the frond is beautifully perforated; these perforations are formed from hollow conical papillæ by which the frond is first covered; the tissue diminishes at the apex of the cones, then bursts, and the opening enlarges as the frond grows. In Macrocystis (Ag.) the stalk-like base of each branch of the frond is swollen out into a large pear-shaped air-bladder. In Nereocystis the air-bladder is barrel-shaped, six or seven feet in length, and crowned with a tuft of fronds. Sieve-hyphæ or trumpet-hyphæ with imperfect sieve-plates occur in all the genera; and Oliver has discovered in the comparatively weak stems of Nereocystis and Macrocystis a structure almost identical with that which occurs in the weak climbing stems of many Flowering Plants, true *sieve-tubes* with perfectly formed *sieve-plates* both in the septa and in the longitudinal cell-walls, provided with a true callus-formation (fig. 217).

Zoosporanges of one kind only—the unilocular—are at present known in the Laminariaceæ; these are distributed uniformly over the surface of the thallus or are collected into *sori*, and are interspersed with simple unsegmented club-shaped sterile hairs or *paraphyses*. Of the mode in which the zoospores act as propagative organs very little is known. Areschoug has observed the germination of the zoospores of Chorda tomentosa (Lyng.) after the coherence of two of them by their beaks; but he does not regard this as a true process of conjugation. Gardiner believes that he has detected the conjugation of zoospores in Alaria (Grev.).

Along with the Fucaceæ, the Laminariaceæ are one of the most important commercial sources of iodine. The species of our own shores are employed in the manufacture of kelp. Alaria esculenta (L.) is used by the inhabitants of Scandinavia and Iceland as an article of food, as also are Laminaria digitata (Lmx.) and other species under the name of 'tangle.' The stems of the last-named species are employed for surgical purposes; those of Ecklonia (Hornem.) and others of the larger genera are used as siphons and for making fishing-nets.

LITERATURE.

Reinke—Pringsheim's Jahrb. wiss. Bot., 1876, p. 317.
Areschoug—Observ. Phycol., iii., 1875, iv., 1883, v., 1884; and Acta Soc. Sc. Upsaliensis, 1875, 1883, and 1884.
Will—(Macrocystis) Bot. Zeit., 1884, pp. 801 *et seq.*

Wille—(Sieve-tubes) Ber. Deutsch. Bot. Gesell., 1885, p. 29.
Gardiner—(Conjugation of Zoospores) Proc. Cambr. Phil. Soc., 1886.
Humphrey—(Agarum) Proc. Amer. Acad. Sc., 1886, p. 195.
Oliver—(Sieve-tubes) Ann. of Bot., i., 1887, p. 95.

In the PUNCTARIACEÆ, SPOROCHNACEÆ, and SCYTOSIPHONACEÆ—the limits of which orders are not settled by systematists—the structure of the thallus varies greatly. In Punctaria (Grev.) and Phyllitis (Ktz.) it is flat and leaf-like, from one to six layers of cells in thickness; in other genera it is slender, cylindrical, erect, and more or less branched, the main axis being either solid or hollow, and consisting of a pseudo-parenchymatous tissue, in which the outermost or the two or three outer rows of cells are much smaller than the inner ones. In



FIG. 218.—*Asperococcus bullosus* Lmx. *a*, natural size (after Bornet); *b*, portion of surface with sorus (× 100); *c*, transverse section through thallus and sorus (× 100). (After Kützing.)

Dictyosiphon (Grev.) the 'frond' branches into delicate hairs. In Arthrocladia (Duby) the branches are arranged in delicate whorls. In Scytosiphon (Ag.) the thallus is elongated, cylindrical, and unbranched, but constricted at intervals, and resembles that of Chorda in habit. In other genera, as Asperococcus (Lmx.) and Hydroclathrus (Bory), it is hollow and bladdery. In most of the genera both kinds of zoosporange are known, while in others one or the other has not yet been detected.

Their arrangement, on which the delimitation of the orders is largely made to rest, varies greatly. They may be collected into wart-like sori on the surface of the thallus, appearing sometimes like dark dots uniformly distributed, or they may spring from the branches. In Sporochnus (Ag.) the unilocular sporanges are collected into a peculiar *receptacle*



FIG. 219.—*Sporochnus pedunculatus* Ag. *a*, natural size; *b*, *c*, receptacle containing zoosporanges; *d*, unilocular zoosporange (× 100). (After Kützing.)

consisting of pear-shaped swellings near the extremity of the branches, composed of a dense mass of filaments on which the sporanges appear as lateral branches. Very little is known of the further history of the zoospores. In Scytosiphon lomentarium (Ag.) Berthold ('Mittheilungem Zool. Stat. Neapel,' ii., 1881) has observed conjugation of the swarm-

spores contained in the multilocular sporanges, which must therefore be regarded as zoogametes, and the phenomena are the same as in the Ectocarpaceæ. Areschoug ('Observ. Phycol., iii., 1875) describes a remarkable kind of conjugation—altogether peculiar as far as the brown seaweeds are concerned—in the swarm-spores of Dictyosiphon hippuroides (Lyng.), somewhat resembling that in the Conjugatæ. Two, or sometimes three, of the zoogametes cohere by their apices, and the contents slowly pass entirely into one of them, but only after they have come to rest. Both then become invested with a thin coat of cellulose, and the one into which the endochrome has passed, which may be called the female element, subsequently germinates. In other cases the conjugating zoogametes put out conjugating tubes not unlike those of the Zygnemaceæ. Some of the zoospores also germinate without conjugating.

LITERATURE.

Reinke—Pringsheim's Jahrb. wiss. Bot., 1878, p. 362.

The MESOGLŒACEÆ or Chordariaceæ (Myrionema, Grev., Leathesia, Gray, Chordaria, Ag., Mesoglœa, Ag., &c.) are seaweeds with a gelatinous or cartilaginous thallus of hemispherical or cylindrical outline, variable in size, and forming small gelatinous or slimy cushions or branching tufts on larger seaweeds. Each filament is composed of a vertical central row of cells, surrounded by a 'cortex' of radial rows at right angles to the central row. On these cortical rows are placed the zoosporanges, which are both unilocular and multilocular, concealed within the periphery of the 'frond.' Nothing is known of the conjugation of the swarm-spores; they appear to germinate directly, giving rise to a creeping branched filament, from which the ascending axes subsequently rise.



FIG. 220.—*Chordaria flagelliformis* Ag. Transverse section through thallus, with unilocular zoosporanges (× 200). (After Kützing.)

The ECTOCARPACEÆ constitute an ill-defined group of small, occasionally microscopic, marine (Elachista, Duby, Ectocarpus, Lyng., Giraudia, Derb.) or rarely fresh-water (Pleurocladia, Br.) Algæ, usually

attached in tufts to larger algæ, and resembling in habit the fresh-water Confervaceæ. The thallus consists of segmented more or less branched filaments, either composed of a single row of cells or corticated. The growing point of the filament does not lie at its apex, but at the extremity of a basal portion, the true thallus, the terminal hair-like portion being deciduous. The zoosporanges are of both kinds, and are either external and stalked, or are ordinary cells of a filament, whether terminal or intercalary. Multilocular sporanges are sometimes produced on the

FIG. 221.—*Ectocarpus investiens* Hauck, epiphytic on *Gracilaria compressa*. (× 250). (After Bornet.)



FIG. 222.—Conjugation of zoogametes of *Ectocarpus siliculosus* Ktz. *I*, *a–f*, female zoogamete, gradually losing its cilia. *II*, male zoogametes swarming round female zoogamete. *III*, *a–e*, stages in the coalescence of the male and female zoogametes (× 790). (After Berthold.)

same individual as the unilocular, but at a later period. The swarm-cells which escape from the unilocular sporanges are non-sexual zoospores, germinating directly after coming to rest, and investing themselves with a cell-wall. Those contained in the multilocular sporanges all escape through a single terminal opening, and partake to a certain extent of sexual properties, or become under certain conditions zoogametes.

Goebel has observed their conjugation in Giraudia sphacelarioides (Derb.) and Ectocarpus pusillus (Griff.), Berthold in E. siliculosus (Ktz.). The process is thus described by Berthold. There is no apparent difference between the male and female gametes. The female swarm-spores lose their cilia and come to rest first. They appear to be in a receptive condition only for a few minutes, during which time they seem to exercise an attractive force on the male gametes, which swarm round them until coalescence takes place. The impregnated gamete immediately clothes itself with a cell-wall, and proceeds to germinate. If unimpregnated it will still germinate, though not so rapidly; as also do the male swarm-spores which fail to conjugate; but in this case the resulting new individuals are weakly, and soon perish. This process in the Ectocarpaceæ may be regarded as the first stage between the conjugation of equivalent zoogametes and the impregnation of a passive oosphere by an antherozoid. Wright (Trans. Roy. Irish Acad., 1877, p. 15) has detected on an Ectocarpus a parasitic Chytridium, the zoospores of which he believes to have been mistaken for sexual organs of the host.

LITERATURE.

Askenasy—Bot. Zeit., 1869, p. 785.
Janczewski—Mem. Soc. Sc. Nat. Cherbourg, 1875, p. 97.
Goebel—Bot. Zeit., 1878, pp. 177, 193.
Berthold—Mittheil. Zool. Stat. Neapel, ii., 1881.

The TILOPTERIDEÆ (Tilopteris, Ktz., Haplospora, Kjellm.) are a small and ill-defined family, probably nearly related to the Ectocarpaceæ.

The SPHACELARIACEÆ (Sphacelaria, Lyng., Stypocaulon, Ktz., Chætopteris, Ktz., Cladostephus, Ag.) are all small marine Algæ, mostly parasitic; Chætopteris plumosa (Ktz.) grows on rocks at a considerable depth below the surface. The thallus usually consists of a number of rows of cells united into a pseudo-parenchyme, and often differentiated into an appearance of a 'medullary' row surrounded by 'cortical' tissue. In Cladostephus and Stypocaulon these corticating rows of cells descend to the base of the stem, and form rhizoids or organs of attachment. The zoosporanges are of both kinds, and are usually placed at the ends of special branches, while in Stypocaulon they are axillary; but very little is certainly known about the germination or possible conjugation of the swarm-spores. The apical cell of each branch is uncorticated, and frequently develops into a hollow chamber of considerable size termed a *sphacele*, filled, when young, with dark mucilaginous contents, which at a later stage become watery. Geyler has described two kinds of sexual



Fig. 223.—*Sphacelaria cirrhosa* Ag. *a*, natural size; *b*, branch with propagules, a (× 100); *c*, filament with unilocular zoosporange (× 100). (After Hauck.)

organ, antherids and 'sexual spores' (oospheres), formed within the sphaceles; but Janczewski believes that the supposed antherozoids are in reality the zoospores of parasitic Fungi (Chytridiaceæ), to whose attacks these seaweeds are especially liable. Pringsheim describes two kinds of fructification produced by Cladostephus, one in the autumn, the other in the winter; but Wollny suggests that both the autumnal fructification and the so-called unilocular sporanges may be due to the attacks of parasitic Chytridiaceæ. The Sphacelariaceæ have a strong tendency to multiply by means of buds, gemmæ, or *propagules*. Janczewski describes the mature gemmæ as consisting of a pedicel and



Fig. 224.—*S. cirrhosa*. *a*, filament with propagule, *b* (× 140). (After Reinke.)

three rays diverging above, with a hair springing from the centre of the rays. They become detached like the basidiospores of Fungi, and are constantly being formed afresh.

LITERATURE.

Geyler—Pringsheim's Jahrb. wiss. Bot., 1866, p. 479.
Janczewski—Mem. Soc. Sc. Nat. Cherbourg, xvi., 1872, p. 337; and Ann. Sc. Nat., 1873, p. 253.
Magnus—Zur Morphologie der Sphacelarieen, 1873.
Pringsheim—Abhandl. Berlin Akad., 1873, p. 137.
Rischawi—Algol. Untersuch., i., 1874 (Just's Jahrb., 1874, p. 13).
Wollny—Hedwigia, 1880, p. 65.

The RALFSIACEÆ (Ralfsia, Berk., Lithoderma, Aresch., &c.) are small seaweeds (with the exception of two species of Lithoderma which grow in fresh running water) with crustaceous thallus, attached to stones, rocks, or the shells of molluscs and crustacea, composed of a pseudo-



FIG. 225.—*Lithoderma fatiscens* Aresch. *a*, vertical section of portion of thallus with unilocular zoosporanges; *b*, with multilocular zoosporanges (× 320). (After Hauck.)

parenchymatous tissue of vertical rows of cells. They have both kinds of sporange, collected into wart-like groups or sori on the surface of the thallus, but nothing is known with regard to the function of the swarm-spores.

The small order of CUTLERIACEÆ, comprising the genera Cutleria (Grev.), Zanardinia (Nard.), and Aglaozonia (Zan.), consists of a small number of seaweeds, nearly all natives of warmer seas, although others from colder climates have been erroneously included in it. The thallus is coriaceous or membranaceous, flat, and either erect (Cutleria), or prostrate (Zanardinia), with the peculiarity that the marginal filaments are dissociated in their growth from the rest of the 'frond.' A true sexual mode of reproduction has been observed by Reinke in Zanardinia, and by Falkenberg and Janczewski in Cutleria. The former genus is monœcious, the latter dicœcious. The oogones and antherids are both collected into sori, the former very dark brown, the latter orange-coloured.

The oogones are divided into thirty-two or sixty-four cells, each of which produces an oosphere. The oospheres are at first biciliated swarm-cells or *zoospheres* endowed with active motion, closely resembling the zoospores of other Phæosporeæ, but are larger and variable in form—one of the very few instances known in the vegetable kingdom of the occurrence of such organs. They are said to be destitute of a nucleus. The antherids are also divided into a number of cells, each of which produces two antherozoids, the normal number in an antherid (in Cutleria adspersa, De Not.) being 128. The antherozoids are also biciliated swarm-cells, but smaller than the oospheres; they have each an orange pigment-spot, and are identical in structure with those of the Fucaceæ. The antherozoids do not approach the oospheres until the latter have come to rest and lost their cilia; the absorption of a single antherozoid into the oosphere is then sufficient to impregnate it; it becomes invested with a cell-wall of cellulose, and begins to germinate at once. Thuret states that in C. multifida (Grev.) the oospheres germinate without having been fertilised. The thallus resulting from the germination of the impregnated oospheres is said to be dorsiventral, producing rhizoids on the ventral side only. Zanardinia produces also non-sexual zoospores in unilocular sporanges; and has another non-sexual mode of propagation, by *budding*. In Aglaozonia reptans (Ktz.) the non-sexual zoospores are the only reproductive organ known; and Falkenberg



FIG. 226.—*Cutleria multifida* Grev. (natural size). (After Hauck.)



FIG. 227.—*C. multifida*; transverse section of portion of thallus with a sorus of multilocular zoosporanges in different stages of development (× 330). (After Bornet.)

FIG. 228.—*Zanardinia collaris* Crouan. *a*, oogones; *b*, antherids (magnified). (After Reinke.)

FIG. 229.—Fertilisation of *Z. collaris*. *a*, swarming oosphere; *b*, antherozoids; *c*, coalescence of antherozoid with passive oosphere; *d*, oosperm (magnified). (After Reinke.)

regards this species as probably a non-sexual generation in the cycle of development of Cutleria multifida. The chief argument for placing the Cutleriaceæ among the Phæosporeæ is the occurrence of non-sexual zoospores, but the mode of sexual reproduction indicates a distinct approach to the Fucaceæ ; in this latter class, however, the structure of the antherids is quite different.

LITERATURE.

Janczewski—Mém. Soc. Sc. Nat. Cherbourg, 1872, p. 345, and Ann. Sc. Nat., xvi., 1883, p. 210.

Reinke—Monber. Berlin. Akad. Wiss., 1876, p. 565 ; and Nova Acta Acad. Leop.-Carol., 1878, p. 67.

Falkenberg—Mittheil. Zool. Stat. Neapel, i., 1879, p. 420.

The order DICTYOTACEÆ, comprising, according to Bornet, the eight genera Dictyota (Lmx.), Zonaria (Harv.), Stœchospermum (Ktz.), Lobospira (Thur.), Spatoglossum (Ktz.), Padina (Adans.), Taonia (Ag.), and Dictyopteris (Lmx.), has been united by some writers with the Cutleriaceæ, while by others it has, with much more reason, been erected into a distinct class, of equal rank with the Phæosporeæ. It differs, in fact, from the other orders of Phæosporeæ in several important points. The thallus of the Dictyotaceæ is membranaceous, usually erect flat and leaf-like, seldom ribbed, often beautifully variegated in zones from the presence of prismatic hairs or from incrustation of calcium carbonate. Growth takes place either by means of a single apical cell (Dictyota), or from a group of equivalent cells. Non-sexual organs of propagation are known, and other organs which are probably sexual. The Dictyotaceæ differ, in the first place, from all other Phæosporeæ in the absence of motile ciliated zoospores. The non-sexual *tetraspores* are produced in *tetrasporanges* developed on the margin of the thallus, or in sori on its surface, on special non-sexual individuals. Although the contents of each sporange usually break up into four spores, when they resemble in all respects the tetraspores of the Florideæ, this is not always the case ; occasionally they divide into only two spores, and still less often the entire contents of the sporange escape as a single naked spore. The spores germinate directly, after investing themselves with a cellulose membrane. The presumed sexual organs, *oogones* and *antherids*, are collected into sori in similar situations, but not on the same individuals as the tetrasporanges. Some species are monœcious, others diœcious. The contents of each oogone consist of a single undivided motionless oosphere. The contents of the antherids, on the other hand, divide into a large number of motionless *pollinoids*. Up to the present time, however, actual impregnation of the oospheres by the pollinoids has not been

observed. As the Cutleriaceæ present an approach towards the Fucaceæ, so the Dictyotaceæ may possibly indicate the point of departure of the



FIG. 230.—*Padina Pavonia* Gaill. *a*, natural size; *b*, portion of surface of thallus with tetrasporanges; *c*, transverse section of upper portion of thallus; *d*, of lower portion (× 100). (After Kützing.)

Florideæ, agreeing with that order in the presence of tetraspores and of non-motile pollinoids; but as the oogone presents no indication of even a rudimentary trichogyne, and there is no process analogous to the forma-



FIG. 231. Stages in the formation of the tetraspores in *Padina Pavonia* (magnified). (After Reinke.)

tion of a cystocarp, it seems best, until more is known of the process of fertilisation, to retain the Dictyotaceæ as an aberrant order of Phæosporeæ, with which they also agree best in the nature of their pigment.

LITERATURE.

Reinke—Nova Acta Acad. Leop.-Carol., 1878.
Thuret & Bornet—Etudes Phycologiques, 1878.
Hauck—(Padina) Hedwigia, 1887, p. 41.

The position of the small family of SYNGENETICÆ, as constituted by Rostafinski, is one of great uncertainty. The two genera of which it is composed have generally been regarded as of a very low type of structure, and it is very doubtful whether they are nearly related to one another. The probability seems to be in favour of both genera having been derived from the Phæosporeæ by retrogressive metamorphosis in different directions. Heckel and Chareyre (Journal de Micrographie, 1885) regard Hydrurus and Chromophyton as presenting a connecting link between the Diatomaceæ and the Phæosporeæ.

FIG. 232.—*A*, *Hydrurus penicillatus* Ag. (natural size). (After Cooke.) *B*, zoospore (greatly magnified). (After Lagerheim.)

Hydrurus Ag. consists of a filamentous thallus, attaining sometimes a foot in length, slimy and affixed to a conical disc, and growing in cold fresh running water. The filaments are simple below but branched above, often with exceedingly fine penicillate divisions, filled with a brown or olive endochrome identical with phycophæin. The surface is naked or densely covered with delicate hair-like appendages, which are occasionally fasciculate. The thallus is composed of cells dispersed through the gelatinous matrix; towards the apex of the branches the cells are in close contact with one another, but in the older parts of the thallus they are some distance apart. Each is surrounded by a very delicate membrane, and Lagerheim states that some of them contain pulsating vacuoles. Propagation takes place by means of zoospores of very peculiar form, produced in the branches only, two or four from each cell. When mature the zoospores are tetrahedral, each angle being

prolonged into a slender colourless beak; in one of the angles is a brown chromatophore; and attached to the centre of the opposite side a single short cilium, and near it two pulsating vacuoles, but no pigment-spot. The zoospores appear to germinate directly without conjugation. Lagerheim has also detected, on different individuals from the zoospores, peculiar resting-spores, through the vitality of which Hydrurus remains dormant through the summer and autumn, its active life extending only through the cold season. Hydrurus is placed by Rabenhorst and Cooke among the Palmellaceæ.

Chromophyton Wor. is an epiphytic organism which vegetates and hibernates within the hyaline cells of the leaves of Sphagnum and other aquatic mosses. In this state it consists of unciliated naked masses of protoplasm with pulsating vacuoles, and endowed with an amœboid motion. While still within the cells of the host, these bodies become invested with a delicate cell-wall, multiply by repeated bipartition, and assume the condition of resting-spores, the endochrome being now of a brownish red colour. From these resting-spores are developed zoospores, minute ellipsoidal or nearly spherical bodies, 8–9 μ. long and 4–6 μ. broad, with a single cilium, a contractile vacuole, and a bright yellow or yellowish brown pigment-disc, consisting of a substance apparently identical with the diatomin of the Diatomaceæ. These zoospores are imbedded in a colourless mucilaginous matrix, in which condition they float in large numbers on the surface of the water of bogs in the form of a fine yellow dust. When completely immersed in water, the zoospores are set free from their investing mucilage, and at once begin to swarm. After a time each zoospore develops a second colourless gelatinous envelope, with a tubular opening below, through which it absorbs water. In this encysted condition, having now lost its cilium, it multiplies by bipartition. Although two forms of zoospore have been observed, one much smaller than the other, no process of conjugation has been detected. Cornu describes a second species of Chromophyton with stalked bodies which may be sporanges, and a siliceous coat like that of diatoms. Although in some respects presenting a resemblance to a degraded form of Phæosporeæ, it is possible that Chromophyton may be a stage in the development of some organism belonging to a totally different class. In some respects it may be compared to the Chytridiaceæ among Fungi.

LITERATURE.

Woronin—(Chromophyton) Bot. Zeit., 1880, pp. 625, 641.
Rostafinski—Hydrurus u. seine Verwandtschaft, Krakow, 1882 (Ann. Sc. Nat., xiv., 1882, p. 5).

Cornu—(Chromophyton) Bull. Soc. Bot. France, 1883, p. xciii.
Hansgirg—Oesterr. Bot. Zeitschr., 1884, p. 31.
Lagerheim—Ber. Deutsch. Bot. Gesell., 1888, p. 73.

Phæothamnion Lagerh. (Bot. Zeit., 1885, p. 604) is a fresh-water alga forming brownish yellow tufts on Vaucheria, Cladophora, &c. Certain cells develop into zoosporanges, each of which produces two biciliated zoospores, and the alga has also a palmella condition. Notwithstanding the brown endochrome, and the fact that the zoospores germinate directly and have not been observed to conjugate, Lagerheim places this genus near to Chroolepideæ and Chætophoraceæ, making it the type of a new family, PHÆOTHAMNIEÆ. It may possibly, however, be more nearly related to the Syngeneticæ.

Class XIV.—Conjugatæ.

The Conjugatæ, as defined by de Bary, constitute an extremely well-marked and natural group, composed of the three families *Mesocarpaceæ*, *Zygnemaceæ*, and *Desmidiaceæ*, with no near affinities (except possibly with the Diatomaceæ). The individual is unicellular in most of the Desmidiaceæ ; but in some genera of desmids, and in all belonging to the other two orders, it consists of a filament of cells, which is almost invariably unbranched. The arrangement of the bright green endochrome, in spiral bands, plates, discs, or stars of beautiful symmetry, is altogether peculiar to this group of plants, and renders them among the most interesting and beautiful of microscopic objects. No formation of zoospores occurs throughout the class, and the ordinary mode of vegetative increase is by simple cell-division, and the breaking up of old individuals in the filiform genera into fragments. They retain their power of life through the winter, when under conditions unfavourable to the formation of zygosperms, by the production of *resting-spores*, or single cells which retain for a long period their vitality. These may be either akinetes or aplanospores in Wille's sense of the terms. Gay states (Bull. Soc. Bot. France, 1886, p. 41) that the filaments of Zygnema (Ktz.), especially when growing in dry situations, have a tendency to break up into *cysts*, i.e. fragments which become enclosed in a mucilaginous sheath, resulting from the gelification of the outer layers of the cell-wall. These cysts may preserve their vitality for months, and then, when moisture again penetrates the sheath, they divide by transverse septa, and develop into new individuals. The single cell of the Desmidiaceæ and the filament of the filiform genera is enveloped in a

thin transparent mucilaginous sheath. According to Klebs (Untersuch. Bot. Inst. Tübingen, 1886, p. 333) this sheath is composed of two distinct portions, a homogeneous substance which is but slightly refringent, and a portion which consists of minute rods placed at right angles to the cell-wall. He regards this mucilaginous sheath as entirely independent of the substance of the cell-wall, and derived from the protoplasmic contents of the cell by diffusion through the cell-wall. The same structure probably prevails also in the Confervaceæ and other filiform algæ growing in fresh water. The Desmidiaceæ possess a remarkable power of apparently spontaneous motion, which will be spoken of in detail under that order.

The only sexual mode of reproduction in the Conjugatæ is the *conjugation* of stationary cells, found nowhere else except in some of the Zygomycetes. This consists, in the unicellular genera, of the complete union or fusion of the protoplasmic contents of two individuals ; in the multicellular genera, of the isogamous union of the whole or a part of the contents of *gametes* or non-motile unciliated cells into a *zygosperm* ; conjugation may take place between cells belonging to the same or to different filaments. Whether the two conjugating cells are physiologically equivalent or not will be discussed under the separate orders, and the process described more in detail. Klebahn finds the union of the two nuclei in the zygosperm to take place only slowly in Zygnema (Ktz.) (Zygnemaceæ) ; while in Closterium (Nitzsch) (Desmidiaceæ) they remain distinct even in the mature zygosperm.

The Zygnemaceæ must be regarded as the typical family of Conjugatæ, from which the Desmidiaceæ have probably been derived by retrogression, exhibited, in most cases, by the reduction of the filament to a single cell. The Mesocarpaceæ display an approach to a higher type of sexual reproduction in the more complicated processes connected with the formation of the zygosperm. The reasons for excluding the Diatomaceæ from the Conjugatæ, contrary to the opinion of some writers, and placing them in a totally different group, will be given hereafter.

Many of the Conjugatæ are extremely abundant in fresh water, whether running or stagnant, to which they are almost entirely confined ; some of the filiform species grow also on moist ground and among moss.

LITERATURE.

De Bary—Untersuchungen über die Conjugaten, 1858.
Gay—Essai d'un Monographe des Conjuguées, 1884.
Bennett—Journ. Linn. Soc., xx., 1884, p. 430.
Klebahn—(Zygosperm) Ber. Deutsch. Bot. Gesell., 1888, p. 160.

Order 1.—Mesocarpaceæ.

The species belonging to this family consist of cylindrical unbranched or very rarely branched filaments of elongated cells, in which the chlorophyll is not arranged, as in the Zygnemaceæ, in stars or spiral bands, but in a thin axile plate occupying one diameter in each cell, and containing a number of conspicuous starch-grains ; those of adjacent cells lying usually or invariably in the same plane. Vegetative propagation takes place by the breaking up of a filament into its constituent cells ; sexual reproduction by a process of conjugation, which may take place either between cells of the same or of different filaments.

The ordinary mode of conjugation in the Mesocarpaceæ is that termed *scalariform*, viz. between the several cells of two different filaments. In most species of Mesocarpus (Hass.) this takes place in the following way. When two filaments lie very near one another side by side, each cell of each filament puts out a short protuberance on the side facing the other filament. While these are forming, the greater part, but not the whole, of the endochrome in each cell passes into the protuberance thus formed, a portion being apparently always left behind As soon as the two protuberances meet, the cell-wall becomes absorbed at the extremity of each, and an open tube is thus formed in which the protoplasm of the two conjugating cells coalesces, with expulsion of cell-sap and consequent contraction into a globular *zygosperm*. The zygosperm is not formed in the centre of the short tube, but at one extremity of it, in contact with what may possibly be regarded as the female filament, although the differentiation is doubtful, and in any case exceedingly slight. The zygosperm may be the result of the coalescence of three cells instead of two. The zygosperm is at once separated from the rest of the conjugating tubes, or from the mother-cells, by a septum on either side. In Staurospermum (Ktz.), where the cells are very long and narrow, four cells take part in the formation of each zygosperm. Two of the slender filaments, lying side by side, bend towards one another convexly so as to bring a part of each filament where there is a septum in contact. Both the septa and the longitudinal bounding-walls become absorbed at this spot, and the greater part of the contents of the four cells coalesce into a zygosperm, which is often of a more or less quadrate form, and is again sharply marked off from the four mother-cells by septa at the truncated corners. *Lateral* conjugation also takes place in some species of Mesocarpus between two adjacent cells in the same filament. In this process each of the two cells puts out a horn-like protuberance at the end adjacent to the other cell ; these protuberances bend towards one

another and meet; the septum between them disappears; the endochromes of the two cells draw towards one another and coalesce, with condensation of the protoplasm, in the connecting tube thus formed (fig. 235). In M. neaumensis (Bennett, Journ. Micr. Soc., 1886, p. 15) the



FIG. 233.—*Mesocarpus parvulus* Hass.; stages in the formation of zygosperm (× 750). (After de Bary.)



FIG. 234.—*Staurospermum gracillimum* Hass., with quadrate zygosperms, showing axile plate of chlorophyll in the cells which have not conjugated (× 400). (From nature.)

zygosperm is not formed in the connecting tube, but in one of the conjugating cells. In Gonatonema (Wittr.) *parthenosperms* are said to be formed closely resembling zygosperms, but not resulting from the

coalescence of the contents of two cells. Several species of Mesocarpus frequently put out long connecting tubes between two filaments, which assume a barrel-shaped form, but without any formation of zygosperms. In this condition they closely resemble Mougeotia (de By.).

The most important point in which the Mesocarpaceæ differ from the Zygnemaceæ is in the processes which take place after the formation of the true zygosperm. Immediately after its formation, it divides into two, three, or more cells, the central one only of which is fertile, germinating after a period of rest; the other sterile cells, which are separated from the fertile cell by septa, taking no part in the germination. The germinating cell is therefore here a resting-spore or *hypnospore*, produced non-sexually, and the whole structure is, as Pringsheim points out, a rudimentary *sporocarp*, indicating an approach to the higher classes of Algæ; while the family is, on the other hand, connected with the Zygnemaceæ through the species of Zygnema in which the zygosperm is formed in the connecting tube; and the best writers are by no means agreed as to the limits of the two orders.

FIG. 235.—*Mesocarpus pleurocarpus* de By.; lateral conjugation (× 200). (From nature.)

Wittrock has described the formation of the 'sporocarp' as taking place in three different ways in the Mesocarpaceæ, viz.:—(1) By the tripartition of the zygosperm into a hypnospore and two sterile cells; when the conjugation is lateral, the sterile cells are not separated from one another by the hypnospore, but are permanently united with one another. (2) By quadripartition of the zygosperm; this has been observed only in the case of scalariform conjugation, the sterile cells being arranged two on one side and one on the other side of the hypnospore. (3) The zygosperm is cruciform or H-shaped, and the sporocarp is formed from it by quinquepartition, two sterile cells bounding the hypnospore on each side; this mode also only takes place in scalariform conjugation. Although these characters have been used by Braun and others for the separation of the genera of Mesocarpaceæ, Wittrock regards them as of very little systematic value, since in one species, Mougeotia calcarea (Wittr.), he observed all three modes of reproduction on one and the same filament. In this same species Wittrock also records the formation of *parthenosperms*, precisely resembling the normal zygosperms, but not resulting from any act of conjugation. An outgrowth springs from a filament, but is not met by any corresponding outgrowth from another filament. It is, however, cut off by a septum, and divides into sterile cells and a hypnospore, just as if fecundation had taken place.

In Gonatonema notabile (Wittr.) parthenogenetic hypnospores are formed by tripartition of the contents of an ordinary cell which swells up into the form of a cask.

The phenomenon of reproduction in the Mesocarpaceæ is regarded by some as a rudimentary appearance of an 'alternation of generations.' The sexual generation or oophyte is completed by the production of the zygosperm as the immediate product of fecundation. This does not germinate directly, but its formation is immediately followed by cell-division, or the development of the non-sexual generation or sporophyte; the sporocarp consists of the germinating hypnospore—in the immediate formation of which no process of impregnation took part—and the investing sterile cells or 'pericarp.' Pringsheim, regarding the process of conjugation in the Mesocarpaceæ as representing a distinctly higher type than that in the Zygnemaceæ, divides the process in the former family into two stages. The first stage, to which he applies the term 'copulation,' consists in the simple union of two cells by the absorption of the dividing cell-wall; the second stage is an intimate coalescence of the protoplasmic contents of the conjugating cells, effected by the motility of the chlorophyll-bodies: and this stage he terms 'connubium.' The hypnospore might, indeed, be correctly termed a 'carpospore,' and we have here a point of departure in the direction of a much more highly specialised type of structure. But the complete similarity of the two conjugating cells before conjugation necessitates the retention of the Mesocarpaceæ among the Conjugatæ.

Limiting the order in accordance with the above-named characters, the genera which the Mesocarpaceæ comprise are Mesocarpus (Hass.), Staurospermum (Ktz.). Craterospermum (Br.), and Gonatonema (Wittr.). Several species of Mesocarpus and Staurospermum are not infrequent in stagnant water, especially in moor pools and among Sphagnum. The filaments are not so copiously invested with mucilage and not of so bright green a hue as those of the Zygnemaceæ; Staurospermum capucinum (Ktz.) has a beautiful violet tinge.

Literature.

Wittrock—Algologiska Studier, Upsala, 1867; Om Gotlands och Oelands Sötwasseralger, 1872 (Quart. Journ. Micr. Sc., 1873, p. 123); On the Spore-formation of the Mesocarpeæ, 1878.

Order 2.—Zygnemaceæ.

The individual consists, as in the Mesocarpaceæ, of a filament of cells placed end to end, which is almost always simple and unbranched. The filaments are cylindrical, and the cells of which it is composed often of comparatively large size ; those of Spirogyra crassa (Ktz.) as much as $\frac{1}{240}$ of an inch (125 μ) in diameter, and twice as long as broad. The chlorophyll is arranged in one or more straight (Sirogonium, Ktz.) or more commonly spiral (Spirogyra, Lk.) bands, or in stars placed in pairs in each cell (Zygnema, Ktz.), or occasionally in an axile plate (Mougeotia, de By.), and encloses large starch-grains and a nucleus, often very large and easily discernible even without the use of staining reagents, connected with the parietal protoplasm by radiating threads. In some species of Zygnema the characteristic appearance of the endochrome is assumed only when the cells are about to conjugate. The ease with which some members of this family, especially species of Spirogyra and Zygnema, are cultivated in fresh-water aquaria, and the beautiful arrangement of the endochrome, not only make these algæ extremely striking objects under the microscope, but afford especially good opportunities for observing, in their details, the processes of division of the nucleus and of the cell.[1] It is also in this family that the interesting process of conjugation is most easily followed.



Fig. 236.—*Spirogyra porticalis* Vauch. ; stages in the formation of zygosperm (× 100). (From nature.)

The filaments increase in length by ordinary cell-division ; and single cells, which become very easily detached, are able to develop in this way into new individuals, corresponding, therefore, functionally to the non-sexual spores of fungi. Like the Desmidiaceæ and other floating algæ, they can obtain their nourishment entirely from the water, and increase without any attachment to the substratum. The formation of *resting-cells* and of *cysts* has already been mentioned. The only other known mode of reproduction is by *conjugation* ; but the reproductive organs of other algæ or fungi, which are sometimes parasitic upon or

[1] See Strasburger, *Ueber Zellbildung und Zelltheilung*; also Sachs, *Text-book of Botany*, 2nd English ed. 1882, p. 16.

endophytic in species of Spirogyra and Zygnema, have been mistaken for zoospores.

The usual mode of conjugation is the *scalariform*, between the cells of two filaments lying side by side. The first stage is the putting out of lateral protuberances at right angles to the axis of growth, from the cells in each filament towards the corresponding cells in the other filament. The protuberances put out by opposite cells at length meet; the protoplasm-mass of each of the two cells has by this time begun to contract, withdrawing itself from the cell-wall, and rounding itself off into an ellipsoidal form, with expulsion of some of its cell-sap, the chlorophyll-bodies at the same time losing their characteristic arrangement. The cell-wall then opens between the two protuberances, and the whole of the protoplasmic contents of one of the two cells passes through the connecting tube thus formed, and glides slowly through it into the other cell-cavity, coalescing with its protoplasm-mass. After complete union the combined protoplast is again ellipsoidal or spherical, and scarcely larger than each of the two before coalescence, further expulsion of water and consequent contraction having taken place. In some species of Zygnema the zygosperm is formed, not in either of the conjugating cells, but in the connecting tube, as in the Mesocarpaceæ. In Sirogonium no connecting tube is formed, but conjugation takes place by *genuflexion*; the two filaments are brought into contact by a knee-like bend in each; at the point of contact the adjacent cells of the two filaments are placed in communication by the disappearance of the cell-walls, and a zygosperm is formed in one of the two cells by the coalescence of the two protoplasts.



FIG. 237. *Zygnema peticulatum* Ag., in conjugation (× 100). (After Cooke.)

FIG. 238. — *Spirogyra bellis* Hass.; lateral conjugation (× 100). (From nature.)

A second mode of conjugation, not nearly so frequent, is the *lateral*, between two contiguous cells of the same filament. Protuberances formed near the adjacent ends of contiguous cells bend towards one another till they meet; the cell-wall between them then disappears,

and conjugation is effected though the curved tube thus formed by the coalescence of the protoplasts of the two cells. The zygosperm is never formed, as in the Mesocarpaceæ, in the connecting tube, but in one of the two cells. Lateral conjugation frequently takes place with groups of four cells, the zygosperms being formed in the two central ones.

The protoplast formed by either mode of conjugation finally secretes a cell-wall of cellulose, and becomes a *zygosperm*. Germination sometimes takes place while still within the mother-cell; but most commonly both filaments perish after conjugation, with the exception of the numerous zygosperms, which fall to the bottom, the green endochrome having in the meantime turned to a brick-red colour. It then remains dormant through the winter as a resting-cell or *hypnosperm*, germinating in the spring. Whether the axis of growth of the new individual is parallel or at right angles to that of the old individual is differently stated by different observers. On the commencement of germination, one end of the zygosperm of Spirogyra attaches itself like a root to a stone or to some other alga, so that in the earliest stage of the new individual there is a differentiation between base and apex; but this soon disappears. The innermost of the three layers of which the cell-wall of the hypnosperm is composed bursts through the other two, and protrudes like a bag. The chlorophyll then arranges itself in spiral bands, with starch-grains within them, and the cell divides by transverse septa into a filament, all the cells being from this time precisely alike. Instances are recorded of filaments persisting through the winter. Hofmeister states that the growth of Spirogyra is intermittent, and that the filaments exhibit a nutation, due probably to differences in the rapidity of growth of different sides of the same cell. Where branching takes place, it appears to be confined to the barren portion of a filament in which zygosperms have been formed.

Although the view is contested by some writers, the process of conjugation is regarded by most as a sexual process, but one of the most rudimentary character, the differentiation of the two conjugating elements being exceedingly slight. As de Bary has pointed out—and his statement is confirmed by nearly all more recent observers—the direction of conjugation is clearly governed by some physiological law, the movement of the protoplasm between the two filaments almost invariably taking place in one direction only, so that one of the two conjugating filaments is entirely emptied, while the other is filled with zygosperms.[1] Designating the former as the *male* and the latter as the *female* filament, it is frequently the case that the cells of the

[1] Hassall, however, asserts and figures the contrary (*British Fresh-water Algæ*, i., p. 130).

female are both longer and broader than those of the male filament; and the contraction of the protoplasm has been observed to begin earlier in the male than in the female filament. It is also stated that the protuberance from the female cell is shorter but broader than that of the male cell, the latter fitting into the former as into a socket. The chief argument against the sexuality of the filaments is the occurrence of lateral conjugation; and when this takes place a sexual differentiation can be assumed only of the individual cells and not of the filaments; but that there is some differentiation of this kind would appear from the fact that when lateral conjugation takes place in a group of four cells the zygosperms are formed in the two centre cells, which may be regarded as female. The phenomena may then be compared to those in Sphæroplea. Bessey states (*in lit.*) that scalariform and lateral conjugation may sometimes be seen in different parts of the same filament. The female filaments are, as a rule, very much more abundant than the male; and it is not uncommon for conjugation to take place between one male and several female filaments, while the reverse is at all events much more rare. Occasionally one cell will conjugate with two others, the zygosperm being then the product of one female and two male cells. Several instances are recorded of hybridism between two different species of Spirogyra. Parthenogenesis, or the formation of *parthenosperms* capable of germination, and in all respects resembling zygosperms, but formed out of the contents of a single cell without any previous process of conjugation, is also stated to occur.

The genera included in the Zygnemaceæ with the above characters are Zygnema (Ktz.), Spirogyra (Lk.), Mougeotia (de By.), Sirogonium (Ktz.), and Zygogonium (Ktz.). Several species of Spirogyra and Zygnema are among the commonest of fresh-water Algæ in both stagnant and running water, forming dense bright green masses, often with a slimy feel, owing to the well-developed mucilaginous sheath in which each filament is enveloped. While conjugation is in active progress, which is mostly in the early summer, the filaments of Spirogyra assume a dull green or even brown colour, easily recognised by the naked eye. The other genera are more frequent in moor pools.

Literature.

Pringsheim—Flora, 1852, pp. 465 *et seq.*
Cleve—Monografi Zygnemaceæ, 1868.
Hofmeister—Würtemb. naturw. Jahresheft, 1874, p. 211.
Overton—Ber. Deutsch. Bot. Gesell., 1888, p. 68.
(For fuller bibliography see Bennett, Journ. Linn. Soc., xx., 1884, p. 430).

Order 3.—Desmidiaceæ.

The Desmids are unicellular organisms, for the most part solitary, and inhabiting almost exclusively fresh water, especially stagnant, where they occur in very large numbers; a very few species are brackish. They always float free without any attachment to the bottom or to other algæ, and many species possess a power of apparently spontaneous motion through the water similar to that of diatoms, though not so strongly marked. In several genera, as Desmidium (Ag.) and Hyalotheca (Ehrb.), the individuals are united into long filaments; and either the separate individuals or the filaments are invested by a more or less dense mucilaginous envelope, species of Desmidium and Hyalotheca frequently forming a green slime on the surface of moor pools. The origin and structure of the mucilaginous sheath appear to be the same as in the Zygnemaceæ. According to Hauptfleisch, the cell-wall of desmids always consists of two distinct layers, sometimes of more, which are then provided with a girdle-band similar to that of diatoms. He also states that in nearly all species the cell-membrane is perforated, and that through these pores proceed threads of protoplasm, connecting the protoplasm in the interior of the cell with the gelatinous envelope which is excreted through the pores from the cell-contents. In the filamentous species the protoplasm is probably in connection throughout the filament.

The cells vary greatly in size and form in different species, the largest (Cosmarium, Cord., Micrasterias, Ag.) being just visible to the naked eye. The individual is usually divided by a deep constriction into two symmetrical halves; and even where this is not the case, the cell-contents—chlorophyll-bodies and starch-grains—are symmetrically arranged in the two halves of the cell. The cell-wall is smooth, or punctated, warty, or even elevated into spines, but has no (or very little) deposit of silica. The cells contain a large quantity of chlorophyll of a bright green colour, never concealed by any pigment, often arranged in bands or stars, and containing much starch. Each genus has its own general form of cell, often of very great beauty. In Docidium (Bréb.), Penium (Bréb.), and Tetmemorus (Ralfs) the individual or 'frustule' is elongated, cylindrical, and usually divided by a constriction into two halves placed end to end; in Closterium (Nitzsch) it is crescent-shaped; in Micrasterias it is very thin and flat, usually with a more or less orbicular or elliptical outline, deeply divided into two symmetrical halves, and each half more or less deeply lobed; in Euastrum (Ehrb.) the individual is usually smaller than in Micrasterias, often very minute, and

the half-cells have a smooth, sinuate, or beaked margin, with circular inflated protuberances: in Cosmarium the half-cells are quite undivided, and the whole outline often nearly orbicular; in Xanthidium (Ehrb.) and in most species of Staurastrum (Mey.) the surface is elevated into prominent tubercles or spines.

The transparency of the cell-wall in desmids enables the rotation of the protoplasm to be distinctly seen; and at the colourless spaces at the extremities of some species of Closterium and Docidium the dancing 'brownian' movement of particles suspended in the cell-sap is very



FIG. 239.—A, *Desmidium Swartzii* Ralfs; B, *Micrasterias rotata* Grev.; C, *Euastrum rostratum* Ralfs; D, *Cosmarium cælatum* Ralfs; E, *Xanthidium cristatum* Bréb.; F, *Staurastrum Arachne* Ralfs; G, *Closterium Dianæ* Ehrb.; H, *Docidium baculum* Bréb. (All after Ralfs and variously magnified.)

evident. Klebs describes four kinds of movement in desmids, viz.: (1) A forward motion on the surface, one end of each cell touching the bottom, while the other end is more or less elevated and oscillates backwards and forwards; (2) an elevation in a vertical direction from the substratum, the free end making wide circular movements; (3) a similar motion, followed by an alternate sinking of the free end and elevation of the other end; and (4) an oblique elevation, so that both ends touch the bottom—lateral movements in this position; then an elevation and circular motion of one end, and a sinking again to an oblique or horizontal position. These movements are, according to this observer, all due to an exudation of mucilage, and the first two to the

formation, during the motion, of a filament of mucilage by which the desmid is temporarily attached to the bottom, and which gradually lengthens. The movements of desmids are especially vigorous when they are in the act of dividing. Stahl found that, like the movements of zoospores, they are affected by light.

Vegetative propagation takes place by *division* or *fission*, a process which can be easily followed out in species of Cosmarium or Staurastrum, the whole being completed in the course of a few hours. When cell-division is about to commence, the endochrome retreats slightly from the band or 'isthmus' which connects the two half-cells with one another; and the two halves then separate from one another, retaining their connection only by a transverse band formed by the gradual broadening of the isthmus; this is after a time divided into two by a septum along the length of the isthmus, midway between the two half-cells and parallel to the constriction between them. The endochrome now passes out of each original half-cell into the half of the band in connection with it, and at the same time the half-band bulges, and, growing rapidly, assumes the form and appearance of an original half-cell. Fresh formation of chlorophyll is at the same time taking place in it, and the half-band becomes a complete half-cell, but sometimes slightly larger. We have now two individuals attached to one another by their larger halves; these frequently remain in contact for a considerable period, but at length separate. In the spiny species of Staurastrum the spines are developed very rapidly on the half-bands while their development into half-cells is progressing.



FIG. 240.—*Staurastrum teliferum* Ralfs, dividing (× 400). (From nature.)

A sexual process of *conjugation* takes place in the following way in the genera where the individuals are quite distinct. Two individuals—which cannot in any way be differentiated as male and female—lay themselves either parallel to or across one another, and the pair become enveloped in a common mucilaginous coating. In each individual the outer of the two layers of which the cell-wall is composed gives way, and a circular opening is formed at the constricted part; the inner layer of the cell-wall of each individual protrudes through the opening in the form of a bladder, and these two protrusions come into contact. The outer cell-wall is then thrown off, and the wall separating the two con-

jugating protrusions disappears. The two protoplasmic bodies then unite in the conjugating tube thus formed into a nearly spherical *zygosperm*, enclosed in a cell-wall which ultimately becomes differentiated into three layers, the innermost and outermost of which are colourless, while the middle one is firmer and brown. The outer surface remains in some species smooth, while in many it becomes, when mature, covered with warts or spines, which are not unfrequently barbed. In those genera where the individuals are associated into filaments, conju-



FIG. 241.—Zygosperms of desmids. A, *Euastrum pectinatum* Bréb. (× 400). B, *Penium margaritaceum* Bréb. (× 300). C, *Closterium rostratum* Ehrb. early stage (× 200). D, *Desmidium Swartzii* Ralfs (× 600). (All after Ralfs.)

gation takes place between the cells of different filaments, and a large number of zygosperms may frequently be seen in the same filament. In Gonatozygon (de By.), where the cells are very long and slender, the process is very similar to that in Zygnema. The statement of the occurrence of zoospores in the Desmidiaceæ is founded on erroneous observation, the antherozoids of parasitic fungi having possibly been mistaken for zoospores.

After remaining for a considerable time at rest, the zygosperm germinates by the bursting of its two outer coats, the protoplasmic contents escaping still enveloped in the very thin innermost coat. In this *embryo*,

as it may be termed, the protoplasm and chlorophyll-corpuscles are already distributed symmetrically into two half-cells, which contract somewhat, and the whole becomes invested by a new cell-wall. A constriction has in the meantime made its appearance between the two halves, and the new individual rapidly assumes its mature form, but is at first of small size. It soon divides repeatedly, and each generation gradually increases in size until the full size is attained.

The number of known species of desmids is not large compared with that of diatoms; they are found in great abundance in the midst of larger algæ in fresh water, especially in moor-pools, sometimes forming a green scum on the surface.

LITERATURE.

Ehrenberg—Die Infusionsthierchen, 1838.
Ralfs—British Desmidieæ, 1848.
Nägeli—Gattungen einzelliger Algen, 1849.
Stahl—Verhandl. Phys.-med. Gesell. Würzburg, 1880, p. 24.
Fischer—Bot. Zeit., 1883, pp. 225 *et seq.*
Wolle—Desmids of the United States, 1884.
Klebs—Biolog. Centralblatt, 1885, p. 353.
Cooke—British Desmids, 1887 (which see for further bibliography).
Hauptfleisch—Zellmembran u. Hüllgallerte der Desmidiaceen, 1888.

Class XV.—Confervoideæ Isogamæ.

In this class the individual still consists of a filament of cylindrical cells, placed end to end, which may be branched or unbranched. As in the Conjugatæ, the only known sexual mode of reproduction is an *isogamous* one between two masses of protoplasm, which are not clearly differentiated beforehand into a male and a female element; but the conjugating bodies are not the contents of stationary cells, but are motile ciliated swarm-spores or *zoogametes*, produced by free-cell formation in ordinary or in slightly differentiated cells of the filament, hence termed *gametanges*, their conjugation resulting in the production of a *zygosperm*. The filament increases in length by the repeated transverse septation of successive apical cells, or less often of intercalary cells. The ordinary mode of multiplication is a non-sexual one, by means of naked ciliated *zoospores*, closely resembling the zoogametes, but often larger, and formed singly or in pairs in a cell. Vegetative propagation also takes place by the formation and detachment of *cysts* or resting-cells, which may be either akinetes or aplanospores. The cells very frequently display a plurality of nuclei, but this is not nearly so strongly

marked as in the unicellular Multinucleatæ. The class includes one large order, the *Confervaceæ*, and three smaller ones, the *Ulotrichaceæ*, *Pithophoraceæ*, and *Chroolepideæ*, though the boundaries between them are not in all cases well defined. In the Confervaceæ and Ulotrichaceæ the filament which springs from the germination either of a zoospore or of a zygosperm resulting from the conjugation of zoogametes, attaches itself to the substratum—a stone, another alga, or some other aquatic plant—by a *rhizoid*, which may consist of a single cell, or may branch into a number of cells. As in the higher algæ, the rhizoid is not a nutritive organ, but simply an organ of attachment. These algæ may, however, continue to grow and retain their vitality for a long period in water without any attachment to the substratum.

Order 1.—Confervaceæ (*including* Chætophoraceæ).

The term Confervaceæ has been very vaguely applied to a variety of green fresh-water organisms, but is now limited to a comparatively small number of genera of fresh-water, and a few brackish and salt-water algæ, in which each individual consists of a segmented branched or unbranched filament of cylindrical or disc-shaped cells, invested by a mucilaginous sheath, and in which multiplication takes place non-sexually by *megazoospores*, or sexually by the conjugation of smaller *zoogametes*. Both kinds of swarm-cell have two cilia, or the former in some cases four; Lagerheim describes, in Conferva bombycina (Ktz.), megazoospores with a single cilium. From each parent-cell are produced either one or two megazoospores.



Fig. 242. *Microspora floccosa* Thur. *A*, *B*, portions of filament. *C*, filament dividing for the escape of zoospores. *D*, zoospores (× 300). (After Cooke.)

In only a few species has the process of conjugation of zoogametes been actually observed, and the systematic position of a large number of the species is therefore at present only conjectural. Areschoug has followed both the conjugation of the zoogametes and the direct germination of the megazoospores in Urospora (Aresch.). In Conferva (L.), Chætophora (Schr.), Draparnaldia (Ag.), and some other



FIG. 243.—*Cladophora gracilis* Ktz, *a*, natural size; *b*, upper, *c*, lower portion (× 200). (After Hauck.)

genera, vegetative propagation takes place by means of resting-spores or *cysts*, usually found in swollen barrel-shaped cells. In Conferva the resting-cells may be either akinetes or aplanospores; and Wille believes that they are produced especially under circumstances unfavourable for the formation of zoospores. The resting-spores of Confervaceæ are formed in three different ways: either (1) by rejuvenescence, and the formation of a new cell-wall round the contracting contents; or (2) by separation of a portion of the cell-substance so as to form a swollen part of the mother-cell, and the thickening of the cell-wall at this portion; or (3) by the simple thickening of the wall of the mother-cell. In the formation of aplanospores, one, two, or four proceed from a single cell by the cell-contents rounding off and enclosing themselves in a cell-wall while still within the parent-cell. They hibernate within the parent-cell, and germinate in the spring. *Resting swarm-cells*, naked masses of protoplasm endowed with an amœboid power of motion, are formed in the same way.

The mucilaginous sheath of the Confervaceæ appears to have the same construction as in the classes of algæ already described, but is often but feebly developed. Wille states that the

spores put out an organ of attachment even before they germinate. In Chætophora and other genera which make up the Chætophoraceæ of Hassall, the terminal cell of the main axis or of its branches is prolonged into a colourless hyaline bristle. These are especially well developed in Draparnaldia, an exceedingly beautiful organism not uncommon in

FIG. 244.—*Draparnaldia glomerata* Ag. (× 100). (From nature.)

fresh water, which exhibits a somewhat higher type of development than the other genera, being differentiated into an axis or central tube, and smaller secondary branches arranged in regular whorls; the zoospores being produced in the latter only. Maupas states (Compt. Rend., lxxxix., 1879, p. 250) that the cells of Cladophora contain a large number of nuclei: and Schmitz (Sitzber. Niederrhein. Gesell., 1879) finds four nuclei in a

cell in certain conditions of Conferva. In this, and in the possession of proteinaceous crystalloids (Klein, Bot. Zeit., 1880, p. 782), these genera show an affinity to Siphonocladaceæ. Binuclearia (Wittr.) (Bot. Centralbl., xxix., 1887, p. 60) appears to have always two nuclei in each cell.

There are still many points to be cleared up in the life-history of the Confervaceæ, although some of the genera are among the most abundant of fresh-water organisms; and the bounds and systematic position of the family are still uncertain. According to some observers, many of the species are connected genetically with forms at present placed under the Protophyta; to this view further reference will be made hereafter. Andersson (Bot. Centralbl., xxxv., 1888, p. 351) believes Palmella uvæformis (Ktz.) to be a resting condition of Draparnaldia. Among the genera now included in the order are Conferva (L.), Microspora (Thur.), Cladophora (Ktz.), Rhizoclonium (Ktz.), Stigeoclonium (Ktz.), Chætomorpha (Ktz.), Draparnaldia (Ag.), Chætophora (Schr.), Urospora (Aresch.), and Binuclearia (Wittr.). Phæothamnion (Lagerh.) (see under Syngeneticæ) ought possibly to be included here; as also Spongocladia (Aresch.) (see p. 290). Several species of Cladophora, Chætomorpha, and Rhizoclonium grow in brackish or even in salt water.

LITERATURE.

Vaucher—Hist. des Conferves d'eau douce, 1803.
Areschoug—Nova Act. Reg. Soc. Upsala, vi., 1868, and ix., 1874.
Reinhardt—Arb. Naturf. Gesell. Charkoff, 1876.
Wille—Bot. Centralbl., xi., 1882, p. 113; and Pringsheim's Jahrb. wiss. Bot., 1887, pp. 437, 459, and 492.
Lagerheim—Ber. Deutsch. Bot. Gesell., 1887, p. 409.
Murray and Boodle—(Spongocladia) Ann. of Bot., ii., 1888, p. 169.

ORDER 2 (?).—PITHOPHORACEÆ.

The Pithophoraceæ must be admitted as a distinct order only with very great doubt, both because the mode of sexual reproduction is at present unknown, and because of their strong resemblance to the Confervaceæ. The family, consisting of a single genus, was founded on *Pithophora Kewensis* Wittr., an inhabitant of warm tanks in the Botanic Gardens at Kew, Oxford, and elsewhere; other species have since been found in tropical America. The thallus is composed of branching filaments of cells resembling Cladophora, but increasing only by bipartition of the terminal cell, and presenting here and there barrel-shaped cells very rich in chlorophyll, in which are formed resting-spores of non-sexual origin. These germinate directly, and in opposite directions, from the two apices. There is another mode of non-sexual propagation by 'prolific cells'; but no zoospores nor any sexual mode of

reproduction have as yet been detected. They are distinguished by the remarkable development of their rhizoids or organs of attachment.

LITERATURE.

Wittrock—On the Development and Systematic Arrangement of the Pithophoraceæ, 1877.

ORDER 3.—ULOTRICHACEÆ.

This small order includes the genera Ulothrix (Ktz.), Hormiscia (Aresch.), and perhaps one or two others, not uncommon in fresh and occasionally in brackish water. The life-history of U. zonata (Ktz.) and other species has been investigated by several observers. They exhibit considerable affinity both to the Confervaceæ and to the Hydrodictyeæ. Each individual is composed of an unbranched filament of short cells, broader than long, and nearly uniform in length. Some of the cells are *megasporanges*, giving birth to 2, 4, or 8 megazoospores with 4 cilia; others are *microsporanges* or *gametanges*, producing 16 or 32 biciliated microzoospores or zoogametes. From the non-sexual megazoospores to the zoogametes there is, however, a gradual transition, the only constant difference between them being the number of cilia. Those microzoospores which do not conjugate, as well as the megazoospores, germinate directly, germination sometimes taking place even within the mother-cell. Their escape is, however, sometimes arrested, when they lose their cilia, invest themselves with a thick cell-wall, and assume a *palmelloid* condition. The plants which spring from the germination of the megazoospores are larger than those which spring directly from the micro-



FIG. 245.—*Pithophora Kewensis* Wittr., branching plant; *sp*, spore (× 20). (After Wittrock.)

zoospores, as also than those which spring from the *zygosperms* resulting from the conjugation of the zoogametes. The escape of the swarm-spores was observed by Cramer to take place, usually in the morning, even in water which froze on the surface every night, conjugation following quickly afterwards. The two kinds of swarm-spore are never produced in the same cell, but in different cells of the same filament, and Cramer believes that conjugation takes place between zoogametes from the same filament. The megazoospores use up in their formation the whole protoplasmic contents of the mother-cell, while in the production of the

FIG. 246.—*Ulothrix implexa* Ktz. *a*, vegetative filament (× 480); *b*, portion of the same (× 800); *c*, palmella-condition (× 480); *d*, escape and conjugation of zoogametes (× 800). (After Dodel-Port.)

microzoospores or zoogametes a portion of the contents forms a bladder which escapes with them, but soon perishes. According to Wille (Bot. Centralblatt, vol. xi., 1882, p. 113) Ulothrix also produces *cysts* or resting-spores, which may be either aplanospores or akinetes. In some other members of the order the filament is branched. Schaarschmidt points out that a state closely resembling the microsporiferous filaments of Ulothrix occurs in the development of the Confervaceæ; and

Hansgirg (Bot. Centralblatt, 1885) believes that the filamentous genera placed in this genus are connected genetically with forms classed under the Chætophoraceæ, Siphonocladaceæ, and Ulvaceæ. To this order belong also Hormidium (Ktz.) and Schizogonium (Ktz.). Wildeman (Bull. Soc. Bot. Belg., 1886, p. 7) traces a genetic connection between Ulothrix and Pleurococcus.

LITERATURE.

Cramer—Vierteljahrschrift Nat. Gesell. Zürich, 1870.
Cienkowski—Mél. biol. Bull. Acad. St. Pétersbourg, 1876, p. 531.
Dodel-Port—Pringsheim's Jahrb. wiss. Bot., 1876, p. 417 ; Bot. Zeit., 1876, p. 177.
Gay—Bull. Soc. Bot. France, 1888, p. 65.
Hansgirg—Flora, 1888, p. 259.

ORDER 4.—CHROOLEPIDEÆ.

This order, as constituted by Borzì, comprises a small group of algæ found on damp walls, the trunks of trees, and similar situations, not unfrequently imbedded in the thallus of lichens, or constituting their gonidial element. The thallus consists of a branched or unbranched filament of cells, usually somewhat rounded or moniliform in outline, and is distinguished by the masking of the colour of the chlorophyll by a golden yellow, orange, or red oily pigment, soluble in alcohol and imparting a strong odour of violets. This pigment, which occasionally occurs also in other lowly organised algæ and protophytes, has been examined by Rostafinski, and found to be a derivative of chlorophyll to which he gives the name *chlororufin*. Microzoospores or zoogametes and megazoospores are produced in *gametanges* or zoosporanges, which are indistinguishable from the vegetative cells except by their somewhat larger size, and which are either terminal or intercalary. Conjugation of the



FIG. 247.—*Trentepohlia Bleischii* Rbh. *A*, filament with swollen cells in which the zoogametes are formed (gametanges), *g*; *B*, stages in the conjugation of the zoogametes (× 330). (After Wille.)

smaller swarm-spores has been observed in Chroolepus (Ag.), but they can also germinate without conjugation. Chroolepus also produces resting-spores. Of this genus one species, C. aureum (Ktz.), is common on walls and rocks, and another, C. umbrinum (Ktz.), occurs on the bark of trees. C. Iolithus (Ag.) is one of the few algæ that grow in perfectly dry situations on gneiss &c., and on a siliceous rock in the Hartz Mountains known as 'violet-stone.' The genus Trentepohlia (Mart.) is now merged by many writers in Chroolepus. Wille has pointed out that the organism described as Gongrosira de Baryana (Rbh.), which grows attached to the shell of fresh-water molluscs as a green velvety coating, is a form of Chroolepus or Trentepohlia.

The genera included under the Chroolepideæ by Borzì in addition to Chroolepus and Trentepohlia are Microthamnion (Näg.), Acroblaste (Reinsch), Leptosira (Borz.), Chlorotylium (Ktz.), and Pilinia (Ktz.). Acroblaste grows in salt water attached to mussel-shells, but its position here is doubtful. Leptosira produces zoospores, some of which germinate directly, while others are said to conjugate, but in a manner different from other zoogametes, by the end which does not bear the cilia. To this family probably belongs also Trichophilus (Weber), a remarkable alga parasitic on the hairs of a sloth, which produces two kinds of zoospore (Bot. Centralblatt, vol. xxxiv., 1888, p. 161).

Literature.

Caspary—Flora, 1858, p. 579.
Hildebrand—Bot. Zeit., 1861, p. 81.
Gobi—Bull. Acad. Sc. St. Pétersbourg, 1872.
Schnetzler—Bull. Soc. Vaud. Sc. Nat., 1879, p. 267; 1880, p. 13; and 1883, p. 53.
Borzì—Studi Algologici, 1883.
Wille—Pringsheim's Jahrb. wiss. Bot., 1887, pp. 426 and 484.
Wildeman—C. R. Soc. Roy. Bot. Belgique, 1888, p. 140.
De Toni—Notarisia, 1888, p. 581.

Class XVI.—Multinucleatæ.

In this newly-constituted group are included the four orders of Siphoneæ, Botrydiaceæ, Dasycladæcea, and Siphonocladaceæ, the near relationship of which to one another is scarcely doubtful, although the first displays sexual reproduction of a high type, with strongly differentiated antherids and oogones, which are not found in the other orders. All the orders are also propagated non-sexually by zoospores. Their common characteristic is the extraordinary development in size

of the single cell, which nowhere else in the vegetable kingdom attains anything like so great dimensions. Under ordinary conditions the individual is entirely unseptate, except where it is about to form reproductive bodies, whether sexual or non-sexual. The very large number of nuclei is universal in the Siphonocladaceæ; and, although their occurrence in the Siphoneæ rests chiefly on the evidence of Schmitz, there can be little doubt that this order also shares in the peculiarity, which does not lead on to cell-division, as in the groups of algæ already described; and this must clearly be regarded as indicative of a lower or more ancestral type of structure. Whether the Siphonocladaceæ and Dasycladaceæ are an earlier form leading up to the Siphoneæ, or whether they have been derived from the latter by retrogression, indicated by the suppression of the sexual organs, is uncertain, though the probability appears to be in favour of the latter hypothesis.

Order 1.—Siphoneæ (Cœloblastæ).

The thallus is in this family ordinarily unicellular, although often copiously branched, until the commencement of the formation of the organs of vegetative propagation or of sexual reproduction.

In the genus *Vaucheria* DC., which alone represents this order, the plant consists, when in a non-reproductive state, of a single elongated cell of a pale green colour, branching in various ways, sometimes as much as a foot in length, increasing by apical growth. Maupas (Comptes Rendus, *l.c.*) and Schmitz (Sitzber. Niederrhein. Gesell., 1879) state that each tube contains a large number of nuclei. The non-sexual organs of propagation are of two kinds, motionless *resting-spores* and motile *zoospores*. The former are produced simply by the abstriction of ends of particular branches, which swell up to an oval form, become cut off from the rest of the tube by a septum, contract, and finally develop a new cell-wall within the old one, thus affording an illustration of the formation of a new cell by rejuvenescence. This process takes place especially as a result of injury to the thallus. In some cases the newly-formed spore is set free simply by the absorption of the original cell-wall, and falls off with the remains of the mother-cell still attached to it, germinating after a few days; or it is thrown out with a jerk, and goes through a period of rest as a *hypnospore* before germinating. Another method is the swelling up to a considerable size of certain branches, which separate at the base, and put out at once one or more germinating tubes. The zoospores are among the most beautiful of those of any class of Algæ, being of considerable size, and entirely surrounded by a fringe of fine cilia. In their formation the extremity

of a branch is cut off by a septum as a *zoosporange*; the dark green endochrome in it gradually contracts into an ellipsoidal form, finally forcing its way out by the rupture of the cell-wall at the apex of the branch. The rotatory motion imparted to the zoospore by its fringe of cilia begins even within the mother-cell. During its escape it sometimes gets nipped in two by the pressure of the cell-wall, and each half then becomes a zoospore, one inside, the other outside, the wall of the mother-cell. The zoospores are formed in the night and escape in the morning; their spontaneous motion lasts for a period varying from less than a minute to several hours. As soon as they have come to rest they



FIG. 248.—*Vaucheria sessilis* Vauch. *A*, *B*, formation of antherids and oogones; *h*, male branch; *a*, antherid; *og*, oogone. *C*, oogone opening and ejecting drop of mucilage, *sl*. *D*, antherozoids. *E*, antherozoids entering oogone. *F*: *a*, empty antherid; *osp*, oogone with fertilised oosperm (magnified). (After Pringsheim and Goebel.)

lose their cilia, and become invested by a cell-wall of cellulose; their germination begins during the same day or the following night. The spore puts out either one or two germinating tubes, formed by its inner coat or endospore bursting through the outer coat or exospore; and the new plant usually fixes itself by a *rhizoid* or root-like organ of attachment.

The sexual reproductive organs of Vaucheria, *oogones* and *antherids*, originate as lateral protuberances on a filament, sometimes even on

the germinating tube which springs directly from a zoospore. Most of the species are monœcious, and the oogones and antherids are usually found very near together. The antherids are the terminal portions of slender branches, in some species straight, in others curved and more or less resembling horns or hooks. They contain but little chlorophyll, and the protoplasm breaks up into a large number of biciliated *antherozoids*, which escape through the ruptured apex. The two cilia are of unequal length, and point, one backwards, the other forwards. The oogones arise as thick swellings, often somewhat resembling a bird's head in shape, and are densely filled with chlorophyll; they are finally cut off from the rest of the tube by a septum. The coarsely-granular green protoplasm collects in the centre of the oogone, leaving a colourless portion at the apex which corresponds to the beak of the bird. Here the oogone opens, and a colourless drop of mucilage is sometimes expelled. When the greater part of the contents has contracted into an *oosphere*, a number of antherozoids force their way in through the open mouth of the oogone, and impregnate the oosphere by coalescing with it. The *oosperm*, resulting when the impregnated oosphere has become invested by a cell-wall of cellulose, assumes a red or brown colour, and passes through a period of rest as a *hypnosperm*. On germination several non-sexual generations are produced, ending in the formation of sexual organs. The formation of antherids and oogones begins in the evening and is completed the next morning; fertilisation being usually accomplished between 10 A.M. and 4 P.M. It is most usual for zoospores to be formed on parts of the thallus which are completely submerged in water; while the sexual organs are more commonly found when the plant grows on moist ground or on the margin of a ditch.



FIG. 249. — *Vaucheria dichotoma* Lyng. *A*, oogone; *B*, germinating spore (× 200). (After Woronin.)

Several species of Vaucheria are frequent in fresh, while some occur also in brackish or salt water; others abound on moist or shady ground, being especially common in flower-pots or on neglected gravel-paths, where they form light green tufts or thick mats. The normally unicellular thallus is liable to segmentation as the result of injury; and, even when uninjured, has a tendency to become septated by thick

gelatinous walls. In this condition it was formerly described as a distinct organism under the name *Gongrosira*. These gongrosira-cells, when isolated, develop into ordinary plants; or sometimes their protoplasm breaks up into fragments which escape from the cell-wall and move about with an amœboid motion. These invest themselves after a time with a cell-wall and remain in this condition as spherical resting-cells or *hypnospores*, finally developing into ordinary filaments. Wille has, however, shown (Bot. Centralblatt, vol. xvi., 1883, p. 162) that the organisms formerly grouped under Gongrosira are states of algæ belonging to a number of widely separated genera, such as Trentepohlia (Chroolepus), Botrydium, Stigeoclonium, &c. In some species of Vaucheria the filaments have a tendency to branch copiously at the extremity, the branches interweaving into a ball. Several species are liable to the formation of singular galls caused by the attacks of a rotifer belonging to the genus Notommata (Benkö, Bot. Centralbl., vol. xiv., 1883, p. 1).



FIG. 250.—*Botrydium granulatum* Wallr. *s*, vesicular portion; *w*, rhizoids (× 30). (After Woronin.)

Phyllosiphon (Kühn) is a truly parasitic chlorophyllous alga occurring in the south of Europe within the leaves of Arisarum vulgare, the position of which appears to be near this family. The single cell contains a large number of nuclei; the mode of reproduction is unknown (Kühn, Sitzber. Naturf. Gesell. Halle, 1878; Just, Bot. Zeit., 1882, p. 2 *et seq.*; Schmitz, id., 1882, p. 523 *et seq.*; Franke, Jahrber. Schles. Gesell., 1883, p. 195).

Endoclonium (Franke) (Cohn's Beiträge, 1883, p. 365), *Chlorochytrium* (Cohn), *Endosphæra* (Klebs), and *Phyllobium* (Klebs) (Bot. Zeit., 1881, p. 249 *et seq.*) are green parasitic algæ found within the cells of the leaves of various land and aquatic plants; and, in the case of Chlorochytrium, also on animals (Bot. Zeit., 1885, p. 605). They are stated to produce megazoospores, which germinate directly, and microzoospores or zoogametes, which germinate only after conjugation; but their exact position is altogether uncertain.

Literature.

Thuret—Ann. Sc. Nat. (Bot.), xiv., 1850, p. 214.
Pringsheim—Monber. Akad. Berlin, 1856, p. 225 (Quart. Journ. Micros. Sc., 1856, p. 63).
Schenk—Würzburg Verhandl., viii., 1858, p. 235.
Walz—Pringsheim's Jahrb. wiss. Bot., 1866, p. 127.
Solms-Laubach—Bot. Zeit., 1867, p. 361.
Woronin—Bot. Zeit., 1869, p. 137 *et seq.*
Nordstedt—Bot. Notiser, 1878, p. 176; and 1879, p. 177; and Scottish Naturalist, 1886.
Stahl—Bot. Zeit., 1879, p. 129.

Order 2.—Botrydiaceæ.

This order consists at present of only a single genus, *Botrydium* Wallr., represented by the single species B. granulatum (Wallr.), differing widely from Vaucheria in its mode of reproduction. This remarkable organism forms minute green blobs on wet clayey ground or in dried-up pools, and is attached to the soil by branching root-like rhizoids. The young plant consists of a single nearly spherical cell, branched hyaline prolongations of which constitute the *rhizoids*, while in the upper swollen part the protoplasm forms a hyaline parietal layer containing chlorophyll. From this protoplasm are ultimately produced a number of zoospores, each provided with only a single cilium, which germinate directly on the damp soil after becoming invested with a double cell-wall of cellulose. If the zoosporange is exposed to drought, the vesicular portion shrivels up and the chlorophyll is driven into the underground portion, which then divides into a number of green cells. These may germinate in three different ways:—(1) Each cell becomes an underground zoosporange, producing zoospores of the ordinary kind; or (2) each cell may develop into an ordinary vegetative plant; or (3) each cell becomes a *hypnosporange*, closely resembling the ordinary vegetative plant, with a vesicular portion above the surface, and hyaline rhizoids, but of an olive-green colour; these may retain their vitality for a whole year and then produce zoospores. The ordinary vegetative plant may also, under certain conditions, break up into a number of *resting-spores*, of a brownish red colour, which have been described as species of Protococcus. These may either give birth to zoospores of the ordinary kind or to biciliated *zoogametes* which conjugate in pairs, or sometimes in larger numbers. The resulting *zygosperm* ('isospore' of Rostafinski) soon rounds itself off and germinates.

Literature.

Janczewski et Rostafinski—Mém. Soc. Sc. Nat. Cherbourg, 1874, p. 273.
Janczewski u. Woronin—Bot. Zeit., 1877, p. 649 *et seq.*

Order 3.—Dasycladaceæ.

The Dasycladaceæ are clearly distinguished from the other orders of the class by their verticillate branches and their external sporanges. In *Acetabularia* Lmx. the thallus has the form of a small hymenomycetous fungus with a hemispherical or funnel-shaped cap or 'pileus' at the summit of a cylindrical 'stipe' or stalk. From the lower end of the stalk proceed a number of root-like branches which fix the plant to the substratum. The whole plant consists of a single ramifying cell,

Fig. 251.—*Acetabularia mediterranea* Lmx. (natural size).

the walls of which are, when mature, permeated by calcium carbonate. The cap is divided by regular radial projections into a large number of chambers of nearly equal size, which are in communication with one another above the insertion of the stalk. The upper part of the thallus perishes at the end of the season, while the lower portion is perennial. After a number of sterile thalli have been produced, a fertile thallus appears, similar in structure to the sterile ones. In the chambers of the cap of this thallus are produced the zoosporanges, a large number in each chamber, of an ellipsoidal form, and furnished at one end with a lid, which subsequently becomes detached. When mature the protoplasm of the sporange breaks up into a number of swarm-spores,



Fig. 252.—*A. mediterranea*. *I*, cap (magnified); *n*, scars of branches; *r*, rudimentary whorl of branches; *w*, ring above the cap; *v*, depressed apex (× 4). *II*, sporange with lid (× 120). *III*, the same, showing the escape of the swarm-spores (× 120). *IV*, conjugation of zoogametes. *V*, *VI*, lower part of stem, showing rhizoids, *b*. *VII*, plant germinating from a zygosperm. *VIII*, origin of branches at summit of the stem (× 120). *IX*, at a later stage (× 90). (After de Bary and Strasburger.)

which escape into the surrounding water by the removal of the lid. It would appear from de Bary and Strasburger's observations that some of these swarm-spores germinate directly, while others conjugate to form a zygosperm; but that conjugation never takes place between zoogametes from the same sporange. One species of Acetabularia is a native of the Mediterranean; the rest are tropical. In *Dasycladus* Ag. the spherical zoosporanges stand singly at the apex of verticillate branches, and are surrounded by branchlets of the second order. The biciliated zoospores or zoogametes are of one kind only, and are flattened and heart-shaped. Those from the same plant show no disposition to conjugate; but as soon as those from different plants are brought together, true zygosperms are formed. In *Neomeris* Lmx. the surface consists of a large number of usually hexagonal facets, and is covered with deciduous hairs. The



FIG. 253.—*Dasycladus clavæformis* Ag. *a*, natural size; *b*, piece of branch of a whorl with a zoosporange (× 50). (After Hauck and Derbès and Solier)



FIG. 254.—*Neomeris Kelleri* Cram.; transverse section through tube. *a*, insertion of the branches; *b*, *b*, primary branches; *c*, central secondary branch or zoosporange; *d*, elongated lateral secondary branches (× 40). (After Cramer.)

very large axial cell is always simple; the lateral branches again divide into three branchlets, of which the central one is ovoid and fertile, the two lateral ones greatly elongated and sterile. The fertile branch is a

zoosporange, no conjugation of swarm-spores having been observed. Klein (Bot. Zeit., 1880, p. 782) has detected fine proteinaceous crystalloids in Dasycladus (Ag.) and other allied forms. To this family belong also Polyphysa (Lmx.), Cymopolia (Lmx.), and some other genera.

FIG. 255.—*Polyphysa peniculus* R. Br., with whorl of zoosporanges, *sp* (magnified). (After Agardh.)



FIG. 256.—*Caulerpa prolifera* Lmx. (natural size). (After Reinke.)

ORDER 4.—SIPHONOCLADACEÆ.

This order comprises a number of very remarkable green algæ, mostly inhabitants of warm and shallow seas, characterised by the thallus consisting of a single cell which is often of very great size and much branched, differentiated into a root-like and stem-like portion. The undivided cell very commonly contains a large number of nuclei; and the cell-wall is

often strongly encrusted with calcium carbonate. In several of the genera of Siphonocladaceæ, but little is at present known as to the mode of reproduction ; and until this has been more fully ascertained, their true affinities are uncertain : and it is possible that the order, as at present constituted, includes forms which are not nearly allied to one another. In none of the genera is a fertilisation of oogones by antherozoids known, similar to that of Vaucheria. The ordinary mode of reproduction appears to be by zoospores which germinate directly without conjugation. Other modes of non-sexual propagation occur in some of the genera, by 'propagules' or by 'prolification.' The following are some of the more remarkable forms included in the group.

The Caulerpeæ include the single genus *Caulerpa* Lmx., characterised by its greatly-branched thallus of remarkably leaf-like appearance. Very little is known about the mode of reproduction. The ordinary process of propagation appears to be by 'prolification' from all parts, the so-called 'roots,' 'stem,' and 'leaves.' Within the tube are solid branched layers of cellulose stretching from wall to wall, and forming a closed net. It often covers enormous tracts of the shore between high and low water marks with a green coating.

Fig. 257.—*Valonia macrophysa* Ktz. (natural size). (After Hauck.)

The Valoniaceæ (Valonia, Gin., Siphonocladus, Schr., Struvea, Sond., Anadyomene, Lmx., &c.) are an ill-defined family, in which the cell is frequently swollen up into a bladder-like structure ; the mode of propagation is apparently by non-sexual zoospores. To these are closely allied the Udoteaceæ (Avrainvillea, Dcne., Penicillus, Ktz., Udotea, Lmx., Halimeda, Lmx., &c.). *Halimeda* has a remarkable Opuntia-like appearance, from the single cell consisting of a large number of pear-shaped branches, each connected with the one below it by a narrow base, and the whole encrusted by calcium carbonate.

In the Bryopsideæ (Bryopsis, Lmx., Derbesia, Sol.) and Spongodieæ (Codium, Stackh.) the thallus is not encrusted with calcium carbonate. *Codium* forms a spongy spherical or cylindrical floating mass, of considerable size, consisting of branched tubes. It is apparently propagated by zoospores. In *Bryopsis* Lmx. the thallus has a branched root-like colourless portion, and an erect cylindrical stem, the upper half of which branches into pinnate leaf-like ramifications with limited apical

growth ; in these branches zoospores are formed ; conjugation between them has not actually been observed. External organs of reproduction known as 'conceptacles' have also been described on several species of Bryopsis, which have been conjectured to be true sexual organs, whether male or female, but their true structure and functions are altogether obscure. If, as is probable, some of the swarm-spores of Codium

FIG. 258.—*Halimeda Tuna* Lmx. *a*, natural size ; *b*, portion with zoosporanges (× 50). (After Derbès et Solier.)

and Bryopsis are in reality conjugating zoogametes, this would bring these two families into close relationship to the Botrydiaceæ ; and this is probably the nearest affinity of the latter group. Spongocladia (Aresch.), described by Zanardini as belonging to the Siphoneæ, appears to be a genus more nearly allied to Cladophora, in which the filaments are remarkably infested with sponge-spicules.

LITERATURE.

(*Dasycladaceæ and Siphonocladaceæ.*)

Nägeli—In Zeitschr. wiss. Bot., 1844.
Derbès & Solier—Ann. Sc. Nat. (Bot.), 1850, p. 240.

Braun—Verjüngung in der Natur, 1851, p. 136 (Ray Soc., Bot. and Phys. Mem., 1853).
Woronin—Ann. Sc. Nat. (Bot.), xvi., 1862, p. 200.
Pringsheim—Monber. Berlin Akad., 1871, p. 240.
Arcangeli—Nuov. Giorn. Bot. Ital., 1874, p. 174.
De Bary u. Strasburger—(Acetabularia) Bot. Zeit., p. 713 *et seq.*
Cornu—Compt. Rend., lxxxix., 1879, p. 1049.
Schmitz—Sitzber. Niederrhein. Gesell., 1879 and 1881.
Berthold—Bot. Zeit., 1880, p. 648; and Mittheil. Zool. Stat. Neapel, 1880, p. 72.
Murray—(Rhipilia Ktz. = Avrainvillea Dcne.) Trans. Linn. Soc., ii., 1886, p. 225
Wakker—(Caulerpa, Versl. Akad. Weten. Amsterdam, 1886, p. 251; and 1887, p. 251.
Agardh—Till Algernes Systematik, Siphoneæ, 1887.
Cramer—Ueber die verticillirten Siphoneen (Neomeris und Cymopolia), 1887.
Noll—Bot. Zeit., 1887, p. 473.
Murray and Boodle—(Struvea) Ann. of Bot., ii., 1888.

Class XVII.—Cœnobieæ.

In this class are included a small number of minute (mostly microscopic) fresh-water organisms, characterised by the cells being associated together into a *cœnobe*, i.e. into a colony of more or less equivalent cells resulting from the division of a common mother-cell. As this division always takes the form of repeated bipartition, the number of cells constituting a colony is necessarily, when perfect, a power of 2, viz. 4, 8, 16, 32, 64, &c. The cells constituting the cœnobe are more or less imbedded in a gelatinous envelope, which is sometimes enclosed in a membrane common to the whole colony; in the higher forms the cells, or some of them, are ciliated, the cilia protruding through the enveloping membrane, and the colony moves about in the water with very great activity; the lower forms are not ciliated, but the colony is nevertheless endowed with a very considerable power of motion. The cœnobe is always of an exceedingly beautiful regular form, spherical, or less often discoid or cubical, or, in the Hydrodictyeæ, in the form of a net. The five orders of which it is composed, the Sorastreæ, Pandorineæ, Pediastreæ, Hydrodictyeæ, and Volvocineæ, form a series of ascending development. Very little is known about the reproduction of the first; the Pandorineæ, Pediastreæ, and Hydrodictyeæ multiply by the conjugation of zoogametes; while in the Volvocineæ, which represent the highest type attained by organisms of the cœnobe type, the mode of sexual reproduction is much more complicated, the male and female reproductive cells being separately formed in distinct antherids and oogones.

Order 1.—Volvocineæ.

The well-known *Volvox globator* L. may be taken as a type of this interesting family. This organism is not uncommon in clear pools, and is visible to the naked eye as a minute pale-green globule, about $\frac{1}{30}$ inch in diameter, rolling through the water, the motion being due to numerous colourless cilia projecting from its surface. It is one of the most beautiful objects that can be observed under the microscope.

Under a sufficiently high power of the microscope, Volvox is seen to be a membranous transparent hollow sac studded with green points; in the interior are darker green globules, the original number of which is apparently always eight. The green peripheral corpuscles or swarm-cells are each provided with a pair of vibratile cilia, which protrude through the enveloping sac; they vary in form, but are usually somewhat pear-shaped, and contain a starch-granule, a reddish-brown pigment-spot, and one or two contractile vacuoles, the cilia being borne at the narrow anterior hyaline end. The internal green globules are young individuals formed within the parent, which thus constitutes a colony or *cœnobe*; but all the cells which make up the colony are not equivalent as respects their reproductive power. The larger number of the cells are sterile or purely vegetative, while a much smaller number, developed at particular spots in the colony, are generative, these again being differentiated into non-sexual propagative and male and female reproductive cells. The sterile cells are the peripheral or *swarm-cells*, 2–3 mm. in diameter, which precisely resemble in structure Chlamydococcus, the motile stage of Pleurococcus, or the zoospores of many algæ. The gelatinous membrane which envelopes each of these swarm-cells is pierced by a number of canals, which lie nearly in one plane, and which are filled by green or colourless extensions of the protoplasmic interior. Since the canals of adjoining swarm-cells correspond in position, they appear as if they were all connected together by a network of fine reticulations. The membrane is also perforated by two pores, through which the vibratile cilia protrude into the surrounding water. These bodies present the unusual phenomenon of cells endowed with spontaneous power of motion, which have, nevertheless, as far as is known, no reproductive function, and are therefore not properly called zoospores. The non-sexual propagative cells, *zoospores* or parthenospores, are similar in structure to the sterile swarm-cells, but from two to three times their diameter. Very early in the development of the young colony the contents of the mother-cells of the zoospores begin to divide by repeated bipartition, all in the same colony being usually at the same stage of development at one time; the daughter-colonies,

or *zoosporanges* thus formed have the form and appearance of the parent-colony, each segment possessing a single chlorophyll-body, which contains starch. Ultimately, while still within the parent-colony, vibratile cilia are developed in its peripheral segments, which cause it to rotate, clothed at first by a transparent mucilaginous envelope, which it at length breaks through. The number of colonies of zoospores thus produced within the parent-colony is normally eight, resulting apparently from the eight cells into which the parent-colony breaks up on its third segmentation. The young colonies complete their growth in a few days, attaining a diameter of from 100 to 150 mm., and have by this time absorbed the greater part of the chlorophyll and starch of the parent-colony, from which they finally escape.

Volvox may multiply by this non-sexual mode of propagation for several successive generations, and these are succeeded by a sexual mode of reproduction. The male and female reproductive cells are formed either in the same or in different colonies; or, according to other observers, the so-called 'dioecious' species are in reality proterandrous, producing antherids at an earlier, oogones at a later stage. While the non-sexual propagation by zoospores goes on through the whole year, the sexual cells appear to be produced only in the autumn. The *oogones* are at first quite indistinguishable from the non-sexual cells except in size, but are much more numerous, and soon manifest a distinction from the fact that they do not divide. On their first appearance they are about three times the size of the sterile cells; their protoplasm increases rapidly, and becomes of a dark-green colour, from copious production of chlorophyll. They have at first a frothy appearance from the formation of vacuoles, but afterwards appear to be densely filled with the dark-green endochrome. They soon become flask-shaped, the narrow end touching the periphery of the colony, and the larger end hanging free inside; but, when ready for impregnation, round themselves off into a spherical form, their contents being an *oosphere* enveloped by a gelatinous membrane. The *antherids* present at first sight a still closer resemblance to zoospore-colonies at an early stage, but are lighter in colour from containing a smaller quantity of chlorophyll. Their contents soon begin to divide, but in two directions only, the young colony thus developing into a plate instead of a sphere of segments enveloped in a gelatinous coating. The colony ultimately resolves itself into a bundle of *antherozoids*, naked fusiform masses of protoplasm, each consisting of a thicker but elongated body, in which the chlorophyll has been transformed into a yellow-brown pigment, and a slender colourless beak, with a pigment-spot at its base, where also are attached on one side two very long cilia. About the time when the oogones, with their



FIG. 259.—*Volvox globator* L. *A*, colony, sexual generation; *a*, antherids; *b*, oogones; *c*, oospheres. *B*, ciliated peripheral cells. *D*, antherid. *E*, *F*, antherozoids. *G*, oogone containing mature oosphere and antherozoids. *H*, mature oosperm. *C*, mode of division of parent-cell of a zoosporange (all highly magnified). (After Cohn.)

oospheres are mature, the movement of the cilia begins to set the entire antherid in motion : but it shortly breaks up, and the separate antherozoids are seen in rapid motion within their gelatinous envelope, which they ultimately break through, and then move about rapidly in all directions within the cavity of the parent-colony. They assemble in large numbers round the oogones, and some of them finally penetrate through the gelatinous membranes of these organs, and coalesce with the oospheres. The number of sexual reproductive colonies within a parent-colony varies greatly : Cohn has observed five or six male and about forty female. The impregnated oosphere now becomes an *oosperm* by the development of a cell-wall of cellulose, which is at first single and smooth, but becomes subsequently differentiated into three distinct layers, of which the two inner ones are smooth, while the outermost becomes ultimately covered with conical or warty elevations, giving it on section a stellate appearance. The chlorophyll of the oosperms gradually disappears, and is replaced by an orange-red pigment dissolved in oil, giving a red tinge to the entire organism as seen by the naked eye. Soon after the maturity of the oosperms, the parent-colony breaks up, and the peripheral swarm-cells escape from the combination and swim about freely in the water ; their further history has not been traced. The orange-yellow oosperms sink to the bottom, and there hibernate as *hypnosperms*. Their contents are said to break up into eight Volvox-colonies, which ultimately escape by the rupture of the outer and absorption of the inner coats of the oosperm, and swarm about in the water.

In *Eudorina* Ehrb. the cœnobe is a hollow ellipsoid body, consisting of usually 16 or 32 cells enclosed in a gelatinous envelope, each provided with two long cilia which protrude through canals in the envelope, and a red pigment-spot. Daughter-colonies are formed non-sexually within the parent-colony by repeated bipartition of its cells, the cells of the daughter-colony being arranged in a disc. The male and female reproductive bodies are formed in special daughter-colonies which may be termed antherids and oogones respectively ; the oospheres contained in the oogones are each provided with two cilia, and are therefore true *zoospheres* : the antherozoids, which closely resemble those of Volvox, swarm round the oogones until their cilia become entangled in those of the zoospheres ; they then force their way into the gelatinous envelope of the oospheres, and finally coalesce with them.

Volvox and Eudorina are regarded by Ehrenberg and Stein as belonging to the Flagellate Infusoria.

Under the name of Scyamina, Van Tieghem (Bull. Soc. Bot. France, 1880, p. 200) describes a singular blackish organism found on the surface of ponds, which he regards as a genus of Volvocineæ destitute of chlorophyll.

Literature.

Ehrenberg—Die Infusionsthierchen, 1838.
Busk—Trans. Micros. Soc., 1853, p. 31.
Cohn—Jahrber. Schles. Gesell., 1856, p. 237; and Beiträge, i., 1875, Heft 3, p. 93 (Pop. Sc. Rev., 1878, p. 225).
Carter—(Eudorina) Ann. Nat. Hist., 1858, p. 237; and 1859, p. 1.
Braun—Bot. Zeit., 1875, p. 189.
Stein—Der Organismus der Infusionsthiere, part iii., 1878.
Henneguy—Journ. d. Micrographie, 1878, p. 485.
Maupas—Compt. Rend., lxxxix., 1879, p. 129.
Kirchner in Cohn's Beiträge, iii., 1879.

Order 2.—Hydrodictyeæ.

The relationship of the Hydrodictyeæ to the other families of Cœnobieæ is somewhat obscure. They differ from them in the form of the cœnobe, which, instead of being minute and globular, ellipsoidal or tabular, is of considerable size, and presents the appearance of a net. The only known mode of reproduction is by the conjugation of zoogametes.

As here constituted the order is limited to the single genus *Hydrodictyon* Roth, one species of which is moderately common in ponds and ditches, and is known under the name of 'water-net.' When the plant is mature, the cœnobe consists of a sac-like net several centimetres in length, composed of a great number of cylindrical cells united at their ends so as to form a 4- or 6-cornered mesh. The ordinary mode of propagation consists in the protoplasm of one of the cells breaking up into from 7,000 to 20,000 *megazoospores*, each furnished with four cilia, which move about with a trembling motion within the zoosporange, come to rest in the course of half an hour, and then arrange themselves in such a way that, by their elongation, they again form a net of the original kind, which is set free by the absorption of the wall of the mother-cell, and attains, in the course of three or four weeks, the size of the mother-colony. In other cells of the mature net the protoplasm breaks up into from 30,000 to 40,000 smaller swarm-cells or *zoogametes*, furnished with only two cilia, which at once leave the *gametange* and swarm about for some hours. Conjugation between these has been observed to take place even within the gametange, more than two sometimes uniting together. The resulting *zygosperm* retains its green colour, and invests itself with a firm cell-wall. After remaining for several months at rest as a *hypnosperm*, it begins to grow slowly, and, when it has attained a considerable size, its contents break up into two or four megazoospores, which come to rest after a few minutes, and assume a peculiar angular form when they have attained a considerable size, putting out horn-like appendages. In each of these *polyhedra* the green parietal

protoplasm again breaks up into zoospores, which swarm about for twenty or forty minutes within a sac which protrudes out of the protoplasm. When they come to rest, they arrange themselves within the sac into a small net consisting of from 200 to 300 cells, which gradually grows to



FIG. 260.—*Hydrodictyon utriculatum* Roth. *A*, net (natural size). *B*, mesh (× 10). *C*, megazoosporange (× 300). *D*, megazoospores (× 600). *E*, gametange with zoogametes (× 300). (After Cohn.)

one of the ordinary size. In some of the polyhedra smaller and more numerous microzoospores are formed; but they also appear to unite again into a net without displaying sexual functions.

LITERATURE.

Nägeli—Gattungen einzelliger Algen, 1849, p. 92.

Braun—Verjüngung in der Natur, 1851 (Ray Soc., Bot. and Phys. Mem., 1853); and Algæ unicellulares, 1855.

Pringsheim—Monber. Berl. Akad., 1860, p. 775 (Quart. Journ. Micr. Sc., 1862, pp. 54 and 104).

Order 3.—Pediastreæ.

The Pediastreæ are most nearly allied to the Hydrodictyeæ in their mode of reproduction. Of the typical genus *Pediastrum* Mey., several species are very common in fresh water, whether stagnant or running, attached in the form of minute (usually microscopic) discs to other algæ or water-plants, or swimming free. Each disc is of a regular symmetrical



Fig. 261.—*Pediastrum integrum* Näg. *A*, younger, *B*, older cœnobe (× 300). *C*, portion of older cœnobe showing resting-cell, *r* (× 600). (From nature.)

form, usually elliptical, and consists most often of 8, 16, or 32 cells, or some larger number which is probably always, when perfect, a power of two. The cœnobe is invested in a very thin gelatinous envelope, and the peripheral cells have commonly horn-like or crescent-shaped appendages. Pediastrum is multiplied either by non-sexual propagation or by sexual reproduction. In the former case one of the ordinary vegetative cells becomes a zoosporange, its protoplasm breaking up into a number of nearly globular megazoospores each furnished with two very fine and inconspicuous cilia, which, after swarming about for a time, lose their cilia and arrange themselves in the form of a plate, which then escapes from the zoosporange invested in mucilage, and develops into a new Pediastrum-disc. Others of the cells become gametanges, the contents dividing in the same way into zoogametes of an ovoid or pear-shaped form, which conjugate after escaping separately from the mother-cell, but apparently only with those from other gametanges. The peripheral cells of the cœnobe appear to have a tendency to develop into *resting-spores*.



Fig. 262.—*A*, polyhedra of *Pediastrum* (× 550). *B*, formation of *Pediastrum*-disc within polyhedra (× 550). (After Askenasy.)

Askenasy has observed the development of the Pediastrum-cœnobe by another method from the *polyhedra* form, previously regarded as a distinct genus under the name Polyedrium (Näg.). In this form each individual consists of a minute flat angular cell often provided with spines or hook-like processes. From this a Pediastrum-disc is developed in precisely the same way as from a Pediastrum-cell. The cell-contents break up into a number of megazoospores, which escape in the form of a plate after swarming about for a time; then, losing their cilia, and placing themselves in a plane side by side, develop into an ordinary Pediastrum. Reinsch, on the other hand (Notarisia, 1888, p. 493), regards Polyedrium as the type of a separate family of Palmellaceæ.

Literature.

Nägeli—Gattungen einzelliger Algen, 1849.
Braun Verjüngung in der Natur, 1851 (Ray Soc., Bot. and Phys. Mem., 1853); and Algæ unicellulares, 1855.
Lagerheim—Bot. Centralbl., xii., 1882, p. 33.
Askenasy—Ber. Deutsch. Bot. Gesell., 1888, p. 127.

Order 4.—Pandorineæ.

In the higher genera of this order, Pandorina (Ehrb.), Gonium (Müll.), and Stephanosphæra (Cohn), the individual is a spherical or tabular cœnobe, the cells of which are united together by a gelatinous matrix with a definite bounding wall. With them are associated also some unicellular organisms, Chlamydococcus (A. Br.) and Chlamydomonas (Ehrb.), which may possibly be connected with them by a process of degeneration. Whether isolated or associated, each cell possesses a pair of whip-like vibratile cilia attached to the anterior pointed end, by means of which it is rapidly propelled through the water; in the case of the social genera these cilia project through the common gelatinous envelope of the colony. Multiplication takes place either non-sexually by simple subdivision of the cells of a colony, or sexually by the union of two (or occasionally more than two) *zoogametes* into a resting *zygosperm*. A characteristic feature of the family is the formation of a colony of cells within each cell in the mother-colony. The organisms here included were described by Ehrenberg as constituting a family of Infusorial Animalcules. They live, associated with larger algæ, in fresh water, running or stagnant, often in such quantities as to impart to it a green colour. The family closely approaches Volvocineæ through Eudorina.

Of the unicellular Pandorineæ *Chlamydomonas* Ehrb. may be taken as a type. In the form in which it is known under this name, it

consists simply of a single motile primordial cell, in other words, of a zoospore or swarm-cell. These are megazoospores, half as long again as broad, each with two contractile vacuoles, a lateral red pigment-spot, and two long cilia; in the posterior half is a nucleus. After the conclusion of the period of swarming, these zoospores become invested with cellulose, and, after a long period of rest as *hypnospores*, multiply non-sexually by division into four—less often into two. According to Rostafinski sexual reproduction takes place by one of these megazoospores dividing, by successive bipartitions, into eight daughter-cells, which are then microzoospores—or, more correctly, *zoogametes*. These have a longish almost elliptical form, and a light green colour, one nucleus, a red pigment-spot, and four cilia. They are distinguished from the true zoospores, not only by their smaller size, but also by the large colourless extremity in place of the two contractile vacuoles. They swarm out, and soon begin to conjugate in pairs, coming into contact by their colourless extremities, and coalescing into a single cell, the ends which bear the cilia at the same time rounding off and approximating. This body has now eight cilia and two lateral pigment-spots; ultimately the eight cilia disappear, and a zygosperm is formed, which multiplies by simple division without swarming. Dangeard finds in Chlamydomonas pulvisculus (Müll.) a differentiation of male and female gametes, the latter being considerably larger than the former. In C. Morieri (Dang.) he describes a peculiar mode of conjugation of zoogametes, which he compares to that of Spirogyra.

Chlamydococcus A. Br. presents a similar life-history; and, according to some observers, the organisms known as Pleurococcus, Glœocystis, and others usually included under Protococcaceæ, are identical with the resting conditions of Chlamydomonas; and, under suitable conditions, can be made to produce biciliated zoospores with two contractile vacuoles and a nucleus.

Under the Volvocineæ, and near to Chlamydomonas, Dangeard (Ann. Sc. Nat., vii., 1888, p. 105) and Stein include also Chlorogonium (Ehrb.).

Among the most interesting of the social Pandorineæ is *Pandorina* Ehrb.. Each family or cœnobe consists of sixteen cells closely crowded together, and surrounded by a thin gelatinous envelope through which the cilia protrude. Non-sexual multiplication is preceded, after the colony has come to rest, by the absorption of the cilia in the sixteen cells, each of which then breaks up into sixteen smaller cells; and these sixteen daughter-families are set free by the absorption of the gelatinous envelope of the parent-colony; each becomes itself invested by a gelatinous envelope, and grows to the size of the original parent-colony,

having in the meantime developed two cilia from each of its cells. Sexual reproduction takes place in the following way. Sixteen daughter-families are first of all formed in the same manner, but the gelatinous envelopes of the young colonies deliquesce, and the separate 256 (= 16 × 16) swarm-cells are set free as zoogametes. These vary in size, but are always rounded and green at the posterior end, pointed hyaline and with a red pigment-spot in front, where they bear two cilia. Among the crowd of these swarm-cells—now swimming about freely—some, irrespective of their relative size, approach one another in pairs, their pointed anterior apices coming into contact, and they finally coalesce into a body which has at first somewhat the shape of an hour-glass, but gradually contracts into a sphere, in which the two pigment-spots and the four cilia are still to be seen for a time, but soon disappear. This whole process occupies about five minutes. The resulting zygosperm is then a spherical cell enclosed in a cell-wall, which remains at rest for some time as a hypnosperm, its green colour becoming changed to brick-red. If the dried-up hypnosperms are placed in water, they begin to germinate after about twenty-four hours. The outer layer of the cell-wall is ruptured, an inner layer becomes gelatinous and swells up, and the protoplasmic contents escape in the form of one, two, or three large zoospores. Each of these, after a short period of swarming, loses its cilia, surrounds itself with a gelatinous envelope, and breaks up, by successive bipartitions, into sixteen portions, which develop cilia, and form themselves into a new cœnobe.



FIG. 263.—*Pandorina morum* Fhrb. *a*, swarming cœnobe; *b*, *c*, swarm-cells; *d*, *e*, conjugation of zoogametes; *f*, zygosperm (× 500). (After Pringsheim.)

A still more remarkable succession of phenomena is exhibited by *Stephanosphæra* Cohn, a rare organism occurring occasionally in the rain-water which collects in the hollows of large stones in mountainous countries. In addition to the process of vegetative or non-sexual propagation, the cells belonging to a family, each of which possesses a red pigment-spot, divide repeatedly into zoogametes, which ultimately become free, and coalesce into resting zygosperms or hypnosperms. Motionless balls, which are probably the result of this conjugation, accumulate at the bottom of the water, and assume a red colour. After

these resting-cells have lain for some time dry, and then again been flooded, the contents break up into four or eight zoospores, which invest themselves with a cell-wall, and, in the course of a single day, divide, by successive bipartitions, into an eight-celled cœnobe, which again, during the next night, gives birth to eight motile families. The enclosed primordial cells of Stephanosphæra are of a bright green colour, fusiform, and are attached to an equator of the investing membrane at both ends by branched strings of protoplasm. The whole family rotates on an axis at right angles to the plane which passes through them all.

In *Gonium* Müll. the cœnobe is a tabular aggregation of cells moving rapidly through the water by the aid of vibratile cilia. Its life-history has not been fully followed out, and very little is known of its mode of reproduction.

LITERATURE.

Henfrey—Trans. Micros. Soc., 1856, p. 49.
Cohn & Wichura—(Stephanosphæra) Nova Acta Acad. Nat. Cur., xxvi., Suppl. 1857, p. 1 (Quart. Journ. Micros. Sc., 1858, p. 131).
Archer—Quart. Journ. Micros. Sc., 1865, pp. 166, 185.
Cienkowski—Bot. Zeit., 1865, p. 21.
Pringsheim—Monber. Berlin Akad., 1869, p. 721; and Bot. Zeit., 1870, p. 265.
Volten—Bot. Zeit., 1871, p. 383.
Rostafinski—Bot. Zeit., p. 785; and Mém. Soc. Sc. Nat. Cherbourg, 1875.
Goroshankin—Nachr. Gesell. Naturf. Moskau, 1875.
Hieronymus—Oesterr. Bot. Zeitschr., 1874, p. 313 *et seq.*
Reinhard—Arbeit. Naturf. Gesell. Charkoff, 1876.
Bréal—Bull. Soc. Bot. France, 1886, p. 238.
Dangeard—(Chlamydomonas) Ann. Sc. Nat., vii., 1888, p. 105.

ORDER 5.—SORASTREÆ.

In this order are included a few genera distinguished by the cœnobe being unciliated. In *Sorastrum* Ktz. the colony consists of a more or

FIG. 264.—*Sorastrum spinulosum* Näg. (× 400). (After Cooke.)



FIG. 265.—*Cœlastrum cubicum* Näg. (× 600). (From nature.)

less spherical aggregation of closely-packed, horned or bifid, somewhat

wedge-shaped cells ; in *Selenastrum* Reinsch the cells are crescent-shaped ; in *Cœlastrum* Näg. the cœnobe is spherical or cubical, composed of a single layer of cells, and hollow in the centre. The species are found occasionally in bog-pools ; although unciliated, the cœnobe swims freely in the water with a kind of rolling motion. Very little is known with regard to the mode of reproduction ; no formation of zoospores has been detected. Lagerheim describes the mode of formation of a cœnobe of Selenastrum by cell-division within the mother-cœnobe. The affinities of this very beautiful family are clearly with Botryococcus among the Protococcaceæ, and upwards with the Pandorineæ. *Scenedesmus* Mey. is very probably a primordial or a retrogressive member of this family.

LITERATURE.

Nägeli—Gattungen einzelliger Algen, 1849, p. 97.
Reinsch—Algen-Flora Mittel-Franken, 1867.
Lagerheim—Bot. Centralbl., xii., 1882, p. 33.
Bennett—Journ. Micr. Soc., 1887, p. 13.

FOSSIL ALGÆ.

All that we know of the relationships between the animal and vegetable kingdoms leads us to the conclusion that the appearance of animal life, both in fresh and in salt water, must have been preceded by that of aquatic vegetation ; and it is almost certain that these primeval vegetable organisms must have had a structure and mode of life which would classify them under the head of Algæ or of Schizophyceæ. But, since even the largest of these organisms would probably consist entirely of cellular tissue, it is not to be anticipated that their remains could be handed down to us in the fossil state except in those cases where the cell-walls were either silicified or impregnated or coated with lime.

As far back as the Laurentian period, beds of graphite occur which must undoubtedly have been the result of the decomposition of vegetable matter, but all traces of the structure of the organisms from which it has been derived are lost. The structures from the Russian coal-fields described by Reinsch as the remains of algæ allied to the Scytonemaceæ, are either the spores of Vascular Cryptogams, or, in many cases, are inorganic crystallisations. Even the very earliest argillaceous deposits, whether from fresh or from salt water, display long rounded trailing impressions, which are believed by some writers to be the remains of algæ ; but it is exceedingly difficult to distinguish between these and

the trails of aquatic animals, or even ripple-marks. Carruthers is disposed to regard the structure which he has called Nematophycus, from the Lower Erian or Upper Silurian beds of Canada, as the earliest algoid remains or impressions which have come down to us. The true nature of the tracings in the still earlier Laurentian rocks known as Eozoon canadense is still a subject of controversy with palæontologists and petrologists.

Coming down to more recent periods, the organisms described as Chondrites and Confervites, found in the Cretaceous beds, were possibly algæ allied to the Confervaceæ, and are thought by some to have had a large effect in the precipitation of chalk from sea water, from the great quantity of carbonic acid which they removed. From nodules in the Pleistocene of Canada, Dawson obtains remains which he places under genera allied to Ulva or Fucus, and possibly also to Laminaria. Various other remains, sometimes of gigantic size, found in strata extending over a very wide geological range, are referred by Saporta and others to algæ allied to the Laminariaceæ and Fucaceæ : but the evidence is generally too imperfect to justify a settled conclusion on the subject.

Of the calcareous remains of algæ which have come down to us, the most remarkable are those described by Munier-Chalmas, including more than fifty genera, which he refers to the Siphonocladaceæ, mostly found in the Triassic, Jurassic, Cretaceous, and Tertiary strata. The calcareous skeleton contains hollow chambers and canals where the rays and the organs of reproduction were seated in life, rendering them very liable to be mistaken for the remains of Foraminifera, under which class of animals many of them have been placed by palæontologists. Among these is the genus Ovulites from the Eocene.

Literature.

Balfour—Introduction to Palæontological Botany, 1872.
Carruthers—(Nematophycus) Monthly Microscopical Journal, 1872, p. 208.
Munier-Chalmas—Comptes Rendus, 1877, p. 814 ; Bull. Soc. Géol. France, 1881, p. 661.
Saporta—Les organismes problématiques des anciennes mers, 1884.
Reinsch—Micro-palæophytologia, 1884.
Nathorst—Nouvelles observations sur des traces d'animaux et autres phénomènes d'origine purement mécanique décrits comme Algues fossiles, 1886.
Delgado—Etudes sur les Bilobites, 1886, 1888.
Dawson—Geological History of Plants, 1888.
(See also under Fossil Vascular Cryptogams.)

FIFTH SUBDIVISION.

FUNGI.

Until recent years Fungi were looked upon as a great group embracing all Thallophytes which do not vegetate by means of intrinsic chlorophyll. With the advance of research and the widening of knowledge, a new classification of Thallophytes was established, mainly by Sachs, which was based on the characters of the sexual organs, and under it groups were constituted composed of Algæ and Fungi alike, in recognition of the principle that a mere physiological attribute, such as the presence or the absence of chlorophyll, should be no bar to the bringing together of organisms associated with each other by morphological characters. With the analogy before their eyes of the relationship of flowering parasites devoid of chlorophyll with green Flowering Plants, morphologists readily accepted the proposed grouping, and until a few years ago it was generally adopted. It then became apparent that this step, though in the right direction, exceeded due bounds; and, with the first publication by de Bary of the classification of Fungi used in this treatise, a new movement, which cannot, however, be justly called a reaction, took place. This classification recognises a main group of Fungi branching off from the Algæ (Chlorophyceæ), and definitely marked by morphological relationship. This main group differs from the old group of Fungi not only in its internal disposition, but in the exclusion from it of *Mycetozoa* and *Bacteria*, though it resembles the old group in its latter days in the inclusion of lichen-forming fungi. As clearly characterising and delimiting the group, the words of de Bary ('Lectures on Bacteria,' p. 2) may here be quoted:—

'The term Fungi denotes a group of lower plant-forms, distinguished by definite characteristics of structure and development, and recognised at once when we see a mushroom or a mould. The members of this group are all, as a matter of fact, devoid of chlorophyll, but they might contain chlorophyll and yet belong to this group, just as a bird may have no apparatus for flight, and yet be allowed to be a bird. To these Fungi, as defined by natural history, and not by physiological characters

only, Bacteria are as little related in structure and development as bats are to birds ; the relationship is even less, because there are a few, though only a few, true Bacteria which contain chlorophyll and decompose carbon dioxide, and which are therefore not Fungi in the physiological sense.'

Histological Characteristics.

The *thallus* of fungi consists of one or more cylindrical *hyphæ* branching in monopodial fashion, rarely dichotomously, and increasing by apical growth. In numerous instances the hypha remains, for a period at least, unsegmented, but most frequently transverse septa are formed either in the apical cell or in segment-cells of the first order. The thallus of the simple filamentous fungi consists of one hypha with its ramifications. The so-called *compound thallus* of larger fungi is formed of hyphæ of which the branches meet and remain in close contact. While all fungi have the filamentous habit to begin with,



Fig. 266.—Clamp-connections: mycele of *Hypochnus centrifugus* Tul. (× 390). (After de Bary.)

Fig. 267. — Pseudo-parenchyme breaking up at end into separate hyphæ : *Nectria cinnabarina* Fr. (highly magnified). (After Reinke)

some remain so throughout their whole development, and others are characterised by both habits in different periods of their life-history. It should also be mentioned that the whole thallus of a fungus may be reduced to a single globular cell. *Clamp-connections* take place, especially in the Basidiomycetes, between adjoining segment-cells of the same hypha by means of a protuberance emitted from the one nearer the growing-point. This protuberance issues immediately above the transverse wall, and effects a junction with the cell beneath. By this means open communication is maintained between the cells for a time, but it is

eventually closed by the formation of a transverse wall in the protuberance. Cross-links, loops, &c., are frequently formed between branches of the same hypha, or between originally distinct hyphæ, by the resorption of the membranes at the point of contact. The coherence of the compound thallus is most commonly effected by the more or less dense interweaving of the hyphæ, and by the cementing of them together in many cases by an intercellular substance. The union of the component hyphæ may proceed so far, and the conditions of pressure, &c., be so favourable, that a tissue is formed resembling the parenchyme of higher plants. Being thus different from it in mode of formation, the tissue in question is termed *pseudo-parenchyme*.



FIG. 268.—*Saccharomyces cerevisiæ* Meyen. *a*, single cell of beer-yeast; *b*, *c*, stages of sprouting; *d*, colony of sprout-cells; *e*, cell with four ascospores; *f*, one with two; *g*, group of ascospores with one sprouting; *h*, further development of a similar group (*h* × 750, the others much more). (*e h* after Reess.)

The form of hypha described is not universal among fungi. *Sprouting*, as it is called, is characteristic of the thallus of the yeast-fungus Saccharomyces, and it occurs in certain stages of other fungi, such as Mucor. This process consists in the production of ovoid or globular cells, by means of excrescences which remain connected by a narrow base with the parent cell, and are eventually cut off at or before maturity by a cross-partition. This process, on being repeated again and again undisturbed, exhibits in the result the appearance of a copiously-branching hypha formed of bead-like cells. This connection, however, is very easily broken, and the usual appearance presented is that of small aggregates of sprouting cells.

Cell-contents.—The cells of fungi are characterised rather by the absence of certain cell-contents than by the presence of peculiar substances. The absence of chlorophyll and of starch-grains is most note-

worthy, and the physiological substitution for the latter of fatty matters *Glycogen* is found in the asci of Discomycetes and Tuberaceæ, and in the vegetative cells of these and other fungi. *Crystalloids* of albuminoid substance occur in the sporophores of Mucorini. *Nuclei* are of frequent occurrence in the cells of the reproductive system of many fungi, and in the vegetative cells of the Saprolegnieæ and other groups, though absolutely satisfactory evidence has not been produced with regard to their common presence in the thallus of fungi, apart from the reproductive system. Unsegmented hyphæ resemble the Siphoneous Algæ in their multinucleate condition, though single cells of segmented hyphæ also sometimes contain several.

Cell-membrane.—The typical cell-membranes of fungi consist of a substance differing from cellulose in its behaviour to the usual reagents, which is therefore known by the special name of *Fungus-cellulose*. Richter contends that this difference is more apparent than real, since he found that after prolonged maceration in a 7–8 per cent. potash solution, membranes exhibited the ordinary cellulose reaction which had refused the reaction after treatment of the usual kind with boiling potash solution, &c. De Bary, who originally gave the name of Fungus-cellulose to the membranes of fungi, welcomes these observations as confirming the undoubted near affinity of such membranes to ordinary cellulose, but points out that it merely proves that the membranes of the fungi operated on by Richter are altered by maceration with potash in the way described. 'Whether this alteration consists in the removal of some substance which was present from the first must remain uncertain; such an explanation has not been proved, and others are at least possible' ('Comp. Morph.,' p. 13). It is specially noteworthy that the membranes of certain filamentous fungi (Saprolegnieæ, &c.) exhibit the ordinary blue reaction with iodine.

The membranes of fungi undergo thickening and stratification, but are rarely pitted. The capillitium-threads of Lycoperdon are pitted, for example, and spiral and annular thickening occurs in the corresponding structure in Batarrea. Membranes very frequently become coloured with age, and undergo sclerosis; while in contrast to that condition other membranes often occur, which, though hard and horny in a dry state, swell up enormously by absorption of water, and acquire a gelatinous consistency.

Calcium oxalate, though of common occurrence in various forms in fungi, is of extreme rarity within their cells; Russula adusta (Fr.) and Mutinus caninus (Fr.) are the only recorded instances known to us.

Differentiation of the Thallus.

It is usually possible to draw a fairly sharp distinction between the vegetative part of the thallus, the *mycele*, and that portion specialised for the bearing of propagative bodies, the *sporophore*. In reduced forms, however, such distinction will be seen to be arbitrary, since propagative cells are formed in such cases from otherwise undifferentiated hyphal cells. The mycele, which serves the same physiological purposes as roots in point of the absorption of nutriment and of attachment, occurs in a variety of forms. The simplest and the typical form is a loose filamentous condition, spreading on or in the substratum, and sometimes provided with special branches termed *haustoria*, which penetrate cells of living plants and act as special organs of nutrition and of attachment. Such are the haustoria of Peronosporeæ and Erysipheæ, &c. A denser condition of mycele is represented by a membranous state to be found in forms vegetating on the surface of fluids and in fungi which attack wood, such as dry-rot (Merulius lacrymans, Fr.) and very many others. Strands of mycele are formed frequently in the Gasteromycetes and other fungi possessing compound sporophores. The strand-mycele of Agaricus melleus (L.), formerly known as *Rhizomorpha*, possesses a highly-developed structure, in which a cortical and a medullary region may be distinguished. Most compact of all forms of mycele are those known as *sclerotes*—densely compacted tuberous bodies. These consist of a central medullary tissue composed of very compact uniform hyphæ, sometimes pseudo-parenchymatous, enclosed by one or more layers of cortical cells with sclerosed walls. They are resting states of mycele, and act as stores of reserve material. With sclerotes may be classed such sclerotioid bodies as Pachyma and Mylitta, the develop-



Fig. 269.—Intercellular mycelial hyphæ (*m*), with haustoria penetrating into cells (*z*), *A*, of *Cystopus candidus* Lév.; *B*, of *Peronospora calotheca* de By. (× 350). (After de Bary.)

ment of which is quite unknown. Sclerotes were formerly classed together under the generic name of Sclerotium before it was recognised that such bodies are mere growth forms, or rather resting forms, of myceles of different fungi. Such a strand-mycele as that of Agaricus melleus (L.), formerly considered an independent fungus (Rhizomorpha), noticed above, has been aptly called a sclerote with a growing-point. B. Frank (Ber. Deutsch. Bot. Gesellsch., 1885) has described masses of fungal hyphæ having a dense sclerotioid structure investing the roots of trees (Cupuliferæ, Salicaceæ, Coniferæ) and intimately associated with the superficial cells of the root. To these bodies, termed *mycorhiza*, a symbiotic relationship with the trees in question is attributed, viz., the



FIG. 270.—*Agaricus melleus* L. Longitudinal section through apex of rhizomorph-strand (× 250). (After de Bary.)

absorbing function of root-hairs, the formation of which mycorhiza suppresses.

The *sporophores* of fungi are either simple or compound. They arise from the mycele, and produce the spores or definite organs of propagation. Simple sporophores are commonly erect branches of the mycele which either remain unbranched or branch again. The spores are borne at the extremities, and with the production of these bodies the growth of the sporophore either ceases altogether, or it may be renewed with farther production of spores. The modes of branching and the forms of sporophore are characteristic of *species* of fungi. Compound sporophores are structures often of considerable size, of

varying external appearance, and of a certain degree of differentiation of tissue. This differentiation is exhibited in the formation of layers, sometimes pseudo-parenchymatous, adapted for special functions. They are

FIG. 271.—Section through sclerote of *Sclerotinia Sclerotiorum* Lib., showing its cortical medullary tissue (× 375). (After de Bary.)

commonly characterised by the bearing of spores, or special spore-mother-cells, on definite regions and in more or less definite arrangement. They form, as a rule, that portion of the whole organism which

FIG. 272.—Simple sporophores of *Phytophthora infestans* de By. *a*, formation of first spores (zoosporanges) at ends of branches; *b*, two ripe spores on each branch and a third being formed (× about 200). (After de Bary.)

is popularly termed a fungus, such as the stalk and cap of mushrooms, the peridium of puff-balls, and the like. The structure of these is, as has been said, very diverse, and it will be found described in more or

less detail under the different groups of fungi. It may be well to note that transition-forms occur between the simple and compound types of

FIG. 273.—*Agaricus dryophilus* Bull. *a*, compound sporophore, longitudinal section showing the course of the hyphæ, a very young complete specimen 1·3 mm. in height, first beginnings of pileus; *b*, older specimen with pileus 2·5 mm. in breadth; *l*, piece of a lamella (slightly magnified). (After de Bary.)

sporophore: for example, Penicillium, though commonly simple, sometimes produces tufts of sporophores formerly supposed to belong to a different fungus under the generic name of Coremium.

SPORES.

The prevailing mode of spore-formation is by acrogenous *abjunction*. The terminal portion of the mother-cell or a special protuberance formed on it is cut off by a transverse wall, and this daughter-cell then drops off as a spore. The *basidiospores* of Basidiomycetes may be taken as an example. Finely pointed processes are formed on the summit of the basid, and these swell into ball-like form at the apex. The globular body is then abjointed and set free as a spore. Series or chains of spores are successively formed in like fashion in Cystopus, Penicillium, Uredineæ, &c. Spores are also endogenously formed within mother-cells—*sporanges*—and these are of two kinds, motile and non-motile. Examples of non-motile spores thus formed are to be found in Mucor, and in the *ascospores* of Ascomycetes. Such spores are either set free by the disappearance of the sporangial wall or by internal causes effecting their ejection. Motile spores or *zoospores* (swarm-spores) possess the power of moving freely in water by means of fine whip-lashes or cilia, and examples of these are to be found in the Saprolegnieæ and Peronosporeæ, the groups presumably most nearly related to Algæ. That the phenomenon of the production of swarm-spores is one nearly akin to that

FIG. 274.—*Puccinia graminis* Pers. *t*, teleutospores; *u*, uredospores (× 390). (After Sachs.)



FIG. 275. *m*, mycelial branch of *Cystopus Portulaceæ* Lév. producing two basids abjointing spores, in series; *b*, sporophore of *Eurotium herbariorum* Lk. with sterigmata; *s* and *t*, portions showing sterigmata, *p*, *p*, with their spores, *n* being the youngest (*a* × 390, the rest × 300). (After de Bary.)

FIG. 276. *Peziza* (*Pyronema*) *confluens* Pers. *a*, small portion of hymenium; *p*, paraphyse attached to, not originating in, hyphal branches from which the three asci spring; *m*, young asci; *r*–*w*, successive stages, according to letters, in the development of ascospores within asci (× 390). (After de Bary.)

of simple germination by the emission of a germ-tube is manifest. The example of Phytophthora infestans (de By.) illustrates this. The acrogenously-formed zoosporange produces zoospores in pure water containing free oxygen in fair amount. In nutrient solutions, on the other hand, no zoospores are formed, and the potential zoosporange simply emits germ-tubes. Germination of spores, however, takes place characteristically in fungi by the emission of *germ-tubes* under conditions of temperature, moisture, and the like presently to be discussed. Germination by the formation of *sprout-cells*, however, occurs in a number of forms.



FIG. 277.—Zoosporanges of *Phytophthora infestans* de By. *a*, division completed; *b*, escape of zoospores; *c*, free zoospores; *d*, spores come to rest and germinating (× 390). (After de Bary.)

Sexual Reproduction.

This subject is incidentally so fully discussed under the different groups, that nothing need be said here beyond calling attention to the fact that it falls under the same types as in Algæ. Such a form as Polyphagus may, however, be mentioned since it exhibits a type apart from ordinary isogamous or oogamous reproduction.

Conditions of Germination.

FIG. 278.—Ascospores of *Helvella esculenta* Pers. Stages of germination in order of letters (× 390). (After de Bary.)

Spores may be divided into two categories with reference to their power of germination, viz. those, by far the majority, which are capable of germination from the time of maturity, and those which must undergo a period of rest. Of the first kind a considerable number, generally speaking thin-walled watery spores, *sporids*, or *zoospores*, do not retain this power for more than a period measured by hours or days. The conditions under which they are kept are, as will be expected, of importance in this respect. Many spores retain the power of germination for a long period if kept in an air-dry

state. This time extends in numerous instances to one or two years, and, in the case of spores of Ustilagineæ especially, to longer periods. The spores of Tilletia caries (Tul.) germinated after eight and a half years, and those of other species after shorter intervals varying from seven and a half downwards. *Resting-spores*, or those of the second category, generally undergo more or less definite periods of rest corresponding to periods of vegetation. While germination cannot be procured before the lapse of this time, they frequently exhibit inability to survive the occasion presumed to be favourable. Such are hibernating spores like the teleutospores of Uredineæ, and the oosperms of Peronosporeæ. Resting-cells belonging to saprophytes again, e.g. the zygosperms of Mucorini, while they undergo a necessary period of rest of varying duration, yet display no partiality for seasons of the year, and this also is intelligible in view of their mode of life.

The power of resistance of spores to external agencies operating against their vitality is in many instances very great. The spores and zygosperms of several Mucorini withstand mechanical injury and repair slight wounds while preserving the power of germination. Short-lived spores and those of aquatic fungi do not bear desiccation ; but a great number of spores retain the power of germinating, as has been stated, for considerable periods in an air-dry state. Spores adapted for hibernation in temperate climates, and, it may be assumed, long-lived spores, withstand very low degrees of temperature, ranging below zero C. ; while such long lived spores, on the other hand, are very sensitive to high temperatures. The capacity for germination after exposure to high temperatures is maintained or not within certain degrees, according to the dryness or humidity of the environment. Thus it has been shown that while no spores are known to withstand a temperature of 100° C. in water or watery vapour, and many perish under these circumstances at much lower temperatures, the same spores can endure a considerably higher temperature in a dry state. Dry spores of some fungi have been found to withstand temperatures up to 120° C. and beyond, but it is probable that 130° C. marks the death-point of all. Others, again, perish at degrees considerably below 100° C. It must, of course, be borne in mind that the duration of the exposure is an important factor in such experiments, and that spores which support a high temperature for a few minutes or an hour are killed by longer exposure. It is probable that much individual variation exists in regard to this matter and to the duration of life under ordinary circumstances, and that in this lies the explanation of conflicting results obtained by different experimenters.

There is not much known as to the maximum, minimum, and

optimum temperatures at which the actual germination of spores takes place. According to Wiesner the minimum for the spores of Penicillium glaucum (Lk.) is from 1·5° C. to 2°, the optimum about 22° C., and the maximum 40° C. to 43°. This may be taken as fairly illustrative of other fungi in temperate countries in the open. Those which germinate, like many Mucorini and fungi inhabiting excrement, in the digestive tract of warm-blooded animals, have a much higher minimum, and an optimum agreeing with the body temperature. A supply of water and of oxygen must accompany the favourable temperature in all cases, and of nutrient substances in some. Speaking generally, parasites germinate freely in pure water or vapour, and saprophytes in nutrient substances, but the spores of many fungi germinate in both.

Conditions of Vegetation.

Under this head it will be necessary to consider little else than the nutritive adaptation of fungi, since they resemble other plants in the general conditions of vegetation, in their dependence on temperature, light, &c. The optimum temperature varies, as might be expected, in the cases of fungi which flourish at different seasons of the year, and in different climates. The optimum temperature for the growth of mycele in Penicillium glaucum is about 26° C., while that of spore-formation is the same as that of germination, about 22° C. These figures may be taken as fairly illustrative.

Luminosity [1] is exhibited by a considerable number of fungi—Agaricus olearius (DC.) and the rhizomorph form of A. melleus, Polyporus annosus (Fr.), and P. sulphureus (Fr.) (Europe), Agaricus igneus (Tul.) (Amboyna), A. noctilucens (Lév.) (Manilla), A. Gardneri (Berk.) (Brazil), A. lampas (Berk.) (Australia), A. Emerici (Berk.) (Andaman Islands), a species of Didymium (Jamaica), and probably by a number of other forms, the evidence as to which is doubtful. It is a phenomenon dependent upon the life of the organism, and the progress in it of destructive metabolism.

As regards nutrition, the absence of chlorophyll and the consequent inability to decompose carbon dioxide drive fungi to seek for organic carbon-compounds. In taking up food, fungi cause chemical changes in the organic bodies which furnish the food, e.g. fermentation. The well-known ferment-fungi need only be mentioned. It is in the highest degree probable that the solvents secreted by such fungi as penetrate dense woody and other structures are ferments. All fungi may be

[1] Vines, *Lectures on the Physiology of Plants*, p. 317; see also Phillips, *Proc. Woolhope Club*, 1888.

divided primarily into such as feed on the decaying bodies of plants and animals and dead organic substances—*saprophytes*—and those which attack living bodies—*parasites*. Between the two extremes of *strict saprophytes* and *strict parasites* there are intermediate forms. Some saprophytes, which ordinarily live throughout their course of development as such, have the power of living as parasites either wholly or during a part of their course of development. Such are called *facultative parasites*. Similarly some parasites which ordinarily live as such have the power of passing at all events a part of their lives as saprophytes. Such are *facultative saprophytes*. The *lichen-forming fungi* which live socially with algæ may be placed in another category. Most fungi are saprophytes, and it is obvious, from the fact that so many are confined to specific substrata, that there is much variation in the nutritive adaptations of such forms. These adaptations are, however, more clearly marked in the case of the smaller number which lead parasitic lives. Some are confined to single species of host-plants; many range over allied species, some of them attacking plants outside the group mostly affected, or exempting from attack certain species within the group. Others, again, may be said to be omnivorous parasites, attacking plants or animals of diverse groups. With regard to the predisposition of the host to the attack of the parasite, it is impossible for the most part to say exactly wherein it lies. Reference may be made to the case of species of Pythium which as facultative parasites attack Phanerogams, &c. The amount of water present in the host determines there the degree of predisposition to attack. While a sickly condition may constitute a predisposing cause to the attack of a parasite in some cases, it is by no means so in the majority of instances. It has been asserted that certain cultivated plants, such as cereals and the potato-plant, have by cultivation acquired an 'inherent tendency' to certain parasitic diseases, whereas it is obvious that the growing together of vast numbers of these plants furnishes opportunity for the spread of diseases which, in the absence of other evidence, may be taken to account for extensive outbreaks. Parasites commonly attack their hosts by the penetration of the membranes of the uninjured host, though cases are numerous where the entry is made by means of the stomates, or of wounded surfaces only. Most are *endophytes*, but a small number—e.g. Erysiphe—are *epiphytes*, which send haustorial branches into the body of the host. The result of attack is either the destruction of the host, or the production of deformities by anomalous processes of growth in the parts affected.

Lichen-forming Fungi.

These are strictly parasitic fungi which, without the aid of algal hosts, do not develop beyond the earliest stage of germination. Their nutritive inter-relations with their hosts, however, mark them off from other parasites. The hyphæ of the lichen-fungus embrace the algal cells, and the two elements together compose a thallus of definite form. The algal cells form by means of their chlorophyll-contents organic carbon-compounds by which the fungal cells benefit; but here the resemblance to true parasitism ceases. The host exhibits no sign of exhaustion, since a reciprocal accommodation exists between the two elements. The rhizoid filaments of the fungus draw from the substratum mineral substances, the raw material of food. The hyphal cells are fed by the exosmose of starch and the like from the algal cells, and the inference is justifiable that the algal cells receive in exchange by endosmose the waste products of the fungal protoplasm. There thus exists a lasting *consortism* or *symbiosis* between the elements, and the result is a thallus which may be treated from the point of view of the systematist as an autonomous organism. It must never be forgotten, however, that it is fundamentally two organisms, one of which, the fungal, cannot live without the other, while the latter can and does exist separately and independently in nature. It is a question not definitely decided whether certain algal forms thus living in consortism can or cannot live separately, and it is also doubtful whether the fungal portion of such lichens as live on the bark of trees or substrata rich in humus, does not live partially as a saprophyte. Evidence certainly points in this direction. The symbiotic relations existing in lichens are comparable with those described by Geddes, Brandt, and others, as in operation in Radiolarians and other animals, the 'yellow-cells' of which are actively vegetating algæ.



Fig. 270. Algal cells of Lichens. *A*, spore of *Physcia parietina* Nyl. germinating on *Protococcus viridis* Ag. *B*, *Synalissa symphorea* Nyl. with *Glococapsa*. *C*, *Cladonia furcata* Hoffm. with *Protococcus*. *D*, *Stereocaulon ramulosum* Ach. with *Scytonema*. (*A*, *B*, and *C*, × 950, *D* × 650.) (After Bornet.)

By far the greater number of lichen-forming fungi are Discomycetes or Pyrenomycetes. A few small tropical genera, Cora (Fr.), Rhipidonema (Mattir.), Dictyonema (Mont.), and Laudatea (Johow.), are Basidiomycetous, and two other tropical forms, Emericella (Berk.) and Trichocoma (Jungh.), have recently been declared by Massee (Phil. Trans. Roy. Soc. Lond., vol. 178, p. 305) to be Gasteromycetous Lichens. The two last-named cases are by no means satisfactorily established, and much more and better evidence must be forthcoming before they can be adopted as lichen-forming fungi. Propagation is effected by the spores of the fungal thallus, and an adaptation exists in certain lichens examined by Stahl for the supply of algæ to the new lichen. Algal cells, the offspring of the thallus algæ, which have been carried up into the hymenium, are cast out along with the spores, so that, both falling in the same neighbourhood, the germ-tubes of the spores find suitable hosts at once. This primary synthesis, however, probably takes place comparatively rarely in lichens as a whole. Propagation is very abundant by means of *soredes* or brood-buds



FIG. 280.—*Coccocarpia molybdia* Pers. Transverse section of thallus. *or*, upper, and *ur*, under cortical layer. *m*, so-called medulla; *g*, algal cells; *r*, rhizoids (× 600). (After Bornet.)



FIG. 281. *Ephebe pubescens* Fr. Branch of thallus with two young lateral branches (*s*); *g*, algal cells; *h*, hyphæ (× 500). (After Luerssen.)

consisting of one or more algal cells surrounded by hyphæ which separate from the parent-thallus. As a rule one species of alga furnishes all the algal cells of a lichen ; more rarely two, and then one prevails in abundance



FIG. 282.—*Usnea barbata* Fr. Development of soredes. *a*, group of eight algal cells attached to hypha ; *b*, similar group with branching hypha ; *c*, sorede with algal cell in optical section ; *d*, sorede with algal cells divided ; *e*, *f*, germinating soredes (× 500—700). (After Schwendener.)

over the other. The same species of alga, however, may be found in consortism with different species of fungus, and taking part in the composition, therefore, of differently formed thalli—different lichens in short.

FIG. 283.—*Cetraria islandica* Ach., a fruticose lichen (natural size).

Stahl experimentally proved this in his successful attempts at lichen synthesis. The algæ which furnish the hosts belong to different groups, and both unicellular and filamentous forms occur.

The thallus of lichens is of two sorts, the *heteromerous* and the

homoiomerous. The heteromerous thallus mainly consists of the fungus body of the lichen differentiated into a *cortical layer* and a *medullary layer*, the algæ occurring either as a definite layer where the cortical and the medullary hyphæ join, or they are scattered throughout the medulla, or form a dense mass in it. Such thalli exhibit considerable variety in forms of growth, and are called *foliaceous*, *fruticose*, *crustaceous*,



FIG. 284.—*Roccella tinctoria* DC. A filamentous lichen. Small plant (natural size).

&c., in descriptive works. The homoiomerous thallus consists of algal cells and hyphæ more or less equally distributed and alike in bulk. Collema, referred to below, is a *gelatinous* lichen, exemplifying this structure. Though the fungus does not actually prevail in bulk, it modifies the form of the thallus.

Until comparatively recent times, lichens were considered to be

independent organisms, the algal portion, the so-called 'gonidia,' being regarded as only specially developed cells arising from the colourless cells of the thallus. In 1868 Schwendener first accurately determined their dual nature, though de Bary had two years before indicated the possibility of this state of things in the case of the Collemaceæ, &c. The discovery so far, though sufficiently convincing, was based on anatomical considerations only, but the matter was finally proved, as well as a thing can be proved, by the experiments of Bornet, Treub, Reess, and Stahl. Reess succeeded in producing the thallus of Collema by synthesis, and Stahl went a step farther, and effected the formation of no less than three species of lichen. His observations on the relations of the algal and fungal elements of the lichen-thallus crowned the work of demonstration of its dual nature. Many systematic lichenologists who have been unable to shake off the traditions of their study still cling to the old view of the independent nature of lichens. It is hardly necessary to point out that the judgment of morphologists on such matters is the one to be trusted, especially as the matter has once and for all passed beyond the state of trust in authoritative opinion into the perfect state of complete proof.

Literature (Books of General Reference).

De Bary—Vergleichende Morphologie u. Biologie der Pilze, Mycetozoen u. Bacterien (Leipzig, 1884). [Translation by Garnsey and Balfour, Oxford, Clarendon Press, 1887, referred to in text as de Bary, Comp. Morph., &c.]

(In the above book a complete guide to the detailed morphological literature will be found.)

Systematic.

Saccardo—Sylloge Fungorum (1882, in progress). This work is intended to include all known Fungi.

The student should also consult the numerous works of Fries, dealing chiefly with Basidiomycetes; Corda's Icones Fungorum (Prag, 1837–54); and for British Fungi, Berkeley's Outlines of British Fungology (1860); Cooke's Handbook of British Fungi (1871); the same author's Illustrations of British Fungi (1881, in progress); Stevenson's Hymenomycetes Britannici (Edinburgh, 1886); and Phillips' Handbook of British Discomycetes (London, 1887).

Diseases of Plants caused by Fungi.

Frank—Krankheiten der Pflanzen (Breslau, 1880–81).

Sorauer—Handbuch der Pflanzenkrankheiten (Berlin, 1886).

Smith, W. G.—Diseases of Field and Garden Crops (London, 1884).

Literature of Lichen-thallus.

This literature is too vast to be quoted here in detail, but the reader is referred to the following essential papers.

Bornet—Recherches sur les gonidies d. Lichens (Ann. Sc. Nat., sér. 5, xvii. and xix.).

Johow—Ueber Westind. Hymenolichenen (Sitzber. Berl. Acad., 1884).

Reess—Ueber d. Entstehung d. Flechte Collema glaucescens (Monber. Berl. Acad., 1871).
Reess—Ueber d. Natur d. Flechten (Samml. wiss. Vorträge von Virchow u. v. Holtzendorff, 1879).
Schwendener—Die Algentypen d. Flechtengonidien (Basel, 1869).
Schwendener—Erörterungen z. Gonidienfrage (Flora, 1872).
Schwendener—Die Flechten als Parasiten d. Algen (Verh. d. Basel. naturf. Ges., 1873).
Stahl—Beitr. z. Entwickel. d. Flechten, ii. (Leipzig, 1877).
Treub—Lichenencultur (Bot. Zeit., 1873).
Treub—Onderzoek. over d. Natuur d. Lichenen (Diss.) (Leiden, 1873).

SPECIAL *literature is quoted under each group.*

GROUP I.—PHYCOMYCETES.

Class XVIII.—Oomycetes.

ORDER 1.—PERONOSPOREÆ.

The thallus of the Peronosporeæ consists of irregularly and copiously branched hyphæ inhabiting for the most part the living, and especially the chlorophyll-bearing, tissues of terrestrial flowering plants of different natural orders. The mode of life in this case is parasitic, and the hyphæ usually follow the intercellular spaces, and in many cases send short processes termed *haustoria* into the adjoining cells. These haustoria are variously formed according to the species. They are generally branched in Peronospora (Corda) and globular in Cystopus (Lév.). The hyphæ of other species (Phytophthora, de By.) traverse the cells of the host-plant. Transverse walls do not commonly appear in the hyphæ until the period of the formation of reproductive organs. The effect of this parasitic mode



FIG. 285.—Intercellular mycelial hyphæ (*m*), with haustoria penetrating into cells (*z*). *A*, of *Cystopus candidus* Lév.; *B*, of *Peronospora calotheca* de By. (× 390). (After de Bary.)

of life on the host is extensive destruction of the tissues, usually ending in death. Hypertrophy is produced in other cases, especially at the time of the formation of oosperms, leading to swellings and distortions of the parts affected.

Of the species of Pythium (Pringsh.) transferred to this order by de Bary (Bot. Zeit., 1881) from the Saprolegnieæ, some are saprophytes inhabiting the dead bodies of both plants and animals, while others are both parasites and saprophytes.

The *oogones* are globular cells with either a smooth or a granulated wall of some thickness, situated, as a rule, terminally, or more rarely interstitially. Soon after separation by a transverse wall from the hypha which bears it, the protoplasm of the oogone, which is rich in drops



FIG. 286.—Fertilisation of Peronosporeæ. *I.—VI.*, *Pythium gracile* Schenk. Successive stages according to numbers (× about 800). *VII.*, *Peronospora arborescens* de By. Oosphere is invested with a thick membrane, outside of which is the periplasm contracting to form outer coat of oosperm (× 600). (After de Bary.)

of fatty matter, begins to collect into a central mass containing the drops and bounded by a hyaline layer. Outside this central body (*oosphere*) there is left over a clear mass of protoplasm (*periplasm*), which fills up the space between it and the wall. While the oogone is thus developing, the *antherid* arises, either from the pedicel-cell of the oogone itself, or as the terminal cell of a neighbouring branch. It has commonly the form of an irregularly bent tube with an unthickened cell-wall, and at first ordinarily granular protoplasm. It applies itself closely to the wall of the oogone, and sends through it a delicate straight *impregnating tube*, which penetrates to the surface of the oosphere. The protoplasm of the antherid also undergoes about this period a differentiation into two masses ; one, threadlike but irregular, and occupying the middle,

contains the more granular particles, and is termed the *gonoplasm*, while the other (*periplasm*) surrounds it. The gonoplasm enters the oosphere through the impregnating tube of the antherid, and thus accomplishes the act of impregnation. Sometimes two, rarely more, antherids arise and apply themselves to the oogone, and this varies both with species and individuals. After impregnation the oosperm assumes a cellulose membrane, and gradually ripens. The fatty contents collect into one body occupying the middle, and the membrane becomes thicker and differentiated into two cellulose layers, the *extine* and the *intine*. The periplasm develops into a brown, often granulated and warty membrane, the extine, enclosing the oosperm, while the original wall of the oogone generally breaks up, but may in some cases persist.

The oosperms germinate in water after a period of rest generally lasting throughout the winter ; and this takes place either by the emission of a *germ-tube* which gives rise directly to a new thallus like the parent one, or the protoplasm divides into a number of *zoospores*, which, extruded together within a globular sac and escaping from it, swim for a short time, and, after settling down, push out each a germ-tube which produces a new thallus. In other species, again, both methods of germination occur, some of the oosperms directly emitting germ-tubes, while in the others the production of zoospores intervenes. In certain species, the oosperms of which produce a germ-tube directly, a short mycele (*promycele*) is formed, which, after bearing a few *conidiospores*, dies, and these conidiospores in turn propagate new thalli.

The non-sexual organs of propagation (*conidiospores*) are borne upon special branches of the thallus (*sporophores*) in a variety of ways characteristic of the genera and in a minor degree of the species. These germinate either by means of a germ-tube directly produced, or the contents break up into zoospores, which, after swarming, settling down, and becoming invested with a membrane, also produce germ-tubes.

The usual course of life is the production upon the thallus of vast numbers of conidiospores, which propagate the species extensively throughout spring and summer, followed in autumn by the bearing of sexual organs, with which the generation terminates. De Bary points out ('Comp. Morph.,' &c., 1884) that only in the instances above mentioned of the production from the oosperm of a promycele bearing a few conidiospores, can a distinct *alternation of generations* be recognised. There is indeed merely the succession of one oosperm-bearing generation to another, the propagating spores being only accessory products of the thallus. In such cases as Pythium vexans (de By.) and Artotrogus (Mont.), for example, there are no such organs of propagation at all, or at least long-continued research has failed to discover them. Other

species appear again to have lost the power of producing sexual organs, and this is notably the case in Phytophthora infestans (de By.), the potato-disease fungus, which succeeds in hibernating by means of a perennial mycele. In such a case the species is entirely dependent upon the propagating spores for distribution.

Cystopus (Lév.).—The thallus consists of hyphæ inhabiting the intercellular spaces of the tissue of flowering plants, and provided with haustoria. The oosperms are resting-cells which germinate in spring by means of the production of zoospores in the usual way. The propagating zoospores are borne in *zoosporanges* at the end of cylindrical or club-shaped *zoosporangiophores* in vertical series. A small broad swelling first appears at the apex, and then a transverse wall cuts off the upper portion, which rounds off and thus becomes the first and oldest zoosporange of the series. Then another is cut off in the same fashion, while the sporangiophore elongates. A series or chain is thus produced, each zoosporange joined to its neighbour by a very short and slender connecting stalk. The first cell at the top of the series has a thicker wall than the others, is yellowish in colour, and is, at least in the vast majority of cases, incapable of germination. If germination does take place, a germ-tube is said to be produced, while all the other members of the series give rise to zoospores. These chains of zoosporanges arise in dense masses side by side below the epiderm of the host, which is gradually ruptured, permitting their escape, the thick wall of the top member of the series serving as a shield in bursting the epiderm. When the zoospores germinate, their germ-tubes enter the host by way of the stomates, by this means attaining directly the intercellular spaces. The disease thus set up in the host is not so active as in the case of species of Peronospora, and the parts affected do not perish so rapidly. During the formation of oosperms in Cystopus candidus (Lév.), regions of the host undergo acute hypertrophy. The commonest species of the genus is C. candidus ('white rust'), which attacks a large number of Cruciferæ. Cabbages and the Shepherd's Purse (Capsella bursa-pastoris) suffer conspicuously from it, while the latter is often affected by Peronospora parasitica (de By.) in company with it. Other well-known species are C. Portulaceæ (Lév.) and C. cubicus (Lév.).



FIG. 287.—*m*, mycelial branch of *Cystopus Portulaccæ* Lév., producing two sporangiophores, *p*, bearing zoosporanges, *n*, in series (× 390). (After de Bary.)

Peronospora (Corda).—The thallus and the sexual organs closely resemble those of Cystopus. In both genera the passage of protoplasm from the antherid into the oogone is not directly visible, and the oogonial periplasm is more abundant than in the other genera. In germinating, the oosperm produces a germ-tube, but the process of germination has not been observed in a number of species, and, as de Bary says (Journ. Roy. Agric. Soc., 1876), it is quite possible that the species of Peronospora which, like Cystopus, produce zoospores from their 'conidia' (zoosporanges) present also the same phenomenon in connection with the oosperms. The sporophores of Peronospora commonly issue from the host-plant through the stomates, and are, for the most part, regularly and copiously branched. At the fine points of the branches the non-sexual propagating bodies are produced singly. These are in some species conidiospores germinating by the emission of a germ-tube, and in others zoosporanges producing zoospores. Conidiospores and zoosporanges are borne in precisely similar fashion, and present the same appearance up to the production of the germ-tube or zoospores, as the case may be. The zoospores are formed within, and escape from the original zoosporangial membrane and not from an extruded sac. Intermediate between these forms are the *plasmatoparous* species (P. densa, Rab., and P. pygmæa, Ung.), in which the whole protoplasm escapes from the spore in a mass through the opening of a papilla-like point in the wall, and, at once becoming globular, secretes a cellulose membrane, and subsequently germinates by the emission of a short thick germ-tube. The germ-tubes both of zoospores and of conidiospores penetrate directly the epiderm of the host and the cells underlying it, until an intercellular space is reached. This genus contains a large number of well-known parasites, such as P. viticola (de By.) on the vine, P. nivea (de By.) on Umbelliferæ, P. parasitica (de By.) on Cruciferæ, P. Schleideniana (Ung.) on onions; P. Viciæ (de By.), P. Trifoliorum (de By), &c. Hypertrophy is frequently produced in the host at the time of oosperm-formation, but not so acutely as by Cystopus. The oosperms of several species are unknown, and of these P. Rumicis (Corda) and P. Schachtii (Fückel) hibernate by means of their perennial mycele, while P. Ficariæ (Tul.), the oosperms of which are known, passes the winter in the same way.

Phytophthora (de By.).—This genus was founded for the reception of P. infestans (de By.), which was formerly placed in Peronospora. Industrious research has failed to discover the sexual organs and oosperms of this species. Mr. Worthington Smith claims to have found them, but the balance of evidence is distinctly against this. The sexual organs of Phytophthora omnivora (de By.) have been

observed, however, and fertilisation takes place in the usual way. A very small quantity of gonoplasm (not visibly differentiated) passes over into the oosphere. The antherids and oogones arise together in this species, and develop in close connection. The oosperms form each a promycele, as described above. The sporangiophores of



FIG. 288.—Simple sporophores of *Phytophthora infestans* de By. *a*, formation of first spores (zoosporanges) at ends of branches; *b*, two ripe spores on each branch and a third being formed (× about 200). (After de Bary.)

Phytophthora, which resemble those of Peronospora in general habit, differ from them in the fact that each branch bears more than one propagating body—not in chains, like Cystopus, but at intervals on the branch. In P. infestans a propagating cell is produced at the apex of each branch; and as it ripens a papilla-like swelling arises beneath it; the branch grows on and turns the cell aside. These propagating cells are usually zoosporanges, but not unfrequently they are conidiospores, differing from them in no other respect than the production of a germ-tube directly instead of zoospores. The zoospores are formed within and escape directly from the zoosporange itself, as in Peronospora.



FIG. 289.—Zoosporanges of *Phytophthora infestans* de By. *a*, division completed; *b*, escape of zoospores; *c*, free zoospores; *d*, spores come to rest and germinating (× 390). (After de Bary.)

Phytophthora infestans has a special economic interest, as the cause of the well-known potato-disease. The disease first appears, as a rule, on the green leaves of the potato plant in July or August, the sporangiophores emerging through the stomates. Sporanges are formed, under favourable conditions of temperature, moisture, &c., in a few hours, are

wafted away, and, falling on other potato leaves, there produce zoospores, or germ-tubes directly, in drops of water formed by dew or rain. The germ-tubes penetrate the epiderm, setting up fresh growths of mycele in new plants, and thus the disease is propagated. Countless numbers of such propagating cells, each potentially the mother of a number of zoospores, may thus be set free from a few diseased plants, and the spread of infection and destruction of tissue in warm moist weather is almost inconceivably rapid. The disease extends to all parts of the plant, including the tubers, in which the mycele often remains in a resting condition throughout the winter (as in certain species of Peronospora mentioned above), and from which a fresh start is made in the following year. The interest attaching to the subject is mainly economic, and an extensive literature bearing upon it has grown up—by far the greater part of it utterly worthless.

Pythium (Pringsh.).—Several species of this genus are saprophytes, inhabiting the dead bodies of plants and animals, while others are true parasites on fresh-water algæ, on prothallia, and on flowering plants. The thallus and sexual organs are of the type described. The oosperms of P. proliferum (de By.), like those of Phytophthora omnivora, form a promycele ; while of P. vexans (de By.) the oosperms only are known. The formation of propagating spores occurs at the end of simple thallus-hyphæ. A terminal cell is cut off by a transverse wall, and usually becomes a zoosporange. This body expands at the apex into a thin globular sac, into which the whole of its protoplasm empties itself. There zoospores are differentiated, and, bursting the sac, escape and germinate. There is some variation according to species in the forms of the zoosporanges ; sometimes they are round or oval and sometimes elongated. They have not the definite arrangement which characterises the other genera. P. intermedium (de By.) and P. de Baryanum (Hesse) sometimes form spores which emit a germ-tube instead of the usual zoosporanges. P. gracile (Schenk), P. entophytum (Pringsh.), and P. Chlorococci (Lohde) inhabit fresh-water algæ, P. Equiseti (Sad.) and P. circumdans (Lohde) attack prothallia, while P. de Baryanum infests seedlings of different phanerogams and fern-prothallia. The last-named is capable of attaining full development as a saprophyte on both dead plants and animals. P. intermedium, also saprophytic, becomes a parasite on fern-prothallia. It is worthy of note that these fungi are parasitic only on seedlings, prothallia, &c., which contain abundance of water ; and though P. de Baryanum causes local injury to grown plants, this power may be raised to one of destruction under water. Pythium vexans is found in diseased potato tubers.

Fossil Form.

Peronosporites (W.G.S.).—This genus was founded by Mr. Worthington Smith for the reception of a fossil fungus Peronosporites antiquarius (W.G.S.), first detected by Mr. Carruthers in the axis of a Lepidodendron from the coal measures. Mycele and bodies which may well be oogones are visible in the preparations. The fungus is probably nearly related to Pythium.

Literature.

De Bary—Recherches sur le développement de quelques Champignons parasites (Ann. Sc. Nat., 4 sér., Tom. xx.). (Contains reference to older literature.)
De Bary—Zur Kenntniss der Peronosporeen (Beitr. zur Morph. u. Physiol. d. Pilze, ii.).
De Bary—Untersuch. über die Peronosp. u. Saprolegn. (*ibid.*, iv.).
De Bary—Research into the Nature of the Potato-fungus (Phytophthora infestans de By.) (Journ. Roy. Agric. Soc., 1876, xii.).
De Bary—Zur Kenntniss der Peronosporeen (Bot. Zeit., 1881).
Cornu—Monogr. d. Saproleg. (Pythium) (Ann. Sc. Nat., 5 sér., Tom. xv.).
Hesse—Pythium de Baryanum, Halle, 1874.
Millardet—Le Mildiou (Paris, G. Masson, 1882; and Journ. d'Agric. pratique, 1881, T. i., No. 6, and 1882, T. ii., No. 27).
Pringsheim—Die Saprolegnieen (Pythium) (Jahrb. wiss. Bot., i.).
Schröter—Peronospora obducens (Hedwigia, 1877, p. 129).
Schröter—Protomyces graminicola (*ibid.*, 1879, p. 83).
W. G. Smith—Resting-spores (so called) of Potato Disease (Gard. Chron., 1875, iv., N.S.; and 1876, vi., N.S.).
W. G. Smith—Peronosporites antiquarius, W. G. S. (Gard. Chron., 1877). [See also G. Murray, Academy, 17 Nov. 1877 and following numbers; and Williamson, Phil. Trans. Roy. Soc. Lond., 1881.]
A. Zalewski—Zur Kenntniss der Gattung Cystopus (Bot. Centralb., 1883, No. 33).

Order 2.—Ancylisteæ.

This order embraces a few genera which, so far as what is known of them indicates, are related most nearly to Pythium. All the members of the group are parasitic in fresh-water algæ (Cladophora, Mougeotia, Spirogyra, Mesocarpus, Closterium, &c.), and they are all farther characterised by simplicity of structure. The thallus is represented by hyphæ at first undivided, which often extend from one end of the host-cell to the other. Ancylistes Closterii (Pfitz.) displaces the chorophyll-plates of its host, and ultimately causes the death of the cell. Lagenidium (Schenk), found in filamentous algæ, causes the separation of cell-contents from cell-wall, and discolours the chlorophyll, which gathers together into a mass.

The sexual organs are formed by the division into cells of the thallus-hyphæ by transverse walls. Of these cells, some swell and become oogones, while others remain small and act as antherids (Myzocytium,

Schenk); or different individuals produce the oogones and the antherids (Lagenidium, Ancylistes). A perforation having been made in the oogonial wall, the whole of the protoplasm of the antherid empties itself into the oogone (there being no periplasm), and the united mass rounds itself off and becomes the oosperm. The germination of the oosperm has not been observed.

Propagation takes place by means of zoospores (Lagenidium), and to this end either the whole thallus-hypha becomes transformed into a zoosporange, or it is divided into a series of such. Each zoosporange sends out through the membrane of the host-cell to the surrounding water a protuberance, through which the contents escape after the fashion of Pythium, forming uniciliated zoospores, which ultimately attack the fresh cells of other algæ. In Ancylistes the only propagation known is a process of extension of its hyphæ from one host to another.

Literature.

Cornu—Monogr. des Saprolegn. (*loc. cit.*).

Cornu—Note sur l'oospore du Myzocytium proliferum, Schenk (Bull. Soc. Bot. France, xvi., 1869, p. 222).

Pfitzer—Ein neuer Algen Parasit (Monatsber. Berl. Acad., 1872).

Schenk—Ueber das Vorkommen contractiler Zellen im Pflanzenreich (Würzburg, 1858).

Zopf—Ueber einen neuen parasitischen Phycomyceten, &c. (Lagenidium) (Bot. Zeit., 1879, p. 351).

Order 3.—Monoblepharideæ.

The single genus Monoblepharis (Corn.), like the preceding group, is closely related to Peronosporeæ and especially to Pythium. The thallus-hyphæ bear both terminal and interstitial oogones, in which there is no preliminary differentiation of periplasm, but the whole protoplasm contracts and forms the oosphere, while the apex of the oogonial wall opens. The antherid (usually a cell adjoining an oogone) produces several swarming antherozoids, which escape, one of



Fig. 290. *Monoblepharis sphærica* Cornu. Filament bearing an oogone, *o*, and antherid, *a*, antherozoid, *s*. The numbers indicate successive stages (× 800). (After Cornu.)

them attaining and entering by the apical opening of the oogone, and uniting with the oosphere. The resulting oosperm has not yet been observed to germinate.

Propagation takes place by the formation of uniciliated zoospores in zoosporanges, from which they escape in the same way as those of Phytophthora.

LITERATURE.

Cornu—Monogr. des Saprolegn. (*loc. cit.*).

ORDER 4.—SAPROLEGNIEÆ.

The Saprolegnieæ, as their name indicates, are saprophytes on the dead bodies of both plants and animals in water ; with at all events the exception of the Saprolegnia of the salmon disease, which is both saprophyte and facultative parasite. The cause of the predisposition to this disease has not yet been exactly determined, as for example has been the case with those species of Pythium which possess a similar mode of life. Prof. de Bary points out with regard to them that susceptibility to disease in the host is in relation to the amount of water present. The problem in the case of the salmon disease has every appearance of being a more complex one. The Saprolegnieæ bear in other respects much resemblance to the Peronosporeæ, and especially to Pythium, which until recently was included among the former. Pythium indeed presents points of relationship with the types of Oomycetes in general ; and the relationship is rendered the more striking by the union in some of its species of both parasitic and saprophytic modes of life. The thallus-hyphæ of the Saprolegnieæ are usually of relatively large size, springing from slender *rhizoids* buried in the substratum.

The oogones arise, as in Peronosporeæ, on branches of the thallus-hyphæ. In most cases, however, several oospheres are formed in each oogone (sometimes as many as thirty or forty), and, no periplasm having been differentiated, the whole of the oogonial protoplasm is included in them. It happens in some cases that only one oosphere is formed, but the number is variable according to species, and also partly according to individuals. Pits arise, but by no means always, in the oogonial wall.

The antherids, which are commonly club-shaped, are produced on slender branches of the thallus ; and each antherid is borne either on the same hypha of the thallus as the oogone to which it is attached, or on a hypha which bears no oogones. The remarkable point about these antherids is their impotency, since no actual observation of the transference of protoplasm from them to the oospheres has ever been made, though they perforate the oogonial wall, and processes, like impregnating tubes sent through, come in contact with the oospheres. These processes grow from one oosphere to another, and may even

emerge again outside the oogonial wall, but they remain closed at all points, and after a day or two perish. In the case of certain species, the antherids never produce these processes, or the oogones may be without antherids. In other cases antherids are never produced at all, or only by way of rare exception. In the meantime the oospheres ripen into oosperms, while the antherids, if present, perish. Pringsheim has recently endeavoured to show that impregnation takes place in certain species by the transference into the oospheres of minute portions of antheridial protoplasm moving in amœboid fashion. De Bary points out that, while Pringsheim has not actually seen this, the sole evidence trusted to is that of stained preparations, which appear to exhibit open communication between antherid and oosphere, &c. In any case the observation does not affect those cases where antherids are either wanting, or do not produce the penetrating tubular processes.



FIG. 291.—*A* to *C*, *Achlya racemosa* Hildebr. At the end of *A* is an empty zoosporange, *s*, with empty zoospore membranes; at *a*, *b*, and *c*, are oogones with antherids, *a*, in an early stage, *b* and *c* as in *B*, oogone with two oospheres and an antheridial tube applied to one. *C*, ripe oosperm. *D*, *E*, *Achlya polyandra* Hildebr. *D*, oosperms germinating. *E*, germinating oosperm which has formed a sporange with a head of spores. (*A* × 145, *B* and *C* × 375, *D* and *E* × 225.) (After de Bary.)

The ripe oosperms thus parthenogenetically produced germinate after a period of rest varying from a few days to several months. Germination takes place, as in the Peronosporeæ, either by means of a germ-tube, or zoospores are produced.

Propagation is effected by the agency of zoospores produced in special zoosporanges, and also exceptionally by means of certain resting-cells formed by the mycele after transverse division of the hyphæ (Saprolegnia). These swell out into globular form, with thick membranes and plentiful protoplasm, and germinate by the emission of a germ-tube, or zoospores are formed in them ('resting sporanges' of

Pringsheim). These resting cells or sporanges are formed only on old myceles, and by no means regularly.

FIG. 292.—Zoosporanges of *Achlya* N. ab E. *A*, with zoospores formed but still enclosed. *B*, with zoospores escaping. At *a* they are invested with a cell-membrane, at *c* they are free, empty membranes at *b* (× about 300). (After de Bary.)

The zoosporanges vary in form with the genus, and in a minor degree with the species. The usual form is a large club-shaped zoosporange containing a great number of biciliated zoospores, which escape from it through an opening at the apex; though in poorly developed individuals (and normally in Aphanomyces, de By.), the zoosporange is more cylindrical, and only one row of zoospores is formed. In Saprolegnia (Nees ab Esenb.) the zoospores are actively motile when they escape. Their activity ceases for the most part after a few minutes; they settle down, assume a thin cellulose-membrane, rest for a short time, only to escape from this membrane, and resume active movement before final settling down and germination. The spores of individuals may on the other hand omit the second period of movement, and germinate directly on first settling down. In Achlya (Nees ab Esenb.) and Aphanomyces the spores escape from the sporange without cilia and active movement. They arrange themselves in globular fashion outside the apex of the sporange, assume each a thin cellulose-membrane, within which they rest for a few hours, and, escaping from it, swim about, and, settling down, ultimately germinate. In the sporanges of Dictyuchus (Leitg.) the spores are each enclosed in net-like cellulose walls, from which they escape, not by any special orifice of the sporange, but by taking as it were the shortest cut through the sporangial wall and empty spore-cases if they come in the way.

In Aplanes Braunii (de By.) the formation of propagating spores is as a rule omitted. When produced they give rise to germ-tubes directly without swarming.

The zoospores produced by oosperms behave, so far as is known like those from the corresponding zoosporanges.

The modes of zoospore-formation in Phytophthora and Cystopus, Pythium Achlya, and Aphanomyces, Dictyuchus, and lastly Saprolegnia, express in an interesting way the relationships of these genera.

LITERATURE.

De Bary – Beitr. zur Kenntniss der Achlya prolifera (Bot. Zeit., 1852).
De Bary—Einige neue Saprolegnieen (Pringsheim's Jahrb. wiss. Bot., ii.).
De Bary – Untersuch. über die Peronosp. u. Saprolegn. (Beitr. zur Morph. u. Physiol. der Pilze, iv.).
De Bary – Zu Pringsheim's Neue Beob. über d. Befruchtungsact der Gattungen Achlya und Saprolegnia (Bot. Zeit., 1883).
Cornu—Monograph. der Saprolegn. (Ann. Sc. Nat., 1872).
Hildebrandt—Mycolog. Beiträge, i. (Pringsheim's Jahrb. wiss. Bot., vi.).
Hartog—On the Formation and Liberation of Zoospores in the Saprolegnieæ (Quart. Journ. Micr. Sc., 1887).
Hartog—Recent Researches on Saprolegnieæ (Annals of Botany, 1888).
Huxley and Murray—Salmon Disease (Reports of Inspector of Fisheries, 1882, 1883, 1884, 1885. See also Quart. Journ. Micr. Sc., 1882, and Journ. Bot., 1885).
Leitgeb—Neue Saprolegnieen (*ibid.*, vii.).
Lindstedt—Synopsis der Saprolegn., Berl., 1872.
Pringsheim—Entwickelungsgeschichte der Achlya prolifera (N. Acta Acad. Leop.-Carol., xxiii., p. 1).
Pringsheim—Beitr. zur Morph. u. Systematik d. Algen, ii. Die Saprolegn. (Jahrb. wiss. Bot., i., ii., and ix.).
Pringsheim—Neue Beobacht. über d. Befruchtungsact von Achlya u. Saprolegnia (Sitzber. Berl. Acad., 8 Juni, 1882). Nachträgliche Bemerk. zu dem Befruchtungsact von Achlya (Pringsheim's Jahrb. wiss. Bot., xiv.).
Reinsch—Beobacht. über einige neue Saprolegn. (*ibid.*, xi.).
Thuret—Rech. sur les zoospores des Algues, 1851.
Ward – On Saprolegnieæ, and also on Pythium (Quart. Journ. Micr. Sc., 1883). Contain histological details.

Class XIX.—Zygomycetes.

ORDER I.—MUCORINI.

The Mucorini are for the most part terrestrial saprophytes, the remainder being parasites on other Mucorini. The thallus consists of a copiously branching hypha undivided up to the time of the production of spores or sporanges, when transverse walls first appear. Sexual reproduction is effected by the formation of a *zygosperm*, while spores and propagating cells (like some of the resting-cells produced by the mycele of Saprolegnia) are also borne, the former regularly and in characteristic forms, the latter only in special cases and under certain conditions. The production of a zygosperm is effected by the conjugation of two specially differentiated cells, *gametes*, not to be distinguished from each other by any mark or power of movement. The two cells thus contributing to

its formation either, by their simple fusion, themselves constitute the zygosperm, or this body is the direct offspring (daughter-cell) of the union. The *spores* are produced either in terminal *sporanges* or singly at the apex of a *sporophore*, or again serially in like fashion to the last. In a considerable number of cases the zygosperms are unknown, and it

FIG. 293.—*B*, *Phycomyces nitens* Kze. Plant grown on decoction of plums; mycele, *m*, sporophore, *g*. *A*, *C*, and *D*, *Mucor Mucedo* L. *A*, sporange in optical-longitudinal section. *C*, zygosperm (*z*) borne on suspensors. *k*, germ-tube; *g*, sporange. *D*, conjugation. *a*, *a*, gametes; *b*, *b*, suspensors. (*B* slightly, *A*, *C*, and *D* more highly magnified.) (After Brefeld.)

may be assumed, on the weighty authority of de Bary, that in certain of these they do not occur, since industrious observation has failed to discover them. They are known, in fact, only in nineteen species, though future research may bring a fair number more to light. Where they have been observed the life-history proceeds as follows. The germinating

zygosperm gives rise directly to a promycele bearing the characteristic spores, and these in turn produce on germination a mycele which bears spores again, and ultimately a zygosperm. It has been observed in an artificially nourished individual, that the germinating zygosperm at once produced a mycele which subsequently bore spores without the intervention of the promycele stage. In another instance (Sporodinia grandis, Link) zygosperms have been observed on a mycele which arose from a spore, before the production of sporophores upon it ; while it sometimes happens in this species that a zygosperm is produced on a mycele arising directly from a zygosperm without the intervention of spores at all. But in the great majority of cases the production of spores precedes the formation of a zygosperm on the same mycele. In many species the zygosperms are of rare occurrence, and an indefinite number of successive spore bearing generations come between zygosperm and zygosperm. Throughout the whole order spores are produced in vastly greater numbers than zygosperms. Syzygites (Ehrenb.) is the generic name given to certain forms of doubtful affinity which produce, so far as is known, zygosperms alone.

Sub-order 1 : Mucoreæ.—The members of this group are for the most part saprophytes on the excrement of animals, fruits, bread, saccharine fluids, &c. The thallus-hyphæ are relatively large and much ramified. The conjugating hyphæ arise either as branches of the mycele or on special hyphæ somewhat resembling sporangiophores, their place of origin being, in different instances, in either morphological or merely local approximation to each other. At an early stage of development they come into contact by their apices, and a firm connection between the two is established. Thus joined the development of each goes on, and soon a transverse wall cuts off the apical portion of each. This portion is a *gamete*, and the rest of the hypha, generally club-shaped, its *suspensor*. A pore next appears in the centre of the original wall separating the two gametes, and gradually the whole wall disappears and the contents conjugate. The *zygosperm* thus formed increases in size, drawing upon the contents of the suspensors. The protoplasm becomes dense, and the fatty contents gather into a large drop. The wall commonly becomes covered externally with warts or spines at all points except where the suspensors are attached. The form of the whole is roundish or drum-shaped, the smooth walls adjoining the suspensors corresponding with the sides of the drum. The wall is divided into two coats, the outer one (*extine*) brown or black, and the inner one (*intine*) stratified, and either entering the corrugations of the extine or remaining smooth along the surface of contact with it. The suspensors usually remain *in statu quo*, but in Rhizopus nigricans (Ehrenb.), where

one gamete is about half the height, though of the same breadth, as the other, the suspensor of the smaller one becomes greatly enlarged after conjugation, while the other remains as it was. In most cases the suspensors eventually decay, but in others (Phycomyces, Kze., and Absidia, Van Tiegh.) an outgrowth of darkly coloured hyphæ takes place from each suspensor and invests the zygosperm. In Mortierella (Coemans), which has a smooth extine, this outgrowth arises from the hyphæ bearing the suspensors (as well as from the suspensors in one case), and forms a compact integument of the zygosperm. In M. nigrescens (Van Tiegh.) this outgrowth begins after conjugation, first from the suspensors, then from the adjoining hyphæ; while in M. Rostafinskii (Bref.) the outgrowth



FIG. 294.—*Rhizopus nigricans* Ehr. Formation of a zygosperm. Stages according to Letters (× about 90). (After de Bary.)

takes place solely from the adjoining hyphæ, and begins so early that an investment is formed before actual conjugation takes place.

A phenomenon resembling that of the parthenogenesis of the Saprolegnieæ is exhibited by a number of the Mucoreæ in the formation of *azygosperms*. This occurs in Absidia, Sporodinia (Link), and Spinellus fusiger (Van Tiegh.), and the formation of these bodies ensues when gametes have failed to conjugate, and even when single gametes only are produced. They possess the structure and power of germination of normal zygosperms, just as the parthenogenetic oosperms of Saprolegnia do. Bainier states that Mucor tenuis (Link) forms only azygosperms, and de Bary suggests that the (as yet little known) Azygites of Fries may be found to exhibit this phenomenon.

On the germination of the zygosperm, as has been said, a *promycele* bearing sporanges is produced directly, and these sporanges have the

same structure as those that follow them. They are globular sacs borne at the end of sporangiophores, and the spores produced within them are never endowed with the power of active movement. The different forms of sporange and sporangiophore afford characters for the genera of the group. Mucor (Michel.), Pilobolus (Tode), Sporodinia (Link), Phycomyces, Rhizopus (Ehrenb.), Circinella (Van Tiegh.), and Absidia possess a peculiar conformation of the basal wall of the sporange. It bulges inwards in a conical or more or less oval form (see fig. 293 A), and presents an appearance which has suggested the name of *columel* for this peculiarity. Of the genera possessing a columel some are distinguished by a fugacious sporangial wall, others by a firm persistent one, while the mode of branching of the sporangiophore (or the absence of branching) and its general form, afford other generic characters. Mortierella has a fugacious sporangial wall but no columel. Thamnidium (Link), Chætostylum (Van Tiegh.), and Helicostylum (Cord.) have two kinds of sporange, the one kind like those of Mucor, and the other smaller (*sporangioles*) with a persistent wall, no columel, and containing but a few spores, which however resemble the others in function.

On old and on badly nourished myceles of some species, accessory propagating bodies are formed (*chlamydospores*, *stylospores*, &c.). All such accessory spores are capable of giving rise to normal characteristic myceles either at once or after a period of rest. In Mortierella single *acrospores* are borne on slender mycelial hyphæ. The old myceles and even the sporangiophores of Mucor break up into *resting-cells* like those of Saprolegnia with thick walls. The chlamydospores (Van Tieghem) of Mortierella are such bodies, and where they occur terminally, de Bary regards them as transitional forms to the acrospores of the same genus just mentioned. Brefeld and Van Tieghem have described (Mucor racemosus, Fres., &c.) another form of accessory propagating spores, which are produced in series or chains through transverse division of the mycelial hyphæ. These either remain joined together in conferva fashion, as Berkeley says, or they part company, and each such cell exhibits a yeast-like vegetation.

Sub-order 2: Chætocladieæ.—The mycelial hyphæ of Chætocladium (Fres.) become attached to the hyphæ of the Mucor-host, and, by the resorption of the cell-wall at the place of contact, effect a direct communication. At such places of attachment a large number of globular protuberances are produced close together, forming a body of considerable size, which may be regarded as a food-reservoir. The act of conjugation and the formation of the zygosperm agree in all essential particulars with the corresponding processes in the Mucoreæ. The intine

of the zygosperm has a smooth surface, not entering the external warts of the extine. Azygosperms have not been observed. The sporophores terminate in a fine hair-point, but below this give rise to a whorl of branches nearly at right angles to each other, terminating again each in a hair-point. These again branch more or less in like fashion. The ultimate branches become swollen, and on these swellings fine short sterigmata arise, each sterigma bearing a spore. The mass of spores thus produced has a bunch-like aspect.

Cunningham's Choanephora, found by him on the flowers of Hibiscus, appears to approach most nearly to Chætocladium.

Sub-order 3 : PIPTOCEPHALIDEÆ.—This very small group (Piptocephalis, de By., Syncephalis, Van Tiegh.) is, like the last, composed of parasites on the Mucoreæ, and to this end the mycelial hyphæ bear *haustoria*, each of which emits from its slightly swollen base a small crop of short delicate *rhizoids* traversing the Mucor-hypha affected. The conjugating hyphæ of Piptocephalis are arched somewhat like an inverted ∩, the point of contact being the summit. Actual conjugation occurs, as in the Mucoreæ ; but when this stage is reached, the product of the conjugation begins to swell at the point of union, and generally on the convex side, into a globular body, which becomes echinulate as it swells. When it has attained its full size and development at the expense of the protoplasm of the united gametes, it becomes separated from them by transverse partition, and remains seated, as it were, on the summit of the arch. Though not the morphological equivalent of the zygosperm of the Mucoreæ, but rather the offspring of the original zygosperm produced by the conjugation of the gametes, it will be most convenient to regard it as the zygosperm. The sporophores bear at their apices series or chains of spores produced by transverse partition. In Piptocephalis the sporophore is dichotomously branched at the summit, and each bifurcation bears a capitulum of chains of spores. Accessory acrospores are sometimes produced by Syncephalis.



FIG. 295.—*Piptocephalis Freseniana* de By. and Wor. Conjugation and formation of a zygosperm, *z*. Stages in the order of the numbers (× 650). (After Brefeld.)

De Bary ('Comp. Morph.,' p. 156) treats the incompletely known Dimargaris and Dispira, both of Van Tieghem, as at present doubtful Mucorini, probably near Piptocephalideæ. Another small group of genera of doubtful position is formed by Kickxella (Coem.), Martensella



FIG. 296.—*P. Freseniana*. *M*, a mycelial tube of *Mucor Mucedo*, the host of Piptocephalis. The mycele of the latter, *m*, penetrates *M* by haustoria, *h*. *Z*, zygosperm. *c*, sporophore (× 300, the rest × 630). (After Brefeld.)

(Coem.), Coemansia (Van Tiegh. and Le Mon.); while Sorokin's Zygochytrium, an aquatic saprophyte on dead insects, the account of which needs confirmation, stands in a like uncertain position.

LITERATURE.

Bainier—Observ. sur les Mucorinées et sur les zygospores des Mucorinées (Ann. Sc. Nat., 6 sér., Tom. xv., 1883).

De Bary und Woronin—Beitr. zur Morph. u. Physiol. der Pilze, i. and ii.

Brefeld—Bot. Unters. über Schimmelpilze, i. and iv.

Brefeld—Ueber Gährung, iii. (Landw. Jahrb., Thiel, v., 1876).
Coemans—Spicilège mycologique (Bull. Soc. Bot. Belg., i.).
Coemans—Quelques Hyphomycetes nouveaux (Bull. Acad. Roy. de Belgique, 2 sér., Tom. xv.).
Coemans—Recherches sur le polymorphisme et les différents appareils de réproduction chez les Mucorinées (*ibid.*, Tom. xvi.).
Coemans—Monographie du genre Pilobolus (Mém. Couronn. de l'Acad. Roy. d. Belgique, Tom. xxx.).
Cunningham—On the Occurrence of Conidial Fructification in the Mucorini, illustrated by Choanephora (Trans. Linn. Soc. Lond., 2 ser., i., 1878).
Fresenius—Beitr. zur Mycologie, i. and iii.
Gilkinet—Mém. sur le polymorphisme des Champignons (Mém. Couronn. Acad. Belg., Tom. xxvi., 1875).
Hildebrand—Ueber zwei neue Syzygites Formen (Pringsh. Jahrb., vi.).
Klein—Zur Kenntniss des Pilobolus (Pringsh. Jahrb., viii.).
Tulasne—Note sur les phénomènes de copulation, &c. (Ann. Sc. Nat., 5 sér., Tom. vi., 1866).
Van Tieghem et Le Monnier—Rech. sur les Mucorinées (Ann. Sc. Nat., 5 sér., Tom. xxvii., 1873).
Van Tieghem—Nouv. Rech. sur les Mucorinées (*ibid.*, 6 sér., Tom. i., 1875).
Van Tieghem—Troisième Mém. sur les Mucorinées (*ibid.*, 6 sér., Tom. iv., 1876).
Zimmermann—Das Genus Mucor (Chemnitz, 1871).

Order 2.—Entomophthoreæ.

This small group of parasites inhabiting the bodies of insects agrees with the Mucorini only in the formation of zygosperms. The mycele vegetates within the body of the insect attacked, and consists either of septate branching hyphæ (Entomophthora, Fres.), or of a yeast-like mass of cells (Empusa, Cohn). Zygosperms are formed, as described by Nowakowski (Entomophthora ovispora, Nowak., and E. curvispora, Nowak.), by the conjugation of adjacent hyphæ which emit corresponding lateral protuberances. These meet, become united in an H fashion (somewhat as in Spirogyra), and enter thus into open communication. As a result of this conjugation, there arises, either on the conjugating branches or near them, a globular body, which develops at the expense of the protoplasm of the united hyphæ, and finally becomes cut off by a wall. This must be regarded as a zygosperm, morphologically the equivalent of that of Piptocephalis. Azygosperms occur in E. radicans (Bref.) and certain species of Empusa, arising either as lateral outgrowths, or sometimes as terminal bodies. The cell-membrane of both zygosperms and azygosperms becomes much thickened and differentiated into a thick extine, generally of regular outline, and a thin intine, while in the contents a large fatty drop appears. Zygosperms and azygosperms both rest for a considerable period within the dead body

of the host, the surrounding mycele disappearing. On germination, which Nowakowski describes in Empusa Grylli (Fres.), a short promycele is emitted which bears a single spore.

Commonly, however, neither zygosperms nor azygosperms are formed, and after the death of the insect, spores are produced on its outer surface. In the case of Empusa—for example, E. Muscæ (Cohn), which attacks the common house-fly in large numbers in autumn—the yeast-like mycelial cells, at the time of the death of the insect, send forth each a tube, which bursts through the skin, and outside becomes a short club-shaped sporophore bearing a single acrospore. Each sporophore bears but one spore, and then perishes. The spores are capable of germination at once, but the power lasts only for a few days. Affected flies in this condition are common enough objects attached to windows, &c., and surrounded by a whitish mass of spores. The mycele of Entomophthora, which is septate, much branched, and often anastomosing, sends branches through the skin to the outer surface, where farther ramification takes place, investing the body of the insect. These branches range themselves at right angles to the insect's body, and terminate together at nearly the same elevation. Each such branch is a sporophore, which, as in Empusa, forms a single acrospore. The spores are capable of germination at once like those of Empusa; and in both genera either a very short tube is formed, bearing a secondary spore, as in the promycele of the zygosperm, which, on germination, may attack a fresh insect, or the germ-tube of the primary spore may do so without the intervention of secondary spores.

Completoria complens (Lohde), found by Leitgeb in fern prothallia, and Conidiobolus utriculosus (Bref.), described by Brefeld as a parasite on Tremellini, are two forms placed here which, unlike the rest of the group, do not attack insects. Brefeld, who has investigated the group minutely, does not accept the conjugation as a real one, and brings forward arguments against it based on the anastomosing of hyphæ and the situation of the zygosperms. His opinion, if accepted, would lead to placing the group elsewhere; but de Bary states ('Comparative Morph.,' p. 159) that Nowakowski's and Brefeld's different observations may be explained by the different behaviour of different species.

Literature.

Brefeld—Untersuch. über die Entwickel. der Empusa Muscæ und Empusa radicans und die durch sie verursachten Epidemien der Stubenfliegen und Raupen (Abhandl. d. Naturforsch. Gesellsch. zu Halle, xii.).

Brefeld—Ueber Entomophthoreen und ihre Verwandten (Sitzungsber. d. Gesellsch. Naturforsch. Freunde zu Berlin, 1877). See also Bot. Zeit., 1877, p. 345.

Brefeld—Bot. Unters. über Schimmelpilze (Ent. radicans), iv., 1881, p. 97.

Cohn—Empusa Muscæ und die Krankheit der Stubenfliege (Nova Acta, xxv., p. 1).
Cohn—Ueber eine neue Pilzkrankheit der Erdraupen (Tarichium megaspermum), (Beiträge zur Biologie der Pflanzen, Bd. i., Heft 1, p. 58).
Eidam—Eine auf Excrementen von Fröschen gefundene Entomophthoree (Bot. Centralblatt, xxiv., 1885).
Fresenius—Ueber die Pilzgattung Entomophthora (Abhandl. d. Senkenberg. Ges., Bd. ii.).
Giard—Deux espèces d'Entomophthora, &c. (Bull. Sc. du Départ. du Nord, 2 sér., 2 Ann., No. 11).
Frey und Lebert—Die Pilzkrankheit der Fliegen (Verhandl. d. Naturf. Ges. zu Zürich, 1856).
Leitgeb—Completoria complens, ein in Farnprothallien schmarotzende Pilz (Sitzungsber. d. Wien. Acad., Bd. 84, Abth. 1).
Nowakowski—Die Copulation bei einigen Entomophthoreen (Bot. Zeit., 1877).
Nowakowski—Entomophthoreæ (Abhandl. d. Acad. d. Wiss. zu Krakau, 1883). Polish, see Bot. Zeit., 1882.
Sorokin—Zwei neue Entomophthora Arten (Cohn's Beiträge zur Biol. d. Pflanzen, Bd. ii., Heft 3, p. 387).

Order 3.—Chytridiaceæ.

The Chytridiaceæ are a group of minute, more or less aquatic, parasitic fungi, embracing forms which may, in the present state of our knowledge of them, be thus classed together; but whether it will eventually appear that these are naturally related to each other, or are merely organisms of different affinities presenting a similar appearance owing to similar environment and ways of life, is but a subject for speculation. However, there are points in which all agree, and their life-history may be briefly summarised thus:—Zoospores—mostly uniciliated (the rest with two cilia), and containing generally a drop of fatty substance, and, in the larger forms at least, a nucleus—are produced in zoosporanges of various forms and sizes. These escape from the apex of the zoosporange, which is provided in some cases with a lid, either successively or in a mass held together by a viscous substance, from which they are gradually set free. An undulating alteration of outline, accompanied by amœboid movement, takes place in the zoospores of certain species towards the end of the period of their activity. The zoospores give rise again to zoosporanges. Resting-spores are known in certain cases, which likewise give rise to zoosporanges; while in the Rhizidieæ a process probably intermediate between oogamous reproduction and isogamous conjugation takes place. Of the four sub-orders, the first (*Rhizidieæ*) is manifestly nearly related to the Mucorini and the Ancylisteæ; the second (*Cladochytrieæ*) may be regarded as allied to the Rhizidieæ; the third (*Olpidieæ*) and the fourth (*Synchytrieæ*) in all probability following the second. De Bary suggests ('Comp. Morph.,' p. 169) that in

this order the whole group may be looked upon as a lateral branch of the Mucorini or Ancylisteæ successively modified (degraded) by aquatic parasitism, with its extremity represented by the Synchytrieæ, Woronina (Cornu), and Rozella (Cornu). De Bary also discusses (*loc. cit.*) the suggested relationship to such algæ as Protococcaceæ, Characium, Chlorochytrium, &c. Apart from the possession of chlorophyll, the conjugation of zoospores in these algæ separates them from the Chytridiaceæ, in which group such a process has not (at least as yet) been discovered. Granting a relationship of the simpler Chytridiaceæ with Protococcaceæ, &c., these might be regarded as leading up to Rhizidieæ, Ancylisteæ, and Mucorini; unless one regards the Chytridiaceæ as composed of two distinct sub-groups, Rhizidieæ and Cladochytrieæ, related to Mucorini and Ancylisteæ; and the Olpidieæ and Synchytrieæ to Protococcaceæ, &c. Only further research may determine these questions of affinity.

Sub-order 1: RHIZIDIEÆ. The life-history of Polyphagus Euglenæ



FIG. 297.—*Polyphagus Euglenæ* Now. *A*, zoospore with drop of fatty matter and nucleus. *B*, young plant attached to resting Euglena, *e*. *C*, zoosporange containing spores resting on empty prozoosporange, *a*. *D*, conjugation: *a*, the receptive individual; *b*, the supplying individual; *s*, the swollen end of conjugation tube (rudiment of resting-spore); *e*, *e*, *e*, the Euglenæ. *E*, portion of *D* five and a half hours later: *b*, empty, *s*, mature; represent the same parts as in *D*. (*A* × 550, *B*, *D*, *E*, × 350, *C* × about 400.) (After Nowakowski.)

(Nowak.), described by Nowakowski, furnishes us with the most highly developed type of the whole group. The mycele consists of very slender branching rhizoids, tapering each to a very thin point, and attaching themselves to the Euglena-hosts. The original germinating zoospore

from which these arose remains, as it were, the centre of the system of rhizoids, and, nourished by them, grows considerably in size. At length, when it has attained full development, it becomes a *prozoosporange*, since from it there grows out a thick, cylindrical, thin-walled process, into which all the protoplasm passes, and within which it breaks up into zoospores. These, escaping, repeat the life-history. Zygosperms (or oosperms) are produced by the conjugation of gametes, which play unequal parts in the process. The one (supplying) individual is round and larger than the other (the receptive) individual. A rhizoid from the receptive individual places its apex in contact with the supplying individual itself, and begins to grow in girth. The cell-wall at the place of contact disappears, the protoplasm of both unites and passes into a swelling which has arisen on the conjugating tube of the receiving individual close behind the place of contact. This swelling then becomes the zygosperm, which is provided with a thick wall, sometimes covered with fine spines, appearing as early as the outset of the swelling process. It happens—though rarely—that the tubes of two or three receptive individuals attach themselves to one supplying plant, and a corresponding number of zygosperms is thus formed. After a period of rest, the zygosperm germinates by producing a zoosporange. Generation after generation of zoosporanges intervene between zygosperm and zygosperm. 'Which of the two should be called the male and which the female, is not easy to determine. . . . It is evident that we have before us an intermediate case between the ordinary forms of oogamous and isogamous conjugation.' (De Bary, *loc. cit.*, p. 163.)

A series of incompletely-known forms may be placed beside Polyphagus (Nowak.), viz.:—Physoderma (Wallr.) (*pro parte*), Rhizidium (A. Br.), Rhizophydium (Schenk), Obelidium (Nowak.), Chytridium (A. Br.), Phlyctidium (A. Br.). Resting-cells of some of these have been found, but their genesis is unknown.

Sub-order 2 : Cladochytrieæ.—This is a small sub-order, the members of which mostly inhabit the tissues of marsh plants, and possess copiously branching mycelial rhizoids, bearing terminal and interstitial zoosporanges. The zoospores give rise on germination to a mycele like the parent one. Resting zoosporanges occur. No process of conjugation is known either between filaments or zoospores. Cladochytrium (Nowak.) and Physoderma (*pro parte*) compose the sub-order.

Sub-order 3 : Olpidieæ.—The Olpidieæ are wholly destitute of a mycele, and the life-history, as described by Fischer for Olpidiopsis Saprolegniæ (Fisch.), and O. fusiformis (Cornu), which inhabits species of Achlya, is a very simple one. The zoospores of O. Saprolegniæ perforate the young mycelial hyphæ of Saprolegnia, and, after a few days of

growth in amœboid fashion, a cell-wall is secreted, and each becomes a zoosporange. Swellings arise on the mycele of the host as the result of this parasitism. The zoosporange then sends forth a cylindrical process, which perforates the wall of the Saprolegnia, and through it the zoospores are discharged. The zoosporanges are either smooth and capable of emission of zoospores at once, or are covered with fine spines and capable either of emission at once or of resting. While the latter are generally formed under adverse circumstances, it seems to occur with some regularity that the zoospores of the smooth zoosporanges give rise to spiny zoosporanges, and *vice versâ*. The life-history of these two species may be taken as typical, and the incompletely-known species of Olpidium (A. Br.) doubtless conform to it.

Sub-order 4 : Synchytrieæ.—The Synchytrieæ inhabit the epiderm of terrestrial Flowering Plants, in which they excite the production of small yellow or dark-red galls, owing to the abnormal swelling of the epidermal cells affected. Like the Olpidieæ, they have no mycele, but they are distinguished from that sub-order by the formation of a *sorus* of zoosporanges. From the germinating zoospore an initial cell is formed, the contents of which break up into a sorus of zoosporanges. In Pycnochytrium (de By.) (= Chrysochytrium, Schroet., and Leucochytrium, Schroet.) the initial cell is a resting-cell, which eventually germinates by the gradual protrusion of its contents into a globular sac seated upon the extine. Within this sac the sorus of zoosporanges is formed by the division of the protoplasm. Each zoosporange produces a considerable number of zoospores, which again give rise to resting-cells. In Eusynchytrium (Schroet.) an indefinite number of sorus-forming generations, which at once produce zoospores, intervene between resting-cell and resting-cell ; while in Synchytrium Taraxaci (de By.) the resting-cell produces a zoosporange without the intervention of a sorus, a process suggestive of the Olpidieæ. No conjugation of zoospores nor any sexual process has been observed in any member of the group.

Woronina and Rozella, which inhabit Saprolegnieæ, may be placed with Synchytrieæ.

De Bary (*loc. cit.*, p. 170) treats as doubtful Chytridiaceæ (1) Tetrachytrium triceps (Sorok.), the zoospores of which are said to conjugate; and (2) Hapalocystis mirabilis (Sorok.), the zoospores of which are described as conjugating within the mother-cell. The observations, however, require confirmation.

Beyond the general reference to Professor de Bary's 'Comparative Morphology,' &c., the student is specially referred to that source with regard to this incompletely-known group.

LITERATURE.

De Bary—Beitr. zur Morph. u. Physiol. der Pilze, i. (Physoderma [Cladochytrium]).
De Bary und Woronin—Beitr. zur Kenntniss der Chytridieen (Ber. Naturf. Ges. Freiburg, Bd. iii.; and Ann. Sc. Nat., 5 sér., Tom. iii.).
Woronin—Neuer Beitr. zur Kenntniss der Chytridieen (Synchytrium Mercurialis) (Bot. Zeit., 1868, p. 81).
Woronin—Chytridium Brassicæ (in his paper on Plasmodiophora, Pringsheim's Jahrb. wiss. Bot., Bd. xi.).
Braun—Ueber Chytridium, &c. (Monber. und Abhandl. Berl. Acad., 1855).
Braun—Ueber einige neue Arten der Gattung Chytridium und die damit verwandte Gattung Rhizidium (Monber. Berl. Acad., 1856).
Cienkowski—Rhizidium Confervæ glomeratæ (Bot. Zeit., 1857).
Cohn—Ueber Chytridium, &c. (Nova Acta Leop.-Carol., xxiv., Pt. i., p. 142).
Cornu—Chytridinées parasites des Saprolegniées (in Monogr. d. Saproleg., Ann. Sc. Nat., 5 sér., Tom. xv., p. 112).
Fisch—Ueber zwei neue Chytridiaceen (Sitzber. d. phys. med. Soc. zu Erlangen, 1884).
Fischer—Ueber d. Stachelkugeln in Saprolegniaschlauchen (Olpidiopsis) (Bot. Zeit., 1880).
Fischer—Untersuch. über die Parasiten der Saprolegnieen (Pringsh. Jahrb., Bd. xiii.). (Berlin Habilitationschrift, 1882).
Fischer—Zur Kenntniss d. Chytridiaceen (Erlangen, 1884).
Kny—Ueber Entwickelung des Chytridium Olla (Sitzber. d. Berl. Naturf. Freunde. See also Bot. Zeit., 1871, p. 870).
Nowakowski—Beiträge zur Kenntniss d. Chytridiaceen (Cohn's Beitr. zur Biol. d. Pflanzen, ii.).
Nowakowski—Polyphagus Euglenæ (*ibid.*, ii.).
Schenk—Algol. Mittheilungen (Verhandl. d. Phys. Med. Ges. zu Würzburg, Bd. viii.).
Schenk—Ueber d. Vorkommen contractiler Zellen im Pflanzenreiche (Würzburg, 1858).
Schroeter—Die Pflanzenparasiten aus der Gattung Synchytrium (Cohn's Beitr. zur Biol. d. Pflanzen, i.).
Schroeter—Untersuch. über die Pilzgattung Physoderma (Ber. d. Schlesischen Ges., 1882).
Sorokin—Einige neue Wasserpilze (Tetrachytrium triceps) (Bot. Zeit., 1874).
Sorokin—Uebersicht d. Gruppe Syphomycetes (Hapalocystis mirabilis) (Arbeiten d. Naturf. Ges. an der Univ. Kazan, 1874, Bd. iv.). (See Just's Jahresbericht, 1875.)
Thomas—Synchytrium cupulatum (Bot. Centralblatt, xxix.).

ORDER 4.—PROTOMYCETACEÆ.

Protomyces macrosporus, Ung., to which there have recently been added a considerable number of species, many of them on insufficient grounds, is a parasite on Umbellifers, especially Ægopodium Podagraria, Meum athamanticum, and more rarely Heracleum sphondylium, inhabiting the intercellular spaces of the leaf, petiole, stem, flower-stalk, and pericarp. It possesses a branching septate mycele, at irregular intervals on which there are formed interstitially large somewhat oval

resting *progametanges*, with a stratified membrane and dense contents. These persist when the mycele dies, and hibernate. After this period of rest, and on liberation from the decaying tissues of the host (in water), the intine, with the contents, bursts the extine, and, becoming free, constitutes the *gametange*. Within it a large number of minute short rod-shaped *gametes* are formed, while a portion of the protoplasm remains unused in the process. These gametes are ejected, and, being without the power of spontaneous movement, they remain

FIG. 298.—*Protomyces macrosporus* Unger. *a*, a mature resting progametange; *b*, gametange; *c*, *d*, and *e*, further stages of the same in the development of gametes. *c* shows the parietal protoplasm, *d* the same divided into gametes, *e* the gametes rounded off and separated from the rest of the parietal protoplasmic layer (× 390). (After de Bary.)

in more or less proximity to each other. Where pairs of gametes come together, they emit fine processes which conjugate, the whole having the appearance of a dumb-bell or of the letter H. The germination of these has been observed on the epiderm of Ægopodium; it takes place by the emission from one of the original gametes of a germ-tube, to the nourishment of which the protoplasm of the united gametes contributes. This germ-tube, on entering the tissues of the host, repeats the life-history.

LITERATURE.

De Bary—Beitr. zur Morph. und Phys. der Pilze, i.

Von Thümen—Eine neue Protomyces Species (Hedwigia, 1874).

Wolff—See footnote to de Bary's paper 'Protomyces microsporus und seine Verwandten' (Bot. Zeit., 1874, p. 82).

Other literature of systematic interest in Saccardo's Sylloge.

ORDER 5.—USTILAGINEÆ.

The parasites which are grouped together under this name affect Flowering Plants of different natural orders, but are especially conspi-

cuous as causing diseases of grasses. As a rule the attack of the parasite is limited to one special region of the host, e.g. the ovary, or the whole flower, or the leaf, or the stem, or even, in a few cases, the root; and when the fungus has attained its maturity, the result commonly is that the part affected has been destroyed with the exception of the epiderm or integument and the remains of vascular tissue, and is replaced by a powdery mass of brown or black resting-spores. The mycele sends its long thin hyphæ mostly along the intercellular spaces, in many cases emitting branched haustoria into the adjoining cells, and the resting-spores are formed either on all parts of the hyphæ or on particular branches.

FIG. 299. *Tilletia caries* Tul., germinating. In *a*, gametes on promycele, *p*. In *b*, gametes, *s*, conjugate in pairs. In *c*, a germ-tube proceeding from pair of gametes, *s*. *s'*, a sporid (× 460). (After Tulasne.)

The life-history, briefly stated, begins with the production from the resting-spore of a *promycele* which bears sporid-like *gametes*; these gametes conjugate in pairs, and the united pair either directly produce a new mycele, or *sporids* which do so. This mycele then bears *resting-spores* again in another host. Variations on this course of life-history will be mentioned later on. Though these conjugating gametes differ from those of Protomyces in their acrogenous origin on a promycele, they may yet be considered homologues of those, just as the acrogenous spores of Chætocladium are undoubtedly homologous with the endogenous spores of Mucor.

In Entyloma (de By.) the resting-spores are borne interstitially at indefinite intervals on the mycele, as in Protomyces; in Tilletia (Tul.) they occur singly, and only terminally on the spore-bearing hyphæ; while in Geminella (Schroet.) they are borne, two together, in series throughout the length of the special hyphæ. In Urocystis (Rabenh.), Sorosporium (Rudolphi), and Tuburcinia (Berk.) they are united, several together, into a kind of coil, which is invested with a transitory or a persistent integument. With the exception, perhaps, of Graphiola (Port.), the exact relationship of which to the Ustilagineæ has yet to be determined, Sphacelotheca Hydropiperis (de By.), formerly Ustilago Hydropiperis, according to de Bary's description, affords the best example of a well-developed *stroma*. This fungus attacks the ovule of

Polygonum Hydropiper, and replaces it by a dense plexus of hyphæ, which exhibits a differentiation of a thick outer wall enclosing the whole, a cylindrical columel, both colourless, and a dark-violet mass of resting-spores filling up the space between. An undifferentiated basal portion discharges the function, as it were, of a meristem, and adds new elements to wall, columel, and spore-mass. The fungus attacks only the ovule ; and as it grows thus in bulk, the unaffected wall of the ovary is burst, and exposes the brittle wall of the sporal mass, which breaks at the slightest touch, and sets the resting-spores free.

The germination of the resting-spores, which are usually round or many-sided, and possessed of both intine and extine, the latter often covered with characteristic fine granulations, varies according as it takes place in water merely or in a nutritive fluid. In water only the resting-spores of Entyloma, Tilletia, Tuburcinia, and Urocystis produce a short promycele-tube, which bears simultaneously on its blunt apex a whorl, or 'crown' of elongated gametes, varying in number. Many species of Ustilago (Pers.) and Tolyposporium (Woron.) produce a promycele-tube which divides into a number of cells, from each of which bud off laterally rod-shaped sporids or gametes, somewhat as a yeast-cell buds off. The resting-spores of Thecaphora Lathyri (Kuhn) and Ustilago longissima (Tul.) give rise to a short slender promycele-tube, which bears an acrogenous sporid, or several serially ; while in Thecaphora hyalina (Fingerh.), Ustilago carbo (Tul.), and U. destruens (Tul.) it divides into several cells, each of which produces, if any, only one sporid.

The conjugation of gametes in pairs takes place either before or after separation from the promycele, by means of short or long cross hyphæ, according to the distance between them. The product of this union is the outgrowth from the connected gametes of a mycelial germ-tube, or, as occurs in Tilletia, Entyloma, Tuburcinia, and Urocystis, sporids are produced on the outgrowth from the connected pair, and these then emit mycelial germ-tubes. In a number of forms gametes are constantly produced which do not conjugate, but give rise to mycelial germ-tubes, as sporids do, while in Sorosporium Saponariæ (Rudolph.) the resting-spore itself emits the mycelial germ-tube directly, without the intervention of promycele, gametes, and sporids. What occurs regularly in these species may occur exceptionally in the case of individuals the gametes of which ordinarily conjugate ; that is, they may fail to produce gametes or sporids at all, or their gametes may fail to conjugate, either all of them, or, where the number of gametes is an uneven one, the odd gamete. In such exceptional instances the

resting-spore or the single gamete, as the case may be, produces a mycelial germ-tube, as a sporid does.

When the resting-spores germinate in nutritive fluids, as the experiments of Brefeld show, the product is, according to the species, either directly a mycele which bears spores, or an abundant yeast-like outgrowth from the promycele. In cases where the resting-spores failed to germinate, they were permitted to do so in water, and the sporids so formed, on being introduced into a nutritive fluid, gave rise to myceles bearing acrospores, resembling, as the case might be, either the gametes or the sporids.

Accessory *acrospores* resembling the sporids are borne on branches of the mycele of certain species of Entyloma before the production of resting-spores. The branches of the mycele which bear them protrude through the stomates and epiderm of the host. They probably give rise to new myceles, as the sporids do. The same thing occurs in Tuburcinia Trientalis (Berk.), only here the accessory spores differ in form from the promycelial sporids. They produce in this case undoubtedly each a mycele which bears resting-spores.

Brefeld regards the conjugation of the gametes as a merely vegetative process, and in no way analogous to any sexual act. De Bary ('Comp. Morph.,' p. 182) subjects Brefeld's arguments to destructive criticism, while stating Brefeld's case in the fairest terms. While the student is referred to the source quoted for the details, it may be shortly stated that de Bary's arguments are broadly based on the regularity with which conjugation occurs, and on the fact that it takes place equally regularly in pairs, under the normal conditions of germination in water. He shows, besides, conclusively how the process differs from the well-known vegetative anastomosing of hyphæ, &c.

De Bary regards the higher forms of Ustilagineæ, such as the coil-forming Urocystis, Sorosporium, and Tuburcinia, and Sphacelotheca with its well-developed stroma, as connected through the simpler forms (e.g. Entyloma) with Protomyces. Both produce (in the one case acrogenous, in the other endogenous) conjugating cells of equal value. The next indicated ally is Cladochytrium, the spores of which are zoospores (a matter not affecting the homology), but these fail to conjugate so far as is known. At all events the nearest ally appears to be, in the present state of our knowledge, that group of Chytridiaceæ to which Cladochytrium belongs, through Entyloma and Protomyces.

The peculiarity is to be noted that in Protomyces and the Ustilagineæ the act of conjugation takes place at a stage of the life-history which bears no homology with the sexual states of other Phycomycetes.

LITERATURE.

De Bary—Untersuch. über die Brandpilze (Berlin, 1853).
De Bary—Protomyces microsporus und seine Verwandten (Bot. Zeit., 1874) (Entyloma).
Brefeld—Botanische Untersuchungen, v. (Leipzig, 1883).
Cornu—Sur quelques Ustilaginées nouvelles (Ann. Sc. Nat., 6 sér., Tom. xv.).
Fischer von Waldheim—Beitr. zur Biologie und Entwickelungsgesch. der Ustilagineen (Pringsh. Jahrb., Bd. vii.).
Fischer von Waldheim—Les Ustilaginées et leurs plantes nourricières (Ann. Sc. Nat., 6 sér., Tom. iv.).
Ed. Fischer—Beitr. zur Kenntniss der Gattung Graphiola (Bot. Zeit., 1883).
Kuhn—Die Krankheiten der Culturgewächse (Berlin, 1858).
Prillieux—Quelques observ. sur la formation et la germination des spores des Urocystis (Ann. Sc. Nat., 6 sér., Tom. x.).
Schroeter—Bemerk. u. Beobacht. über einige Ustilagineen (Cohn's Beiträge zur Biologie der Pflanzen, Bd. ii.).
Tulasne—Mém. sur les Ustilaginées comparées aux Uredinées (Ann. Sc. Nat., 3 sér., Tom. vii.).
Tulasne—Second Mém. sur les Uredinées et les Ustilaginées (*ibid.*, 4 sér., Tom. ii.).
Ward—Entyloma Ranunculi, Proc. Roy. Soc., xli. (1886).
Winter—Einige Notizen über d. Fam. d. Ustilagineen (Flora, 1876).
Wolff—Beitr. zur Kenntniss der Ustilagineen (Bot. Zeit., 1873).
Wolff—Der Brand des Getreides (Halle, 1874).
Woronin—Beitr. zur Kenntniss der Ustilagineen (Tuburcinia). (De Bary und Woronin's Beitr. zur Morph. u. Phys. d. Pilze, v.).

The older literature will be found quoted in Tulasne, Fischer von Waldheim, and in de Bary's 'Brandpilze.'

GROUP II.—SPOROCARPEÆ.

Class XX—Ascomycetes.

THE SPOROCARP.

This large class is distinguished by the universal formation of spores in *asci*, for the most part tubular in shape, but sometimes broadly ovate or roundish, and borne terminally on special hyphæ termed *ascogenous hyphæ*. When an ascus has reached its full size, and only then, the formation takes place within it of *ascospores* by free-cell formation. The young ascus is at first filled with finely granular protoplasm, which contains a nucleus, and within it a smaller body, probably a nucleole. The protoplasm next gathers itself together at the upper part of the ascus, while a watery fluid occupies the remainder of the cavity except a thin

layer of protoplasm coating the wall. At this stage the full growth of the ascus is commonly reached, and the formation of ascospores begins with the division of the nucleus into two ; then by the same process four appear, then eight, which in the majority of cases represents the number of ascospores. In many Ascomycetes, however, other numbers typically prevail, e.g. one, or two, or four, or sixteen, forty, fifty, and so on to over a hundred. Dothidea (Fr.), for example, has two to four, Sordaria (Ces. and De Not.) four, sixteen, sixty-four, and one hundred and twenty-eight. Whatever the number, the nuclei always possess the same



FIG. 300.—*Peziza* (*Pyronema*) *confluens* P. *a*, small portion of hymenium ; *p*, paraphyse attached to, not originating in, hyphal branches from which the three asci spring ; *m*, young asci ; *r*—*w*, successive stages, according to letters, in the development of ascospores within asci (× 390). (After de Bary.)

structure in all stages of multiplication, but they become smaller in size as the number advances. Round each nucleus there gathers a clear mass of protoplasm, and ultimately this becomes enclosed by a membrane, and, growing in size, thus develops into a spore. The ascospores are arranged in a series, one over the other within the ascus. The protoplasm left over within the ascus and outside the spores differs from that within the spores in exhibiting a reddish or violet-brown colour after treatment with iodine solution. De Bary originally proposed the term 'epiplasm' for this portion of the contents ; but, Errera having

shown that it contains a relatively large proportion of glycogen, he has more lately ('Comp. Morph.,' p. 77) adopted for it the term glycogen-mass or simply the *glycogen*. In certain cases the separation of the glycogen from the protoplasm takes place before the formation of spores, the former occupying the lower part of the ascus, and in some instances both the apical and the basal portions on either side of the protoplasm in which the nucleus is situated and the ascospores are formed.

Such are the asci which characterise the great class of Ascomycetes. They are borne as a rule in considerable numbers mostly between hair-like bodies, the *paraphyses*, and united into *hymenia* within special *sporocarps*. The sporocarps are either seated on a fine mycele or they are borne on a large *stroma* (thallus), which assumes in different genera diverse characteristic forms. The whole body of the Ascomycetous type is always built up of septate hyphæ.

The *sporocarps* are composed of two main elements—of the asci with the ascogenous hyphæ, and of the *envelope* including paraphyses. These two elements, though much interwoven with each other, are of distinct origin. The paraphyses, for example, are never borne by the ascogenous hyphæ, and the asci never by the cells of the envelope; at all events the one thing certain about so-called exceptions is their doubtful character. According to the structure of the mature sporocarp, the Ascomycetes may be classed in three divisions: the *Discomycetes*, with *discocarps* or *apotheces* (hymenia exposed); the *Pyrenomycetes*, with *pyrenocarps* or *peritheces* (hymenia within flask-shaped bodies open at the neck); and a third assemblage of forms with *cleistocarps* (enclosed hymenia).

The hymenial disc of the *apothece*, consisting of asci and paraphyses placed perpendicularly to the surface, is when ripe fully exposed. The asci are imbedded, so to speak, in the paraphyses, and the whole mass terminates in a regular surface. The paraphyses originate beneath from a dense plexus of hyphæ called the *subhymenial layer* or *hypothece*, directly continuous with the outer envelope termed the *exciple* or with the tissue of the stroma as the case may be. The paraphyses thus form a part of the envelope-tissue as distinguished from the ascus-apparatus. The ascogenous hyphæ are interwoven with those hyphæ of the hypothece which bear the paraphyses. Terminating upwards in the asci they are barely otherwise distinguishable in the mature apothece, unless in certain cases by their greater size and the blue colour they assume after treatment with potash. The asci grow up between the paraphyses, and reach the surface, as has already been indicated, about the period of ripening. As a rule the apothece exhibits a regular progressive marginal growth, with, in a considerable number of cases, an intercalary

growth in addition, giving rise to irregularities of shape. The basin-shaped apotheces of Peziza (Discomycetes) and of the gymnocarpous lichens are the characteristic forms, while with them may be placed the large stromata of Morchella (Dill.) and allied genera, which are club-shaped and in the form of stalked caps. The Hysterineæ and the Phacidiaceæ, Ascobolus (Pers.), Pyronema (Fckl.), &c., which show no marginal extension, approach the Pyrenomycetes in this respect.

The *perithece* does not differ essentially from the apothece. As a rule it is much smaller, rarely more than one millim. in diameter, and it consists of an outer wall enclosing an ascogenous hymenium at all points but one narrow opening, the *ostiole*. The whole perithece is round or flask-shaped, and the ostiole is simply a pore in the wall or a channel through the neck of the flask. In other words, while an apothece is generally larger, it does not differ in any marked way from a perithece, except that its hymenium has an open surface, while the margin of the perithece is arched over, leaving only a narrow opening for the escape of the spores. The asci arise from ascogenous hyphæ, either exclusively at the base of the perithecial cavity or at all points of the inner surface, and in successive crops, producing in all a considerable number of ascospores. The envelope consists of a wall which, when a stroma is absent and the peritheces appear singly on the mycele, is composed of a dense pseudo-parenchyme clothed sometimes externally with hairs. Where the peritheces are borne on a stroma there is no sharp differentiation of wall-structure. The neck, when present, is but a prolongation of the wall, and the ostiole arises (for at first the cavity is fully closed) as an intercellular passage either schizogenously by the separation of tissue through unequal growth, or lysigenously by the absorption of the tract of tissue originally occupying its place. In many cases it is hard to say by which process it arises, and it is likely enough that both sometimes have a share in it. One distinguishes here between *paraphyses* which arise and stand in relation to the asci as in the apothece, and *periphyses*, by which is meant other hairs of like origin arising from the hymenium at places where there are no asci—for example, in the region of the neck. Sometimes the periphyses protrude through the ostiole. They are always present in greater or less numbers, except, according to Füisting, in Massaria (De Not.); while paraphyses are absent from a number of genera, both of fungi and of lichens. In Chætomium fimeti (Fckl.) the perithece remains closed, and this transition form leads us to the division which possesses cleistocarps.

Just as the peritheces are essentially folded over apotheces, so the *cleistocarps* may be described in general terms as peritheces which

remain wholly closed. The spores escape from them by the decay or the rupture of the wall by external agencies. The Tuberaceæ, which possess undoubted asci of peculiar form, are necessarily included among the Ascomycetes, and, since little more is known of them than the structure of the sporocarps, they may be most fittingly dealt with here at the conclusion of the consideration of the mature sporocarp.

The formation of ascospores within the asci of *Tuberaceæ* (truffle family) differs from the typical mode already described. In Tuber (Mich.) the asci are globular, and a differentiation of glycogen from protoplasm takes place, the glycogen forming as it were a layer next the wall. In the protoplasm cavity, which is separated from the glycogen by a sharply outlined

FIG. 301.—*Tuber rufum* Pico. *a*, section seen in reflected light. The white veins, *l*, contain air, the dark ones, *v*, fluid, *h*, hymenial tissue. *b*, a thinner section seen in transmitted light, lettering as in *a*, white and dark appearance of veins reversed. (*a* × 5, *b* × 15.) (After de Bary.)

layer, the nuclei (4–6 in number) are formed, and young ascospores appear—one or more only, however, developing, while the rest are arrested and ultimately disappear. In Elaphomyces (Nees ab Esenb.) the process is somewhat similar, though here no glycogen is differentiated.

The sporocarps of the Tuberaceæ, which are mostly large subterranean bodies, are borne on a fine mycele, usually vanishing at the time of maturity. They are either attached to the mycele only at their bases or are completely invested by it. Very little is known as to the life-history of any of the forms, but the structure of the sporocarps has been accurately and exhaustively detailed. These are enclosed by a *peridium*, the outer surface of which is frequently corrugated irregularly (Tuber) or regularly (Hydnobolites, Tul., Genabea, Tul.), sometimes covered

with wart-like excrescences, or quite smooth. This peridium consists of a pseudo-parenchyme of densely compacted hyphæ, showing, in the case of Stephensia (Tul.), separate layers. The external cells are denser and thicker-walled than those within, but a gradual transition takes place from one to the other, and again from the latter to the cells composing the veins or seams of tissue which traverse the interior and divide the spore-bearing tissue into chambers. The walls of such chambers in Genabea (Tul.) possess a pseudo-parenchymatous structure like the peridium, but in general such walls consist of a looser tissue. Only by the perishing of this peridium or its rupture by external agencies do the spores escape.

Certain authors divide the Tuberaceæ into two families, the *Tubereæ*, containing Tuber, with an assemblage of allied genera mentioned below; and the *Elaphomyceæ*, containing Elaphomyces, Onygena (Pers.), and Penicillium (Link). Though the sporocarps of Elaphomyces and Penicillium undoubtedly resemble each other, there is at all events as yet hardly sufficient ground for going so far as to assume for Elaphomyces (or Onygena) the life-history of Penicillium. Onygena again, though it may be placed beside Elaphomyces, is hardly near enough morphologically to deserve other than a provisional juxtaposition.

Of the *Tubereæ*, Hydnobolites is the simplest, with asci distributed through the internal tissue, and the peridium represented only by the outer layer of sterile hyphæ. Genabea, with a regularly corrugated peridium and asci arranged in groups imbedded in tracts of sterile hyphæ, forms another type with Terfezia (Tul.), in which the large masses of fertile hyphæ are separated by branching white seams of sterile tissue. These sterile tracts or seams proceed inwards from the thick peridium. There is a third type, characterised by a stout peridium enclosing an internal mass consisting of thick plates of tissue, which proceed from the peridium, and are separated by many narrow channels or chambers. The wall of these chambers bears the hymenium. This type includes the remaining genera Tuber (truffles), Balsamia (Vitt.), Choiromyces (Vitt.), Pachyphlœus (Tul.), Stephensia, Hydnocystis (Tul.), Hydnotria (B. and Br.), and Genea (Vitt.).

Elaphomyces has a thick, hard, corky peridium consisting of two distinct but united layers. The external one is thin, and either smooth or warty; the inner thicker, and composed of a dense plexus of thick-walled hyphæ. When ripe the internal substance consists of a dark mass of spores traversed by a delicate cobweb-like *capillitium* which arises from the inner surface of the peridium. Before maturity the ascogenous hyphæ may be recognised by their greater diameter than the capillitium-hyphæ, but they vanish with the ripening of the spores.

According to Boudier (Bull. Soc. Bot. France, xxiii., 1876), Elaphomyces is probably parasitic on roots; but this is not certainly established.

In *Onygena*, which inhabits animal remains, the peridium is stalked, and the ascospores (eight in each ascus) become free through the disappearance of the ascus when ripe, and ultimately escape on the rupture of the peridium.

Origin of the Sporocarp.

The following types, selected by de Bary ('Comp. Morph.,' p. 197) as illustrating the origin of the sporocarp of the Ascomycetes, show an amount of variation in this process which, it may be anticipated, will be extended with farther research into the subject, while certain gaps between these types may be filled up. Taking the most simple instance of the origin of such a sporocarp, that of *Eremascus albus* Eid. described by Eidam, the sexual act preceding the formation of the sporocarp is manifestly the conjugation of two sexual elements of identical structure. Two cells of the septate mycele send forth, close by the septum which parts them, each a short lateral tube, the two tubes being not only exactly alike but in close contact from the outset. They continue to grow outward, winding round each other in a spiral course, performing three or four revolutions apiece. At this stage a septum grows across the base of each tube, and the apices conjugate. As a result of this act, a globular swelling arises at the apex, the protoplasm of the tubes is withdrawn into it and enclosed by transverse walls growing across the tops of the tubes. Within this body (an ascus), which so strikingly resembles a simple zygosperm, there are subsequently formed eight ascospores. [The *carpogone* itself becomes an ascus here.]



Fig. 302.—*Eremascus albus* Eidam. Fertilisation and formation of sporocarp. Stages in order of letters. In *f* the ascospores are formed (× 900). (After Eidam.)

The next type embraces such forms as the Erysipheæ, Eurotium, Penicillium, Sordaria, Melanospora from among cleistocarpous and pyrenocarpous forms; and Gymnoascus, Pyronema, Ascobolus from gymnocarpous and discocarpous fungi, and the Collemaceæ (discocarpous)

from the lichens. It is characterised by the production on the mycele or the thallus-hyphæ of a body consisting sometimes of a cell, sometimes of a chain of cells, called the *carpogone*, or with reference to its function the *ascogone*, since from it there arise the *ascogenous hyphæ*. (In Podosphæra, a genus of Erysipheæ, a single stalked ascus only is formed.) In this type the sexual act which precedes the farther development of the carpogone is distinctly of a higher character. In such forms as Pyronema and Eurotium an antheridial filament is produced which fertilises the *trichogyne*, as the special conjugating portion of the carpogone is called; and in the Collemaceæ detached male cells, *pollinoids*, formed in special *antherids*, often described as 'spermatia' and 'spermogones,' are conveyed passively to the trichogyne, and there, becoming attached, fuse their contents into those of the trichogyne. There is a distinct difference here between the sexual organs in the parts they play and in their structure; though in the Erysipheæ, Penicillium, Sordaria, and Gymnoascus, the actual process of fertilisation has not been observed, yet the constant presence of the organs of sexuality justifies our regarding the function as extremely probable. The case is not so strong with regard to Melanospora and Ascobolus, where the antherid is either not constant in its appearance, or with difficulty to be distinguished from the hyphæ of the envelope. A third type is that of Polystigma, which closely resembles the Collemaceæ in the origin of the sporocarp. The carpogone, trichogyne, and antherid with pollinoids are here very much the same as in that group, only the sexual union has not been observed, and there is present in the early development of the carpogone and surrounding it, a special coil of pseudo-parenchymatous hyphæ which is ultimately concerned in the formation of the envelope-tissue of the sporocarp. Conforming also to the type represented by the Collemaceæ, and more especially allied to Polystigma in this respect, is the process as described for Xylaria. Here too there is a special coil of hyphæ, and in the interior of it a chain of large cells (like the carpogone of Polystigma) called by Füisting 'Woronin's hypha.' This hypha vanishes, and no connection has been discovered between it and the ultimate ascogenous hyphæ in the hypothece. Again no trichogyne has been observed to arise from 'Woronin's hypha,' and therefore no union with it of pollinoids, though bodies are recorded for Xylaria which resemble these male organs. The difference in the origin of the sporocarp between this type and Polystigma is essentially this: that here no distinct carpogone gives rise to the ascogenous hyphæ which have their origin in the special coil mentioned, since 'Woronin's hypha' disappears before this stage in the development of the sporocarp. Sclerotinia sclerotiorum (de By.) affords another type, since the ascogenous hyphæ and those of the

envelope, which are unconnected with each other though interwoven in the hypothece, may be traced down to the apparently uniform tissue of the stalk. De Bary thinks, however, that these may have a distinct origin—the ascogenous hyphæ in those proceeding from the original coil formed in the sclerote, and the envelope-hyphæ in others springing with them from the sterile tissue of the sclerote—that the coil in fact may be (or may contain) a carpogone, though there is no sign of antherid or pollinoids. The tracing of the continuity of these hyphæ is impossible throughout the long stalk with its uniform tissue. With this type are to be classed the lichen-fungi investigated by Krabbe (Sphyridium, Flot., Bæomyces, Pers., and Cladonia, Hill) the sporocarps of which exhibit a clear distinction between ascogenous and envelope hyphæ from an early stage. The former, however, have not been traced to a carpogone or other special initial organ, and neither antherid nor pollinoid has been discovered. Such genera as Claviceps, Epichloe, Pleospora, &c., in which all that has been observed is a gradual specialisation of uniform hyphæ to the functions of ascogenous and envelope hyphæ follow this last type as the farthest removed from those forms described under the second type.

Course of Development.

Within this large class several more or less distinct types of the course of development may be recognised. Just as in the case of the origin of the sporocarp, *Eremascus albus* is among the simplest of all. This remarkable type (the sporocarp of which is destitute of envelope-tissue) possesses no other reproductive bodies than the ascospores produced in the single ascus which represents the sporocarp. That is to say, the germinating ascospore gives rise to a thallus which directly bears the sporocarp again. In the same case are also the Collemaceæ with probably all lichen-fungi, and Pyronema, Ascobolus, Gymnoascus, Hypocopra, and Sclerotinia sclerotiorum, which last, however, possesses other propagating bodies in the shape of mycelial resting-cells comparable with the soredes of lichens. Another type (to which Sclerotinia Fuckeliana, de By. and Wor., belongs) is but one remove beyond this, since the life-history may go directly from sporocarp to sporocarp, though in the typical state *acrospores* intervene. Here the germinating ascospore gives rise to a thallus which bears acrospores, or, as has been said, the sporocarp again ; though never both together or successively on the same thallus. The germinating acrospore in turn produces a thallus with precisely the same properties and capabilities as that arising from the ascospore, except that in some cases there appears to be a tendency to

go on producing other acrospore-bearing thalli. In a third type acrospores are always formed (so far as experience goes) on the primary thallus produced by the germinating ascospore. This type de Bary subdivides into two as follows : (*a*) the primary thallus arising from the ascospore is only a promycele bearing *sporids*, which in turn produce a definite thallus, which either bears a sporocarp at once, or develops as sub-type (*b*). There is here then a necessary intervention of *acrospores* (sporids) before the sporocarp is again formed. (*b*) The primary thallus arising from the ascospore is a definite one, which ultimately bears the sporocarp, it is true, in all cases of complete development, but not until acrospores have been formed on it. The sub-type to some extent suggests the first type, since the acrospores are not *morphologically* a necessary intervention, though their appearance is invariable. The thallus produced by the germinating acrospore resembles the primary one in all respects. The development in this type may stop short with the formation of acrospores, and this is often repeated in succeeding generations.

The spores of such intervening states are invariably acrospores, and they appear either singly or on hymenia on the free surface of the thallus ; or they are produced in *pycnids*, bodies resembling peritheces. The spores so produced are termed stylospores, or better, *pycnospores*, as de Bary proposes. A species may produce only one of these kinds of acrospore, e.g. Erysiphe ; or under favourable circumstances more than one kind, e.g. Pleospora.

1. Erysipheæ.—The mycele of the Erysipheæ infests the surface of green living plants, through the epiderm of which it sends down *haustoria* into the tissues beneath. The mycele is delicate and cob-web like in appearance and consists of branching septate hyphæ, and is secured, so to speak, to the host-plant by means of the haustoria. In the course of the branching of these hyphæ they frequently cross each other, and at such points of contact the sporocarp is formed. If Podosphæra (Kze.), which has only a single ascus, be taken for the sake of simplicity, its development may be described as taking place in this fashion. From one of these crossing hyphæ, at the point of contact there springs an oval cell, the carpogone, which is separated by a transverse wall from the hypha. From the other hypha, likewise at the point of contact, there springs also a cell, longer and thinner than the other, which is similarly cut off by a septum. It overtops the carpogone to which it adheres, and the upper portion, which is slightly bent over the carpogone, is farther cut off from the lower by a transverse wall. The upper portion is the antherid and the lower its stalk. From the hyphæ at the base there now grow up a number of tubes, which envelop

the carpogone and form the single-layered outer perithecial wall. From the inner surface of the cells composing this wall there subsequently arise a number of other cells forming an inner wall several cells thick. The growth of these separates the antherid from the carpogone, and it takes part in the formation of the outer wall. From the outer wall, the cells of which have become larger and brown in colour, fine *rhizoids* are produced near the base, and in some species a few fine hairs at the apex termed *appendiculæ*. Meanwhile the carpogone has divided into two cells, one the ultimate ascus, and the other its pedicel-cell. Within the ascus finally eight ascospores are formed.

In Erysiphe (Hedw.) the chief points of difference from Podosphæra



FIG. 303.—*I*, *II*, *Podosphæra pannosa* de By. and Wor. *I*, chain of spores on sporophore and mycele. *II*, ripe sporocarp with ascus, *a*, emerging through wall of sporocarp, *h*. *III*—*V*, *P. Castagnei* de By. and Wor., fertilisation. *III*, *c*, carpogone; *p*, antherid. *IV*, older state; *h*, hyphal branches of envelope. *V*, still older state in optical longitudinal section; *a*, ascus (× 600). (*I*, *II*, after Tulasne, *III*—*V*, after de Bary.)

to be noted are these. The antherid winds spirally round the club-shaped carpogone, which divides into a series of cells—produces a number of asci—and the inner wall of the perithece is more developed.

The germinating ascospore gives rise to a mycele provided with haustoria, on a suitable host, and from this thallus there spring short *sporophores* which produce successively a series of acrospores. The acrospores in turn produce a mycele exactly like the primary one from the ascospore—which like it, if completely developed, ends by bearing the sporocarp again. But owing to external conditions such as varying weather, nutrition, and the like, this consummation is frequently not reached, and acrospores only are then formed generation after generation.

For example, the acrospore form called Oidium Tuckeri (Berk.), which occurs abundantly, and is well known as vine-mildew, never produces in Europe, so far as is known, peritheces. These it is believed have been found on native vines in North America, which is supposed to be the home of the disease, the perithecial form being the fungus described as Erysiphe (Uncinula) spiralis (Berk. et Curt.).

The Erysipheæ are parasites infesting living flowering plants of many natural orders. Among the best known and most destructive are the above-mentioned vine-mildew; E. lamprocarpa (Lk.) on Compositæ, Plantago, Verbascum, Labiatæ; E. graminis (Lev.) on grasses; E. Martii (Lév.) on Umbelliferæ, clover, lucern, lupins, &c.; E. communis (Lk.) on Polygonum, Rumex, Convolvulus, Dipsacus, Lathyrus, Delphinium, Aquilegia, Ranunculus, &c. Podosphæra Kunzei (Lév.) attacks species of Prunus, and Podosphæra Castagnei (de By.) is a well-known mildew of hops, though it also attacks many other plants of different natural orders.

2. Eurotium (Link).—The carpogone is formed by the rolling up in corkscrew fashion of the tip of a mycelial hypha, the turns of which, four or five in number, gradually come into closer contact till they present the appearance of a hollow screw. It is then divided by transverse walls into as many cells as there are turns in the screw. From the bottom turn of the screw there grow up two or three branches of irregular diameter, which take an irregular course towards the apex, but remain in close contact with the outside of the carpogone. Sometimes one ascends by way of the inside of the screw. One, however, climbs faster than the others and reaches the apex first; this is the antherid. Impregnation by it having taken place at the apex of the carpogone after the absorption of a minute portion of the wall, both the antherid and the other branches of the basal turn of the carpogone which follow it proceed to branch copiously, the hyphæ being septate, until the carpogone is completely enveloped. In this way the outer perithecial wall is formed as in the Erysipheæ. From it, as in the last type too, the inner wall grows inward, filling up the space between the outer wall and the carpogone with several layers, and pressing apart the turns of the screw. The wall-cells are of a pseudo-parenchymatous appearance, and the membrane of the outer wall becomes covered externally by a golden-coloured substance. The whole of these envelope-cells, it should be mentioned, increase in volume considerably. From the carpogone there now proceed numerous ascogenous hyphæ, which press among and suppress the inner wall-cells, and, branching plentifully, bear at the ends of the branches oval asci. These contain each eight ascospores. So copiously does this take place that, of the ascogenous hyphæ soon only

the traces may be seen, and by the time of maturity even the ascus-walls disappear and the perithece contains little but ripe ascospores.

When the ascospore germinates it produces a mycele, on which there shortly arise upright sporophores with round swollen apices bearing numerous short *sterigmata* over the surface. On the sterigmata chains of acrospores are formed successively, which, proceeding radially from the

FIG. 304. *Eurotium repens* de By. *A*, branch of mycele with sporophore, *c*, and sterigmata, *st*; early stage of carpogone at *as*. *B*: spirally twisted carpogone, *as*, antherid, *φ*, and an envelope-hypha. *C*, older state with more envelope-hyphæ. *D*, young sporocarp. *E* and *F*, young sporocarps in optical longitudinal section. In *E* the inner wall is beginning to be formed; *w*, the outer wall; *f*, the nner wall and other cells filling space between it and carpogone. *G*, ascus with spores. *H*, ascospore of *E. herbariorum* Lk. (*A* × 190, the others × 600.) (After de Bary.)

apex of the sporophore, surround it with a globular mass of acrospores. The course of development is the same here as in the Erysipheæ, and generation after generation of acrospores is usually formed in succession without the myceles attaining to the formation again of the sporocarp—this being the result of the external conditions of life of the fungus.

The species of Eurotium are saprophytes, and are found inhabiting decaying plants, fruits, &c., and forming in such situations a loose mycele of delicate thin-walled cells. The common mould formerly called Aspergillus glaucus (Link) is the acrospore stage of Eurotium herbariorum (Link), and was believed to be an independent form before de Bary discovered the pleomorphism of this fungus and identified it with the sporocarp stage (Eurotium).

3. Penicillium (Link).—The sporocarp of Penicillium takes its origin in the winding round each other once or twice of two lateral branches of mycelial hyphæ. These are so like each other that it is impossible, from the observations made on them, to say which is male and which female—a question on which the ultimate development throws no light, since it is uncertain whether the ascogenous hyphæ proceed from either or from both, and moreover, besides being alike in formation, they are equal in activity. Neither has any observation been made of the conjugation of these presumptive sexual elements—though their position towards each other signifies a sexual union. Together with the outgrowths from these of numerous short ascogenous hyphæ, there arises from the neighbouring hyphæ of the mycele a dense growth which completely envelops the presumptive carpogone and becomes interlaced with the ascogenous hyphæ proceeding from it—these being at first thicker hyphæ than those of the envelope-tissue. However, with the growth of the whole body, the cells of the envelope increase considerably in volume, especially the central mass, and acquire thickened pitted cell-walls, while the layers nearer the circumference form themselves into an outer wall the cells of which have yellowish-brown membranes. The whole has a pseudo-parenchymatous appearance. The originally outermost cells are cast off, owing to their taking no part in the growth. While this development of the envelope has been going forward, the ascogenous hyphæ have been pushing in between the interstices of the cells, and sharing in the process of thickening of the cell-membranes. At this stage of the history of the sporocarp a period of rest intervenes lasting about six or seven weeks. This past, the ascogenous hyphæ begin anew their growth in vigorous fashion, and, branching copiously at the expense of the cells of the envelope, ultimately produce at the ends of the branches short thick twisted terminal branches, which bear serially strings of asci containing each eight ascospores. So far is this process carried that finally not only is the whole interior envelope-tissue used up, but the asci themselves disappear, leaving enclosed by the outer wall only a dense mass of ascospores. De Bary compares the existence here of two forms of ascogenous hyphæ—viz. the relatively slender form which traverses and uses up the envelope-tissue, and the short

twisted thicker form which bears the asci—with the occurrence of the two forms of hyphæ in the ripening sporocarp of Elaphomyces. Some authors, it may be remembered, place Penicillium and Elaphomyces side by side.

The germinating ascospore produces a mycele in all respects like that which bore the sporocarp, a much-branching anastomosing septate flocculent mycele, which bears acrospores serially in succession at the end of their characteristic sporophores. The sporophore arises from the mycele in the form of an upright septate stalk, which bears at its summit cymose branches ending in sterigmata of equal height. On the sterigmata are the chains of acrospores. Such a sporophore has a brush-like appearance, the stalk being the handle, and the branches, sterigmata, &c., the hairs. The sporophores arise in dense masses, and in all produce enormous numbers of spores. So densely do they occur in exceptionally favourable situations—in the case of Penicillium glaucum (Lk.)—that they are sometimes bound together in bundles, fasciated as it were, and bear at the summit a dense crown of chains of spores. This form was originally described as a distinct genus by the name of Coremium glaucum (Lk.). As in Erysiphe and Eurotium, so in Penicillium, the course of development, after the production of acrospores, may omit the formation of the sporocarp on the same thallus through external conditions being unfavourable to its development. This, in fact, is the usual case in Penicillium, and generation after generation of thalli bearing acrospores only, and there stopping short, intervene as a rule between sporocarp and sporocarp. Perhaps the commonest of all moulds is Penicillium glaucum (Lk.), occurring on decaying fruits, on bread, &c., &c., in the acrospore-bearing condition. The sporocarp occurs, very rarely, in dark places where there is a poor supply of oxygen, and mostly on bread.

4. Gymnoascus (Baranetsk.) and Ctenomyces (Eidam) do not differ in any very striking peculiarity from types already discussed. The origin of the sporocarp is characterised by the fact that while one sexual hypha entwines the other, it is the entwining one which is the carpogone — which subsequently produces the ascogenous hyphæ—and the other, round which the carpogone winds, is the antherid. The relative positions of these certainly recall the case of Eurotium, where the antherid occasionally ascends by way of the inside of the screw ; but here, on the other hand, the carpogone takes, as it were, the active step, and winds round the antherid. These have their origin as lateral shoots either of the same hypha or of different hyphæ ; or it may be that only the entwining one (carpogone) is a lateral shoot, and the other merely the intercalary portion of a hypha. The ascogenous hyphæ branch

copiously, and ultimately bear at the ends asci containing each eight ascospores, and the envelope-tissue is contributed by shoots from the neighbouring mycele and from the base of the carpogone.

As already mentioned above, no acrospore stage intervenes here between sporocarp and sporocarp, i.e. the ascospore, on germinating, produces a thallus, which again bears the sporocarp directly. Gymnoascus is a saprophyte growing on dung.

5. Ascobolus (Pers.).—The carpogone arises on the mycele in the form of a thick curved sausage-shaped lateral hypha, which becomes divided by transverse walls into six or seven cells, about as long as they are broad. While in this stage the farther end of it is clasped by the branching end of a much thinner hypha—like one of the ordinary mycelial hyphæ—which, it may be assumed, is the antherid, from analogy with those types already discussed. At all events, it soon loses its identity in the dense growth of envelope-hyphæ which, immediately after this stage has been reached, are produced both from the ordinary mycele and from the hyphæ which bear the carpogone and presumptive antherid. These envelope-hyphæ, which soon enclose the carpogone in a round mass with a differentiated rind, next proceed to develop in the upper region over the carpogone (which is situated in the basal portion) the subhymenial layer of a discocarp. From this subhymenial layer there rise upward the straight perpendicular paraphyses. By the time the development has gone so far, the carpogone gives rise to a dozen or more ascogenous hyphæ from a cell near the middle of the row, which has manifestly obtained from its neighbours contributions from their contents. The ascogenous hyphæ grow upward to the subhymenial layer, where they branch and spread about among the roots, so to speak, of the paraphyses, and here bear asci. The asci grow straight upward among the paraphyses to the hymenial surface. The portion of the rind (envelope-tissue) immediately over the hymenial surface is ruptured, in consequence of the expansion of this surface



Fig. 305. *Ascobolus furfuraceus* Pers. Young sporocarp in longitudinal section (diagrammatic). *m*, mycele; *h*, hymenium; *c*, carpogone with ascogenous hyphæ, *s*, in the subhymenial layer, and *a*, asci (shaded); *l*, antherid; *p*—*r*, tissue of envelope giving rise to paraphyses. (After Janczewski.)

during growth, to permit the escape of the ascospores, and as new asci are produced (mostly taking the place of older ones), the expansion often continues till the hymenial surface becomes convex.

Ascobolus, like Gymnoascus, has no intervening acrospores, and the germinating ascospore gives rise to a thallus which bears the sporocarp directly. It is a saprophyte, and the species abound on dung.

6. Pyronema (Fckl.).—Pyronema confluens (Tul.) (or Peziza confluens, Pers., as it was formerly called), which, when mature, forms a discocarp like Ascobolus, differs considerably from that genus in the structure of the carpogone and antherid, though both, doubtless, belong to



Fig. 306.—*Pyronema confluens* Tul. *A* : *c*, carpogones; *a*, antherids; *t*, trichogynes. The trichogyne marked *t* has not yet become united with *a*. *B*, older state. The trichogyne *t*, proceeding from carpogone *c*, and cut off by a transverse wall, is in open union with *a*. *C* : the antherid, *a*, is in communication through *t* with carpogone, *c*, which is swollen and emitting ascogenous hyphæ (× about 300). (From de Bary, after Kihlman.)

the same main type in this respect. On the mycele of Pyronema confluens there arise clusters or rosettes of more or less club-shaped cells by forked branching at the summits of erect hyphæ, occurring generally in pairs ; these pairs in turn having their origin in densely branched groups of hyphæ. The rosettes consist each of three kinds of cell : the broad club-shaped, slightly curved cells are carpogones, and are usually borne on two pedicel-cells ; the antherids are also club-shaped, of the same height, but of about half the breadth ; and the third kind are sterile cells of cylindrical form. Two or three pairs of carpogones and antherids are included in each rosette. From the top of the carpogone there grows forth a slender curved trichogyne, with plentiful supply of

protoplasm ; and this organ, instead of behaving passively in the operation, bends over till it touches the top of an adjoining antherid, to which it adheres. When this has taken place, the trichogyne becomes separated by a transverse septum at the base from the carpogone, and then, through the dissolution of the intervening membrane, the contents of trichogyne and antherid are mingled. After impregnation the carpogone increases in volume, and from numerous points of its surface there are emitted ascogenous hyphæ. From the sterile cells and the whole basal region of the rosette, the envelope-hyphæ now grow forth and form a large *stroma* enclosing the carpogones and antherids—the latter remaining almost unaltered, full of protoplasm, and taking no part in the formation of the envelope—and upon the stroma a free hypothece bearing the paraphyses. In the production of asci and the farther development of the apothece, Pyronema agrees with Ascobolus.

Pyronema, like Ascobolus and Gymnoascus, produces no acrospores; and sporocarp follows sporocarp without intervention on successive thalli.

7. Sordaria (Ces. and de Not.) and Melanospora (Corda), both Pyrenomycetous forms, are placed next Ascobolus by de Bary with respect to the morphology of their sporocarps—of course excluding such differences of form as are peculiar to their being Pyrenomycetes, while Ascobolus is, as has been shown, a Discomycete. Fundamentally there is little real difference in the mode of origin of the sporocarp ; and Chætomium and Ascotricha may eventually prove to belong to the same series.

8. Collemaceæ.—In the Collemaceæ, a group of discocarpous *lichens*, the structure, development, and mode of behaviour of the male sexual element is wholly different from any hitherto described, while the carpogone remains of similar structure, though, of course, modified to meet the requirements of the changed conditions. The male cells are *pollinoids* formed in a flask-shaped antherid, somewhat resembling a perithece. It is sunk beneath the surface of the thallus, and opens by means of a narrow neck. The flask consists of a wall of densely-compacted hyphæ, giving off towards the interior a layer of numerous delicate hyphæ (sterigmata), which converge towards the central portion of the flask. These form at their apices successively in series numerous pollinoids, which soon fill the central space. The pollinoids are thin-walled rod-shaped cells, with an outer membrane of a gelatinous kind, readily swelling and dissolving in water. In damp rainy weather water gets access to the pollinoids, and through the swelling action mentioned they are forced out through the neck of the antherid and dispersed over the surface of the thallus. The development of the antherid

generally somewhat precedes the origin of the carpogone. Under the surface of the thallus a hypha not distinguishable from its neighbours gives off a broader lateral branch, which coils itself up two or three times, and then sends forth the tip of the coil, which, growing upward, emerges through the surface of the thallus into the open. The tube is commonly somewhat swollen as it passes through the superficial tissue, and for some short distance above it, and attains a height above the surface of four or five times its breadth. This is the *trichogyne*. The coil as it grows is divided by transverse walls into about a dozen thin-walled cells, and the trichogyne likewise into a similar number. Its development having taken place, and the suitable conditions of moisture having dispersed the pollinoids over the surface, these, wher-



FIG. 307.—*A*, *Leptogium microphyllum* Ach., section of thallus. *a*, point of trichogyne; *g*, algal cells; *h*, hyphæ. *B*, *Collema pulposum* Bernh., young carpogone. *C*, trichogyne with pollinoids of *L. microphyllum*. *D*, a similar one showing union with pollinoid. (*A* × 350, *B*, *C*, *D* × 750.) (After Stahl.)

ever they come into contact with a trichogyne, stick to it, sometimes in considerable numbers, and an open communication between pollinoid and trichogyne is established by means of a short minute process from the pollinoid. When this has been accomplished the cells of the trichogyne collapse, remaining distended only where a transverse septum occurs in its course, while the cells of the coil increase in volume and in number through the growth of fresh transverse walls. The neighbouring thallus-hyphæ then give out numerous shoots, which not only grow round the coils, but press them asunder. The hyphæ on the side next the surface then give off branches in that direction, the end shoots of which form the first paraphyses, after displacing the intervening tissue in their course. The enveloping hyphæ extend laterally until a basin-

shaped pseudo-parenchymatous exciple is formed, the margin of which reaches the surface. The interior of this basin is then soon filled with upright paraphyses like those which originally attained the surface. The turns of the original coil become unloosed, and eventually there are given off from it ascogenous hyphæ which, after branching in the sub-hymenial zone like those of Ascobolus, finally produce successive asci in the mature apothece.

In *Physma* Massal. the carpogone is produced from the hyphæ which form the wall of the antherid, and the trichogyne rises to the surface outside the wall. Eventually the paraphyses and asci are produced in the place formerly occupied by the sterigmata of the antherid,

FIG. 308.—*Leptogium microphyllum* Ach., young apothece in section. *h*, hyphæ; *g*, a lgal ce *a*, thalloidal exciple; *b*, exciple proper; *c*, hypothece. The apothece is filled with paraphyses, among which may be seen bladder-shaped ascogenous hyphæ with three young asci (× 530). (After Stahl.)

and the antherid is thus transformed into the sporocarp. The species of Collema (Ach.) have no acrospores, and resemble in the course of their life-history Ascobolus, Pyronema, &c. For structure of lichen-thallus, see p. 318.

9. POLYSTIGMA (Pers.) is a genus of parasitic fungi bearing a striking resemblance to the Collemaceæ in its sexual reproductive apparatus. Antherids and pollinoids are formed differing in no material respect from those of the Collemaceæ—the pollinoids being in this case more thread-like and bent. The genesis of the sporocarp is characterised here, however, by the formation of a fundamental coil of hyphæ smaller than the thallus-hyphæ. The carpogone appears among these, consisting of a spirally-wound hypha of two or three turns, with broad short cells.

It traverses the fundamental coil, and its apex grows out as a trichogyne, which protrudes through a stomate of the leaf of the host-plant. Pollinoids have been observed attached to it, but so far no actual communication has been detected. Accompanying the trichogyne are a few fine hyphæ, which, after the collapse of the trichogyne, protrude through the stomate like the tip of a small brush. The farther development of the sporocarp (the envelope-tissue arising from the fundamental coil) is like that of the last type, with this difference, that, instead of an apothece, it is a perithece which is here produced.

The ascospore produces on germinating a short promycelial tube, the end of which bears a sporid. The sporid, on germinating in turn on the leaf of a suitable host (Prunus), sends its germ-tube through the outer wall of the epiderm, and branches within the epidermal cell, these branches again penetrating into the parenchyme of the leaf. Here a thallus is formed, which remains covered by the epiderm of the leaf of the host until the sporocarp is again produced.

10. Xylarieæ, &c. In Xylaria (Hill), Hypoxylon (Bull.), Ustulina (Tul.), Diatrype (Fr.), Stictosphæria (Tul.), Eutypa (Tul.), Nummularia (Tul.), and Quaternaria (Tul.), we have the occurrence of a fundamental coil preceding the formation of the peritheces, followed by the gradual disappearance, as mentioned above for Xylaria, of 'Woronin's hypha,' which is formed in it, without its taking part in the formation of the ascogenous hyphæ. These arise, together with the paraphyses, from the perithecial wall. No trichogyne is formed from 'Woronin's hypha,' and there are no antherids. Before the formation of the peritheces in Xylaria there are borne on the same stroma in dense hymenia bodies which resemble acrospores, or it may be pollinoids. In this genus they have not been observed to germinate, though similar bodies germinate freely in Ustulina, e.g., and other genera. In Xylaria at all events they may be functionless pollinoids persisting in an organism in which indications of a carpogone (in the form of 'Woronin's hypha') also still remain.

11. Sclerotinia (Fckl.).—The sporocarp of Sclerotinia sclerotiorum (de By.) is in the form of a disc at first cup-shaped, borne at the summit of a stalk arising from a sclerote. It takes its origin from an entangled primordial coil of hyphæ with gelatinous membranes situated just beneath the dark peripheral cells of the sclerote. There are many such coils in each sclerote, but they do not all attain this farther development. The bundle of hyphæ constituting the stalk of the sporocarp breaks forth, as has been said, from this region of the sclerote, the central portion of the hyphæ arising from the coil, and the external hyphæ from the ordinary tissue of the sclerote. In the growth of the stalk it has

been found impossible to trace any anatomical distinction between these elements, and therefore it amounts to no more than a probability that the hyphæ from the coil are the ultimate ascogenous hyphæ; while those from the tissue of the sclerote may furnish the envelope-tissue of the sporocarp. The main difference between the development of S. sclerotiorum and S. Fuckeliana in this respect consists in the primordial coil of the latter originating not within the sclerote, but in the central portion of the bundle of stalk-hyphæ after it has reached the external surface.

S. sclerotiorum possesses, so far as is known, no intervening acrospores. The germinating ascospore produces a mycele on which sclerotes are formed, and on these only the sporocarps again. S. Fuckeliana sometimes does the same, and de Bary has observed a single instance of the sporocarp being produced on the mycele without even the formation of a sclerote. The sclerotes of this species, however, frequently produce filamentous sporophores bearing acrospores, this stage being that formerly known as Botrytis cinerea (Pers.). It never happens that a sclerote bears both acrospores and sporocarps, either together or after each other. The primary mycele may bear acrospores directly without interfering with the subsequent production of sclerotes, but this does not happen often. The mycele produced by the germinating acrospores is similar in all respects to the primary one arising from the ascospore, with the reservation that it has a greater tendency than the other to the production of sporophores. In this species there are also often formed certain abortive acrospores, or it may be pollinoids.

12. Pleospora (Rabenh.).—In the origin of the perithece of Pleospora herbarum (Rabenh.), the traces of sexuality disappear from view, and indeed it is stated that the asci arise among the paraphyses as branches of the basal cells of the latter. In the life-history of this fungus a considerable number of forms are embraced. Besides the ascospores, which are compound multicellular bodies, there are in the second category acrospores of three sorts, viz. (*a*) Double or multicellular acrospores of rounded short cylindrical form, previously taken to be an independent species (Macrosporium Sarcinula, Berk.). Each such compound spore appears as a rule singly on its sporophore. (*b*) Conical pear-shaped likewise compound spores appearing in series, often in branching series. This was also formerly described as an independent species of Alternaria (Nees ab Esenb.). (*c*) A small form of acrospore recorded by Bauke, but not, according to de Bary, Cladosporium herbarum (Link), which, though placed in genetic connection with Pleospora herbarum by Tulasne, does not belong here. In the third category there are pycnospores formed in pycnids interstitially arising on mycelial hyphæ. The pycnids consist of a wall of several layers, from the inner surface of

which there converge series of cells producing successively (terminally and laterally) pycnospores. These are about twice as long as broad, and very thin-walled, but surrounded by a hyaline gummy substance. Not only are all these forms on record, but the mycele shows a tendency to pass into a resting state, and single cells or groups of cells becoming detached add to the means of propagation. There is some contradiction involved in the accounts of the occurrence of some of

FIG. 309.—*Claviceps purpurea* Tul. Longitudinal section of portion of sclerote, *m*, in *Sphacelia* stage, producing conidiospores, *c* (much magnified). (After Tulasne.)

these forms and the order of their succession given by Bauke and by Gibelli and Griffini, and further research may be expected to throw light on the matters in dispute, as also on the question whether or not we have here one species very rich in spore-forms or two species resembling each other, but each less rich in forms.

13. CLAVICEPS (Tul.).—The peritheces of Claviceps purpurea (Tul.) appear as it were in a kind of capitulum (immersed in the same stroma),

FIG. 310.—Ear of rye with two mature sclerotes (ergot) of *Claviceps purpurea* (natural size). (After Luerssen.)

borne on the summit of a stalk arising from a sclerote (the well-known *ergot* of grasses); the ascospores are filamentous in form, and on germinating produce hyphæ at several points. When this takes place under suitable circumstances on the pistil of grasses, the tubes enter, and, besides penetrating the tissue, form at length a white hymenium on the surface. This hymenium consists of numerous cylindrical sterigmata, which bear acrospores at their apices. (This stage was described by Léveillé by the name of *Sphacelia*.) These are given off in drops of a sugary juice which oozes out between the flower-leaves in which the pistil lies. This juice, 'honey-dew,' is eaten greedily by insects, which doubtless eat the acrospores with it. These propagate the Sphacelia state. At the base of this acrospore-forming body the formation of the sclerote proceeds, and it ultimately replaces the ovary, emerging from the flower of the grass with its base seated on the floral receptacle. It gives rise in turn to the sporocarp again.

The allied genera Cordyceps (Fr.) and Epichloe (Fr.) agree with Claviceps in the development of the sporocarp.

Besides the above distinctly marked types, the origin of the sporocarp and the life-history of a considerable number of other forms have been more or less completely investigated. It would be entirely beyond the proportions of the present book to enter into a description of these, and a discussion of the controversies that are bound up with the accounts of the multitude of incompletely known forms. The citation of the literature at the end of this section will guide the student to original papers, but he may also be referred here, as in so many other cases of difficulty, to de Bary's 'Comparative Morphology and Biology of Fungi,' &c., 1887, as containing an exhaustive and critical treatment of these more or less obscure forms.

He will find there also more information on the development of the sporocarps (both discocarpous and pyrenocarpous) of the types



FIG. 311.—*Claviceps purpurea* Tul. *A*, sclerote which has produced seven stromata. *B*, upper portion of a stroma in longitudinal section. *cp*, perithecia. *C*, longitudinal section of perithece. *cp*, ostiole; *sh*, cortical tissue; *hy*, inner tissue of stroma. *D*, ascus isolated. *sp*, ascospores issuing. (*A*, natural size, *B* slightly, *C* and *D* highly magnified.) (After Tulasne.)

quoted than it is possible to give in this place without burdening the section beyond proportion with matter of no striking morphological significance.

HOMOLOGIES OF THE ORGANS.

De Bary, the founder of the system of classification of fungi adopted in the present book, calls attention to the striking parallelism between Ascomycetous forms on the one hand and the Mucorini, Peronosporeæ, and Saprolegnieæ on the other. From the oosperms and the spores of the sporocarps there arises a thallus which closes its development with the formation of sexual organs and the oosperm and sporocarp resulting from their union. In many cases the life-history is confined to this; for example, from the Phycomycetes, Pythium vexans, &c.; from the

Ascomycetes, Eremascus, Pyronema, and Ascobolus. In most cases, however, there intervenes the formation of non-sexual spores which may be in a species all of one sort, e.g., Erysiphe, Peronospora ; or of several sorts ; and such spores are, moreover, in many cases exceedingly alike.

Eremascus might almost belong to the Mucorini (Piptocephalideæ), while on the other hand it is not wanting in the essential attributes of an Ascomycete. In the form of its sexual organs it completely resembles Penicillium, Gymnoascus, Eurotium, &c. Again, great agreement is to be recognised between the thallus, spores, and sexual organs of the Erysipheæ (Podosphæra especially) and those of the Peronosporeæ. The resemblance ceases with the farther development of the results of sexual union, and at this point the groups diverge from the point of contact of Eremascus and Podosphæra with the Mucorini and Peronosporeæ. Of course the envelope-apparatus of the asci is not included in any comparison, as being of purely secondary importance ; and since such envelope is actually wanting in Eremascus, the case is made the clearer. It is also pointed out that though the oogones of the Peronosporeæ have no envelope, it is by no means impossible that such may be found, while the zygosperms of Mucorini, as has been shown, are sometimes provided with an envelope. On such evident grounds as these, cited from de Bary ('Comp. Morph. and Biol. of Fungi,' p. 232), who enters minutely into the matter (as well as the subject of the sexuality of the Ascomycetes), the relationship of these groups with each other is abundantly established.

Doubtful Ascomycetes.

1. Laboulbenieæ.—The Laboulbenieæ are a small assemblage of remarkable parasites on insects, attacking mostly water-beetles, but also other insects, including the house-fly. They possess no mycele, and occur fixed on the chitin of the insect attacked by means of a short process which serves as a *haustorium*, if that name may be applied to it. Above this rises a stalk consisting usually of two cells, one above the other, bearing at the summit a simple perithece and a few lateral hairs (the *appendage*) composed of seriated cells sometimes having minute round swellings at their apices. Before the complete development of the perithece has been reached, there is emitted from its summit a short fine process, which may be a trichogyne ; and according to Karsten (Stigmatomyces, Karst.) the minute round swellings on the hairs become free and attach themselves to the trichogyne. Doubt, however, has been cast upon the accuracy of this observation by the investigation of Peyritsch, who also attaches no great value to the suggestion that the so-called

trichogyne is fertilised by the contact with it of one of the young hairs. The perithece contains a number of asci, and these eight or twelve double ascospores. The ripe double ascospore attaches itself to a fresh host by one of its ends, and develops into the new plant.



FIG. 312.—*A*; *b—h*, *Stigmatomyces Baeri* Peyr. (*St. Muscæ* Karsten). *A*, optical longitudinal section of ripe specimen with organ of attachment at base; the asci are seen through wall of perithece. *a*, everywhere the appendage; *b*, an isolated ascus with spores; *c—h*, stages of development of perithece and appendage in order of letters. *B*, *Laboulbenia flagellata* Peyr. *a*, the appendage. (*A*, *c*, *g*, *h* × 350; *b*, *d*, *e*, *f* × 450; *B* × 125.) (After Peyritsch.)

2. EXOASCUS (Fückel).—The species of Exoascus mostly attack fruits, and set up in them sometimes conspicuous deformities. While some of them possess a mycele which penetrates the parenchyme of the fruit, &c. (e.g. E. Pruni, Fckl., E. deformans, Fckl.), others extend no farther than between the cuticle and the epiderm-cells. In the former case the terminal cells of the hyphæ which emerge from the surface

become asci. In the latter case either certain cells become asci while others remain sterile or the whole body of hyphæ form asci. In E. alnitorquus (Sadeb.) these asci have a pedicel-cell; in E. aureus (Sadeb.) there is nothing but asci left at maturity. When the ascospores germinate they give rise to a yeast-like sprouting.

3. SACCHAROMYCES (Meyen).—The species of Saccharomyces occur in fermenting substances, and are well known from their power of converting sugar into alcohol and carbonic acid. Among the familiar species are S. cerevisiæ (Meyen) (ordinary yeast), S. ellipsoideus (Reess), S. Pastorianus (Reess), alcoholic ferments which are apparently mere form-species. With these should be placed S. Mycoderma of Reess, and Chalara Mycoderma of Cienkowski; and the 'thrush' fungus S. albicans (Reess), which lives parasitically on the mucous membrane of the human digestive organs, but is also capable of exciting a feeble alcoholic fermentation in sugar solutions. With the exception of the last-mentioned forms in which hyphæ occur, the species of Saccharomyces are unicellular fungi which increase by sprouting. The cells, in which a nucleus has not been demonstrated, are round or oval in form, and the sprouting takes place in the form of a protuberance, which gradually swells and becomes constricted and finally cut off by a wall at the point of origin. These new cells either separate at once, or chains or groups remain united as they have been formed. When such cells are cultivated, on the cut surface of a potato for example, certain cells may form asci, each containing two to four ascospores. These are at once capable of germination, which takes



FIG. 313.—*Saccharomyces cerevisiæ* Meyen. *a*, single cell of beer-yeast; *b*, *c*, stages of sprouting; *d*, colony of sprout-cells; *e*, cell with four ascospores; *f*, one with two; *g*, group of ascospores with one sprouting; *h*, further development of a similar group (*h* × 750, the others much more). (*e*–*h* after Reess.)

place by the sprouting process described, though they may retain the power of germinating for a longer period.

This cell-increase by sprouting is, as has been seen, by no means confined to Saccharomyces, but occurs in other groups of fungi (e.g. Mucor), and the special character which entitles them to this place in the classification of fungi is the production of asci, which they share only with the Ascomycetes. They may therefore be regarded as much degraded Ascomycetous forms—the other alternative, that they are early forms from which typical Ascomycetes have developed, being disposed of by the establishment of the connection of this great group with the Phycomycetes as already described.

LITERATURE.

Barenetski—Entwickel. d. Gymnoascus Reessii (Bot. Zeit., 1872).
De Bary—Ueber d. Fruchtentwickel. d. Ascomyceten (Leipzig, 1863).
De Bary—Beitr. zur Morph. u. Physiol. d. Pilze, iii., 1870.
De Bary—Exoascus Pruni (Beitr., i.).
De Bary—Ueber einige Sclerotinien, &c. (Bot. Zeit., 1886).
Bauke—Zur Entwickelungsgesch. d. Ascomyceten (Bot. Zeit., 1877).
Bauke—Beitr. zur Kenntniss d. Pycniden (Nova Acta Leop., xxxviii., 1876).
Borzi—Studii sulla sessualità degli Ascomicete (N. Giorn. Bot. Ital., x., 1878).
Brefeld—Botan. Untersuch., ii. and iv. ; compare also v.
Brefeld—Mucor racemosus und Hefe (Flora, 1873).
Brefeld—Ueber Gährung (Thiel's Landwirthsch. Jahrb., 1875, 1876).
Cienkowski—Die Pilze der Kahmhaut (Mélanges Biol. Acad. St. Pétersb., viii.).
Cornu—Reproduction des Ascomycètes (Ann. Sc. Nat., 6 sér., Tom. 11).
Currey—On the Fructification of certain Sphæriaceous Fungi (Phil. Trans. Roy. Soc., 1858).
Eidam—Beitr. zur Kenntniss d. Gymnoasceen (Cohn's Beitr., iii.).
Eidam—Zur Kenntniss d. Entwickel. d. Ascomyceten (*ibid.*).
Eidam—Ueber Pycniden (Bot. Zeit., 1877).
Engel—Les ferments alcooliques, 1872.
Fisch—Zur Entwickelungsgesch. einiger Ascomyceten (Bot. Zeit., 1872).
Füisting—De nonnullis Apothecii Lichenum evolvendi rationibus (Diss. inaug. Berol., 1865).
Füisting—Zur Entwickelungsgesch. d. Pyrenomyceten (Bot. Zeit., 1867, 1868).
Füisting—Zur Entwickelungsgesch. Lichenen (*ibid.*, 1868).
Gibelli—Sugli org. reprod. del gen. Verrucaria (Mem. Soc. Ital. di Scienc. Nat., i.).
Gibelli e Griffini—Sul polimorfismo della Pleospora herbarum (Arch. d. Laborat. di Bot. Crittog. Pavia, i., 1875).
Gilkinet—Recherches sur les Pyrénomycètes (Bull. Acad. Belg., 1874).
Hansen—Oidium lactis, &c. (Medd. fra Carlsberg Laborat., Bd. i.).
Hansen—Untersuch. über d. Physiol. u. Morph. d. Alkoholfermente (*ibid.*).
Hansen—Untersuch. über d. Organismen welche sich zu verschiedenen Jahreszeiten in d. Luft finden, &c. (*ibid.*).
Hansen—Untersuch. über d. Physiol. u. Morph. d. Alkoholgährungspilze (*ibid.*, Bd. ii.).

Hansen—Bemerk. über Hefepilze (Allg. Zeitsch. f. Bierbrauerei, &c., 1883; Bot. Centralbl., xvii.).
Hartig—Wichtige Krankheiten d. Waldbäume, p. 101.
Hartig—Untersuch. aus d. Forstbot. Institut zu München, i.
Janczewski—Morphol. d. Ascobolus furfuraceus (Bot. Zeit., 1871).
Karsten—Stigmatomyces (Laboulbeniaceen) in Chemismus der Pflanzenzelle, 1869.
Kihlman—Zur Entwickelungsgesch. d. Ascomyceten (Acta Soc. Sc. Fennicæ, xiii., 1883).
Krabbe—Entwickel., Sprossung und Theilung einiger Flechtenapothecien (Bot. Zeit., 1882).
Krabbe—Morphol. u. Entwickelungsgesch. d. Cladoniaceen (Ber. d. D. Botan. Gesellsch., 1883).
Kühn—Mittheil. d. Landw. Instituts Halle, i.
Lauder Lindsay—Spermogones and Pycnides of Lichens (Trans. Roy. Soc. Edinb., xxii.).
Low—Ueber Dematium pullulans (Pringsh. Jahrb., Bd. vi.).
Mattirolo—Sullo sviluppo e sullo sclerozio della Peziza Sclerotiorum Lib. (N. Giorn. Bot. Ital., xiv.).
Millardet—Mém. Soc. d'Hist. Nat. Strasbourg, vi., 1868 (Myriangium, &c.).
Pasteur—Mém. sur la fermentation alcoolique (Ann. Chim. et Phys., Tom. lviii., 1860).
Pasteur—Etudes sur la bière (Paris, 1876).
Peyritsch—Ueber die Laboulbeniaceen, &c. (Sitzber. Wiener Acad., Bd. 64, Abth. 1, 1871; Bd. 68, Abth. 1, 1873; and Bd. 72, Abth. 3, 1875).
Pirotta—Sullo sviluppo della Peziza Fuckeliana (N. Giorn. Bot. Ital., xiii.).
Rathay—Ueber d. Hexenbesen d. Kirschbäume (Sitzber. Wiener Acad., Bd. 83, Abth. 1).
Reess—Botan. Untersuch. über d. Alkoholgährungspilze (Leipzig, 1870).
Reess—Zur Kenntniss d. Exoascus d. Kirschbäume (Sitzber. d. Phys. Med. Gesellsch. zu Erlangen, 1882).
Reess—Ueber den Soorpilz (*ibid.*, 1877 and 1878).
Reess—Untersuch. ü. Elaphomyces granulatus (Deutsch. Naturf. u. Aerzte, Strassburg, 1885).
Reinke u. Berthold—Die Zersetzung d. Kartoffel durch Pilze (Berlin, 1879).
Schwendener—Ueber d. Entwickelung der Apothecien von Cœnogonium (Flora, 1862).
Schwendener—Ueber d. Apothecia primitus aperta u. d. Entwickel. d. Apothecien im Allgemeinen (*ibid.*, 1864).
Sorokin—Quelques mots sur l'Ascomyces polysporus (Ann. Sc. Nat., 6 sér., Tom. iv.).
Stahl—Beiträge zur Entwickelungsgesch. d. Flechten, i. (Leipzig, 1877).
Van Tieghem—Chætomium (Comptes Rendus, 81).
Van Tieghem—Nouv. observ. sur le développement du fruit, &c., des Ascomycètes (Bull. Soc. Bot. France, xxiii., 1876). (See also Bot. Zeit. same year.)
Van Tieghem—Sur le développement du fruit des Ascodesmis (*ibid.*).
Van Tieghem—Nouv. observ. sur le développement du périthèce des Chætomium (*ibid.*).
Van Tieghem—Sur le développement de quelques Ascomycètes (*ibid.*, xxiv.).
Van Tieghem—Monascus, genre nouveau de l'ordre des Ascomycètes (Bull. Soc. Bot. France, 1884).
Tulasne—Fungi hypogaei (Paris, 1851).

Tulasne—Selecta Fungorum Carpologia, i.–iii. (Paris, 1861–65).
Tulasne—Rech. sur l'organisation des Onygena (Ann. Sc. Nat., 3 sér., Tom. i.).
Tulasne—Note sur l'appareil reproducteur des Lichens et des Champignons (*ibid.*, 3 sér., Tom. xv.).
Tulasne—Mém. pour serv. à l'hist. organograph. et physiol. des Lichens (*ibid.*, 3 sér., Tom. xvii.).
Tulasne—Discomycètes (*ibid.*, 3 sér., Tom. xx.).
Tulasne—Mém. sur l'Ergot des Glumacées (*ibid.*).
Tulasne—Note sur l'appareil reprod. des Hypoxylées et des Pyrénomycètes (*ibid.*, 4 sér., Tom. v.).
Tulasne—Nouv. observ. sur les Erysiphés (*ibid.*, 4 sér., Tom. i.). (See also Bot. Zeit., 1853.)
Tulasne—Note sur les Isaria et les Sphæria entomogènes (*ibid.*, 4 sér., Tom. viii.).
Tulasne—De quelques Sphéries fongicoles (*ibid.*, 4 sér., Tom. xiii.).
Tulasne—Note sur les phénomènes de copulation d. l. Champignons (*ibid.*, 5 sér., Tom. v.).
Tulasne—Super Friesiano Taphrinarum genere (*ibid.*).
Vittadini—Monographia Tuberacearum (Mediolani, 1831).
Vogel—Gymnoascus uncinatus (Bot. Centralblatt, xxix.).
Wilhelm—Beitr. zur Kenntniss d. Pilzgattung Aspergillus (Diss. Berl., 1877).
Woronin—Entwickelungsgesch. d. Ascobolus pulcherrimus und einiger Pezizen (Beitr. zur Morph. u. Physiol. d. Pilze, ii.).
Woronin—Sphæria Lemaneæ, Sordaria, &c. (*ibid.*, iii.).
Zopf—Zur Entwickelungsgesch. d. Ascomyceten (Chætomium) (Nova Acta Leop., xlii.).
Zopf—Die Conidienfrüchte von Fumago (*ibid.*, xl.).

Class XXI.—Uredineæ.

The Uredineæ are a class of parasites on flowering plants and ferns. They resemble the Ascomycetes in many points, as will be seen from this short account of them. The mycele is septate and much branched, follows the intercellular spaces of the host-plants, and penetrates the cells themselves by means of short branches.

Puccinia graminis (Pers.) may be taken as a type of the course of development followed by some of the forms. Owing to the change of host involved in the course of development of this and other forms, and to the different appearances presented by succeeding stages of the organism, it was formerly supposed that these stages constituted distinct fungi. Thus no less than three form-genera (Æcidium, Pers., Uredo, Pers., and Puccinia, Pers.) were established to denote the stages of the life-cycle of Puccinia graminis, the well-known corn mildew. The sporocarp (Æcidium) is formed in spring on the barberry. In its



FIG. 314.—*A*, diagrammatic transverse section of barberry leaf with æcidia, *a*, *b*, and antherids, *s* (slightly magnified). *B*, uredospores, *u*, and teleutospore, *t*. *C*, germinating uredospore (*B* and *C* × 390). *D*, teleutospores. *E*, germinating teleutospore: promycele, *p*, and sporids, *sp* (× 400). (*A*, *B*, *D*, and *E* of *Puccinia graminis*, *C* of *P. straminis* Fckl., *F* teleutospore of *P. coronata* Cord., × 300). *G*, teleutospore of *Phragmidium incrassatum* Link. (× 300). (*A*, *F* and *G*, after Luerssen; *B*—*D*, after de Bary; *E*, after Tulasne.)

earliest stage it appears to consist of a densely interwoven mass of hyphæ situated in the subepidermal parenchyme of the leaf. This gradually increases in bulk and displaces the surrounding tissue. As it grows the hyphæ increase in size, and the shape of the whole becomes more definitely spherical. Within the base of this sphere the hymenium is developed, and consists of a continuous layer of club-shaped *basids*, from the summit of each of which is produced in basipetal succession a single series of *æcidiospores*. Enclosing the sporal mass is a single layer of pseudo-parenchymatous cells arising from the margin of the hymenium, and arching over the top of the spores. The cells composing the envelope are larger than the spores, and possess thicker walls, while their clearer contents contrast with the orange-coloured spores. The enlargement of the whole body increases until the epiderm is ruptured, while the tissues of the host are pushed aside and compressed. After the rupture of the epiderm the envelope bursts at the apex, and curves back, forming the lip of a cup-shaped body; the sporal mass is farther elevated and the spores escape. These sporocarps are situated on the under surface of the leaf, but accompanying them on the upper surface there are to be found numerous flask-shaped antherids containing pollinoids produced at the apices of sterigmata, and in all points recalling

those already described in Collema. They are orange-coloured like the sporocarps, and the pollinoids have never been known to germinate. No corresponding female sexual organ occurs in any Uredine, though the early stages of the development of the sporocarp are not sufficiently known. No such body as Woronin's hypha in Xylaria, for example, has ever been observed in the Uredineæ, and the only suggestion of a female sexual organ is to be found in the occasional occurrence in some Uredines of short obtuse hyphæ, projecting through the stomates of the host like the trichogynes of Polystigma. These may be traced, it is true, to young æcidia, but there may well be nothing more in the suggestion than the mere protrusion of mycelial hyphæ, since observations connecting such filaments with an act of fertilisation are wholly wanting. Massee ('Annals of Botany,' 1888, p. 47) has recently published an account of observations of a supposed sexual process in Uredineæ, involving the fertilisation of a carpogone by an antheridial branch; but the subject stands in great need of farther investigation.

FIG. 315.—*Puccinia graminis* Pers. *t* teleutospore; *u*, uredospores (×390). (After Sachs.)

The spores from the ripe sporocarp (*æcidiospores*) germinate only on the leaves or stems of grasses, and the germ-tubes entering by way of the stomates give rise to myceles, which attack the tissues of the host. In the course of a week or more, cushion-like masses of mycelial hyphæ situated beneath the epiderm give rise to erect basids, each of which bears a red *uredospore* (Uredo) at the apex. On the rupture of the epiderm the uredospores escape, and these alighting on grass plants germinate, again enter by way of the stomates, and renew the same generation. This process may and does go on indefinitely, and thus much damage is annually caused by the attack of this fungus on the corn crop.

Later on there are developed on the same mycele, first side by side with the uredospores, then gradually replacing them altogether, two-celled spores called *teleutospores*, and with their production the development of the fungus ceases for a period. In this condition the winter is passed. With spring they germinate, each of the two cells of the teleutospore (Puccinia) giving rise to a short promycele, the terminal cells of which bear on slender stalks a single sporid apiece. These sporids germinate in turn on the leaves of the barberry, the germ-tubes piercing the epi-

derm, and giving rise within to the mycele which ultimately bears the sporocarps and antherids.

Gymnosporangium (DC.) represents another type of the course of development in Uredineæ. The sporocarps (corresponding to æcidia, but here denoted by the form-genus Roestelia, Reb.) appear in summer on the leaves and fruits of Pomeæ. No uredospores are formed. The next generation (teleutospores) is produced in spring on juniper in odd-shaped mucilaginous brown or yellow masses. Promyceles are formed which bear sporids, and these again set up on the leaves, &c., of Pomeæ the sporocarp generation.

A farther reduced type is to be found in Endophyllum (Lév.). In this case the germ-tube of the spores of the sporocarp (æcidiospores) becomes a promycele, and, dividing up into several cells, each of these bears at the end of a sterigma a single sporid. The sporid on germinating renews the sporocarp generation.

Two cases of exceptional structure may be noted. In Phragmidium (Link) the sporocarps have no proper envelope, the place of the wall being taken by a circle of club-shaped paraphyses surrounding the margin of the hymenia. On the other hand the uredospores of Melampsora populina (Jacq.) and of Cronartium (Pers.) are enclosed in an envelope resembling that of the sporocarp. The development of the paraphyses in the one case and of the envelope in the other requires investigation.

Besides the sporocarp-forming Uredineæ there is another group known as the *tremelloid Uredineæ*, which do not possess a sporocarp generation. These are not to be confounded with those Uredineæ in which presumably from want of investigation the sporocarps are unknown. The course of development of the tremelloid Uredineæ is perfectly well known in a number of cases (Leptopuccinieæ and Leptochrysomyxa, de By.), and consists of a teleutospore-bearing generation with commonly softer and more gelatinous spore-membranes. These teleutospores germinate as a rule at maturity and not after a period of rest. The sporids formed on the promycele produce a mycele which again bears teleutospores. Leptopuccinia malvacearum (Schroet.), L. Dianthi (Schroet.), &c., bear the same relation in appearance, &c., to Puccinia as Leptochrysomyxa Abietis (Ung.) bears to Chrysomyxa (Ung.), the species of which form sporocarps, uredospores, and teleutospores.

Enough has been said in this brief account to indicate a probable connection of the Uredineæ with the Ascomycetes through their sporocarps. Those forms—the tremelloid Uredineæ—in which the sporocarp generation may be presumed to have been lost, sufficiently resemble the complete types to be necessarily bound up with them ;

while on the other hand, as will be seen, they furnish a valuable link with the next class.

LITERATURE.

De Bary—Untersuch. über die Brandpilze (Berlin, 1853).
De Bary—Rech. sur le développement de quelques Champignons parasites (Ann. Sc. Nat., sér. 4, xx.).
De Bary—Ueber Cæoma pinitorquum (Monatsber. Berl. Akad., 1863).
De Bary—Neue Untersuch. über d. Uredineen (*ibid.*, 1865, 1866).
De Bary—Ueber d. Krebs u. d. Hexenbesen d. Weisstanne (Bot. Zeit., 1867).
De Bary—Æcidium abietinum (*ibid.*, 1879).
Dietel—Beitr. z. Morph. u. Biol. d. Uredineæ (Bot. Centralblatt, xxxii., 1887, p. 54; *ibid.*, 1888, No. 33).
Farlow—The Gymnosporangia or Cedar Apples of the United States (Mem. Boston Soc. Nat. Hist., 1880).
R. Hartig—Wichtige Krankheiten der Waldbäume (Berlin, 1874).
R. Hartig—Lehrbuch der Baumkrankheiten (Berlin, 1882).
Léveillé—Sur la disposition méthodique des Urédinées (Ann. Sc. Nat., sér. 3, viii.).
Oersted—Om Sygdomme hos Planterne, &c. (Kjöbenhavn, 1863).
Oersted—On Podisoma and Roestelia (Oversigt k. Danske Vidensk. Selskab. Forhandl., 1866; and K. Danske Vidensk. Selskab. Skrifter, ser. 5, vii.).
Parker—On the Morphology of Ravenelia glandulæformis (Proc. Amer. Acad. Sc. and Arts, 1886).
Rathay—Untersuch. über d. Spermogonien d. Rostpilze (Denkschr. d. Wien. Akad., 1883, Bd. xlvi.).
M. Reess—Die Rostpilzformen d. deutsch. Coniferen (Abhandl. Nat. Gesellsch. zu Halle, Bd. xi.).
Schroter—Die Brand- u. Rostpilze Schlesiens (Abhandl. Schles. Ges. vaterland. Cultur, 1869 [1872]).
Schröter—Entwickelungsges. einiger Rostpilze (Cohn, Beitr. i., Heft 3; *ibid.*, iii., Heft 1).
Schröter—Ueber einige amerikanische Uredineen (Hedwigia, 1875).
Schröter—Beobacht. über d. Zusammengehörigkeit von Æcidium Euphorbiæ u. Uromyces Pisi (*ibid.*, 1875).
Tulasne—Mém. sur les Ustilaginées et les Urédinées (Ann. Sc. Nat., sér. 3, vii.; *ibid.*, sér. 4, ii.).
Ward—Researches on the Life-history of Hemileia vastatrix (Linn. Soc. Journ. Bot., xix.).
Ward—On the Morphology of Hemileia vastatrix (Quart. Journ. Micr. Sc., N.S., xxii., 1882).
R. Wolff—Æcidium pini u. seine Zusammenhang mit Coleosporium Senecionis Lev. (Festschrift, Riga, 1876).

In the above papers will be found references to older literature and farther memoirs on the group. For a systematic account the student is specially referred to Winter's Die Pilze Deutschlands, Oesterreichs und der Schweiz, in the new edition of Rabenhorst's Kryptogamenflora von Deutschland, &c.; and to Plowright's British Uredineæ and Ustilagineæ (London, 1889).

Class XXII.—Basidiomycetes.

The Basidiomycetes are a large class comprising forms of the utmost diversity in appearance, mostly saprophytes living on humus, rotten wood, or the old wood and the bark of trees. A small number are parasites. They all agree in the production of spores (*basidiospores*) acrogenously on *basids*, which are club-shaped and disposed as a rule parallel to each other, thus forming *hymenia*. The spores produced on one basid are two or four in number, more rarely eight, though divergences from these numbers occur. They vary in shape, but consist, except in some Tremellineæ, of a single cell. Among the basids there commonly occur sterile hyphal branches—*paraphyses*. Besides these spores thus borne on definite hymenia there are also others produced more or less indefinitely on the myceles of certain members of the group, and their character will be described below. The Basidiomycetes are divided into two sub-classes, the HYMENOMYCETES with gymnocarpous, and the GASTEROMYCETES with angiocarpous fructification.

Sub-class 1.—Hymenomycetes.

The Hymenomycetes are characterised by the possession of a *hymenium* on the free exposed surface of the compound structure which bears it—the *sporophore*. The forms embraced in this sub-class range from very simple to highly complex structures, the latter being repre-

FIG. 316.—*Tremella mesenterica* Retz. (natural size). (After Tulasne.)

sented by such types as the common mushroom and the like—in short those fungi to which the name is popularly applied.

EXOBASIDIUM VACCINII (Woron.) may be taken as the simplest type. Its mycele is parasitic on the leaves and stems of Vaccinium vitis-idæa,

and forms on the surface a hymenium of club-shaped basids each of which produces four basidiospores. The spores divide at maturity transversely into four cells, only the two end cells of which germinate, doubtless at the expense of the contents of the remaining two. The germ-tubes penetrate the epiderm of the leaf of the host, and a new mycele is formed which again bears basids. If, however, germination takes place elsewhere than on the proper host-plant, and conditions for the vegetation of the fungus be otherwise favourable, the germ-tube begins to sprout indefinitely by means of elongated *sprout-cells*, giving rise to others only at the ends. This condition has been maintained in nutrient solutions for a considerable time, but the sprout-cells have never been observed actually to give rise to a new mycele like the one produced by the basidiospores.

The TREMELLINEÆ (Tremella, Dill., Exidia, Fr.) present another simple type. They are gelatinous fungi of not very definite form, commonly of wavy outline, and are saprophytic on old and dead wood. The hymenia are formed on the surface of the gelatinous mass. The basids vary in appearance, and are usually provided with fine elongated sterigmata and reniform spores. Certain forms such as Sebacina (Tul.) and Hypochnus (Fr.) do not possess gelatinous membranes. The course of development is much the same as in Exobasidium. The germinating basidiospore gives rise under ordinary conditions to the compound sporophore again. Under other conditions, it has been observed in Dacryomyces, the germ-tubes do not grow to any great length, but produce *secondary spores*, or they form sprout-cells. The basidiospores of the same form divide transversely at maturity, usually into four cells, each of which may germinate. It should be added that on germination these secondary spores give rise to myceles. The hyphæ of such myceles, moreover, as well as those proceeding from basidiospores, sometimes give rise to tufts of rod-like cells, which in turn produce myceles. Similar phenomena have been observed in other Tremellineæ.



FIG. 317.—*Exidia spiculosa* Sommerf. Longitudinal section of portion of hymenium (much magnified). *s*, spores; *b*, basids; *h*, hyphæ of thallus. (After Tulasne.)

It is manifest therefore that in these simple types we have repeated very much the same order of things as in the tremelloid Uredineæ. It is perhaps most striking in the case of Exobasidium, from which the transition is easy to the Tremellineæ. The layer of basids and basidiospores may be compared with the layer of teleutospores, while the transverse division



FIG. 318.—*Coprinus stercorarius* Fr. *A*, *B*, and *C*, germinating spore in successive stages. *D* portion of mycele, *m*, with five early stages of development of fungus. *E* and *F*, further stages. *G*, longitudinal section through germinating sclerote, *s*, with young fungus still within volva, *v*. *H*, fully developed fungus with sclerote, *s*, and rhizoids, *r*. (*A*—*C* × 300, *D* × 200, *E* × 120, *F* × 50, *G* and *H* natural size.) (After Brefeld.)

of the basidiospores into four cells, two of which germinate, heightens the resemblance. Farther the production of secondary spores on the short germ-tubes of basidiospores recalls the formation of sporids on the

promyceles from teleutospores. From the Tremellineæ another easy step leads us on to the Thelephoreæ, and it may be borne in mind in this connection that certain Tremellineæ, as mentioned above, do not possess gelatinous membranes.

The THELEPHOREÆ (Corticium, Pers.) may be shortly described as recalling in point of simplicity of structure the teleutospore-layer of Uredineæ, while they approach very closely the club-shaped Hymenomycetes such as the Clavarieæ, in which the hymenium is disposed on the outer surface of erect club shaped cylindrical and often much-branched compound sporophores. Through a series of intermediate forms, the completeness of which may be recognised from a systematic study of the group, we proceed to the more perfect types of Hymenomycetes which possess sporophores of more complex structure.

In the higher forms of Hymenomycetes, the sporophore consists of a cap or *pileus* borne on the summit of a stalk or *stipe*. The mycele commonly vegetates in a soil rich in humus or on old wood or the like, and though usually of loose filamentous texture it is in

FIG. 319.—*Agaricus melleus* L., in different stages of development on branched rhizomorph-strands. The upper portion of rhizomorph represents that formerly known as *Rhizomorpha fragilis* Roth, while the lower strap-shaped portion is var. *subcorticalis*. (After Hartig.)

certain instances of more compact character. Such are the *sclerotes* which are resting states of Coprinus stercorarius (Fr.) (fig. 318), and the *rhizomorphs* of Agaricus melleus (L.) (fig. 319), composed of root-like branched strands of mycelial hyphæ, parasitic on the pine. The rhizomorphs are simply sclerotes with growing-points. From the mycele, of whatever character it be, there arises the *compound sporophore* by the continued apical or marginal growth of a bundle of hyphæ. It is not certain, but it may very well be, that intercalary growth also, in some

FIG. 320.—*Agaricus campestris* L. The common mushroom (natural size). Stages of development from *a* to *e*; *b* and *c* in section. (After Luerssen.)

cases at least, assists in the development. The hymenial surface, which is commonly situated on the under surface of the cap or pileus, is characterised in different genera by being spread over teeth-like projections (Hydnum, L.), radial plates in the numerous species of Agaricus (L.), concentric plates in the small genus Cyclomyces (Kze.), reticulated folds or pores (Polyporus, Mich. Boletus, L.); such typical characters being united by a wealth of intermediate forms. As a rule these projections are very symmetrical and of regular occurrence, and on them the chief generic characters are based in the classification of the group;

they are termed *gills* or *lamellæ* in the AGARICINI, *pores* or *tubuli* in POLYPOREÆ, and *teeth* in the HYDNEÆ. In many of the forms the hymenium is exposed from the first; in a series of others a membrane (*velum partiale*) connects the edge of the pileus all round with the stalk, and on its rupture by the extension of the pileus, part of it is left attached to the stalk, when it is termed the *annulus* or ring (fig. 320), though this does not occur in all cases. In a third series a membrane (*velum universale* or *volva*) (fig. 318) encloses the whole sporophore, pileus and stalk alike, and in the species belonging to Amanita, a sub-genus of Agaricus, both velum universale and velum partiale are present. In these latter cases, therefore, in which a membrane is present, the sporophore differs from the truly gymnocarpous forms. The development of Amanita is especially noteworthy, since the gills are not developed on the free inner surface of the pileus, but during an early condition from tissue common to both stalk and pileus.



FIG. 321.—*Coprinus stercorarius* Fr. Longitudinal section of the end of a gill in complete spore-bearing. *t*, trama; *p*, sterile palisade cells; *b*, basids with spores; *c*, cystids (× 300). (After Brefeld.)

Immediately beneath the hymenium is a layer of tissue called the *subhymenial layer*, distinguished from the rest of the tissue of the sporophore by the greater density of the ramifications of the hyphæ and by the more abundant protoplasmic contents.

FIG. 322.—*Polyporus igniarius* Fr. Transverse section of the under surface. *h*, the plexus of hyphæ forming the walls between the pores; *s*, the hymenium (× 270). (After Luerssen.)

The *trama* is that portion of the projection which bears the subhymenial layer, and consists of hyphæ running parallel to the surface from the insertion of the projection to its margin, which in many cases is uncovered by the hymenium.

The hymenium itself consists of parallel rows of club-shaped basids surmounted by sterigmata and basidiospores. The basids are the terminations of the subhymenial hyphæ, but the latter also frequently end in sterile cells, which are termed *paraphyses*, from the fact that they stand in the same relation to the basids as the paraphyses do to asci. Large inflated cells, often of relatively great dimensions, called *cystids*,



FIG. 323.—*Polyporus igniarius* Fr. Upper surface (half natural size). (After Luerssen.)

are frequently found emerging from the hymenial surface (fig. 321). They are very variable in form, club shaped, flask-shaped, cylindrical; pointed, hooked, or knob-shaped at the tip. They may be regarded as merely prominent hymenial hairs with the probable function of protecting the basids, or of parting the appressed lamellæ. They have been the subject of much idle speculation, and among other erroneous views they have been regarded as male organs.

In the sporophores of many Agaricini, notably of Lactarius (Fr.), *laticiferous hyphæ* occur, which yield considerable quantities of milky, generally acrid, juice when the tissue is bruised.

Sub-class 2.—Gasteromycetes.

The Gasteromycetes very closely resemble the Hymenomycetes in the essential points of the structure of the basids. At all events the agreement is close in this respect between the higher Hymenomycetes with cap and stalk, and the Hymenogastreæ, a section of the Gasteromycetes ; while other subordinate sections, such as the Lycoperdaceæ and Phalloideæ, diverge from the Hymenogastreæ only in minor points, as, it was seen, the lower Hymenomycetes do from the higher forms. The possession of a trama with a hymenial layer on either side of it may be here noted. In the external conformation, however, of the members of the group, a great variety is displayed, and, but for the existence of numerous intermediate forms, the group would appear to lack coherence in this respect, so great is the range of variation.

FIG. 324. *A*, *Octaviana asterosperma* Vitt., in section (× 5). (From Luerssen after Tulasne.)

The mycele is very frequently in the form of root-like strands, though there is no constancy in this respect, and the simple filamentous mycele occurs abundantly. The compound sporophores frequently grow to a great size in some of the sections, but the character which unites the whole is the possession of an investing membrane, the *peridium*, within which, and springing from it, are plates of tissue dividing the interior into chambers where the hymenium is produced.

At the outset there may be noted the remarkable genus Gautieria (Vitt.), which has no peridium. The peripheral chambers are therefore exposed on the free surface. The peridia of other forms vary considerably in thickness and other characters, and a tendency exists towards excessive thickening in the basal region, which develops outwards, forming a stalk in some instances, e.g. Lycoperdon (Tourn.) ; or inwards, in which case either a cushion-like body is produced, e.g. Hymenogaster (Vitt.), or a central column, e.g. Phalloideæ. The whole chambered structure is termed the *glebe*.

The HYMENOGASTREÆ may be regarded as an assemblage of the simplest forms of Gasteromycetes, possessing usually the simple structure indicated, but including among its members Gautieria without a peridium, and Secotium (Kze.), a genus with a central column traversing the body of the fungus. These two forms but heighten the resemblance which it has been remarked exists between the Hymenogastreæ and the Hymeno-

mycetes, the one being an approach to gymnocarpous forms, and the other noteworthy in respect of its stalked and pileate appearance. De Bary, in comparing the groups, says ('Comp. Morph. and Biol,' p. 337) : 'If we could attribute a decisive value to the habit of the plants, we should dwell upon the great resemblance between the stalked Hymenogastreæ, like Secotium erythrocephalum (Tul.), and a veiled Boletus. . . . But among the Polyporeæ there is a remarkable form Polyporus volvatus (Pk.), the Polyporus obvallatus (Berk. and Cooke), which considered by itself must be placed with or close to the Hymenogastreæ. Its sporophore, which lives in the bark of trees, is a hollow spherical body flattened at the poles and about the size of a hazel nut, with a thick closed wall of leathery texture ; its interior surface is



FIG. 325.—*Batarrea Steveni* Fr., longitudinal sections. *a*, a younger specimen, but with most of its spores ripe. *b*, a mature specimen, of which only apex and base are shown. *p* and *h*, the outer, *i*, the inner peridium ; *g*, the glebe (one-third natural size). (After de Bary.)

FIG. 326.—*Batarrea Steveni* Fr. Isolated threads of the capillitium (× 390). (After de Bary.)

covered with the hymenium of a Polyporus on the part next the substratum, and is sterile on the opposite side.'

In the LYCOPERDACEÆ (Puff-balls) the peridia are often developed to a colossal size, and in structure they agree in the main with the Hymenogastreæ. The chief distinction lies in the existence of two kinds of hyphæ in the trama ; slender segmented hyphæ with dense protoplasmic contents, the terminal members of which compose the hymenium, and stouter hyphæ running not only in the trama, but crossing the chambers. Eventually the slender hyphæ and the hymenium disappear, leaving only the stout hyphæ, now called the *capillitium*, and the masses of spores between. As examples of the possession of both inner and outer peridium in this section, there may be cited Geaster (Mich.), in which the outer one

becomes recurved after splitting longitudinally and acquiring a stellate aspect, and Batarrea (Pers.), which possesses an axile column immediately beneath the middle of the inner peridium. It develops into a stout stalk, which raises the closed inner peridium on its summit and ruptures the outer one, which now resembles in appearance the velum universale of Hymenomycetes. In Scleroderma (Pers.) the development of the glebe is intermediate between Hymenogastreæ and Lycoperdaceæ. While the trama is disorganised, and a portion persists as a fine network together with the masses of spores, it yet forms no true capillitium.

The NIDULARIEÆ, though very different in outward aspect from the



FIG. 327.—*Crucibulum vulgare* Tul. *A*—*C*, longitudinal section through ripening sporophores; stages of development in order of letters (slightly magnified). *D*, ripe sporophore in which the epiphragm is beginning to disappear (natural size). (After de Bary.)



FIG. 328.—*Crucibulum vulgare*. Section through upper part of sporophore of about same age as *B* in Fig. 327 (more highly magnified). *ap*, the outer, *ip*, the inner peridium; *rf* and *af*, its hairs; *n*, funiculus; *t*, the layer which forms a sheath round it, and belongs to a peridiolum divided through the middle. (After Sachs.)

other sections, are yet readily comparable with them. The chambers of the glebe possess very stout walls, and ultimately become separated from each other. The wall of the peridium becomes transformed into a gelatinous substance over the apical region, and on its disappearance the chambers of the glebe (peridiola) are left exposed in the interior of the bowl-shaped lower portion of the wall. Free and detached they resemble comparatively large sporanges. In Crucibulum (Tul.) a thin white membrane termed the *epiphragm* temporarily covers the summit (fig. 327).

The PHALLOIDEÆ are an assemblage of very remarkable and strange forms, in which the Basidiomycetes find their highest development.

Great variety of external conformation exists within the group, as the student will at once recognise on viewing such members of it as Phallus (L.), Hymenophallus (Nees ab Esenb.), Clathrus (Mich.), Ileodictyon (Tul.), Aseroe (La Bill.), &c.

Specimens of Phallus impudicus (L.) while yet enclosed within the peridium exhibit the following structure : The peridium consists of an outer white membrane and an inner white thinner one, and between these two a thick layer of tissue which has become gelatinous. Immediately within the inner membrane lies the glebe, situated in the upper capitate portion, and bounded on its inner surface by a conical membrane



FIG. 329. — *Mutinus caninus* Fr. Young sporophore. *m*, mycele ; stages of development in order of letters *u*—*y*. *y*, a specimen with ripe spores, but before elongation of stalk. *a*, the outer wall ; *i*, the inner ; *g*, gelatinous layer of peridium ; *b*, the basal portion ; *k*, the cone ; *s*, the stalk ; *gb*, the glebe (natural size). (After de Bary.)



FIG. 330.—A nearly mature specimen of *Phallus impudicus* L. before elongation of stalk, in longitudinal section. *m*, mycele ; *a*, outer, *i*, inner wall ; *g*, gelatinous layer ; *st*, stalk ; *k*, its cavity filled with mucilage ; *t*, lower margin of pileus ; *sp*, glebe ; *n*, the cup-shaped basal portion ; *x*, the spot where the peridium bursts (two-thirds natural size). (After Sachs.)

belonging to the central axis. This membrane gives off outwards into the glebe numerous walls arranged honeycomb fashion and dividing the glebe into compartments. The structure of the glebe itself recalls that of the Hymenogastreæ and Lycoperdaceæ. Below the glebe, and surrounding the base of the central axis, is a cup-shaped mass of comparatively firm tissue, in which the base of the stalk is fixed. It connects with the lower portion of the inner peridium, and sends a thin projection of tissue of less consistency upwards between the conical membrane and the stalk. The base rests on the outer layer of the peridium. The stalk itself is hollow at maturity, and is composed of air-containing

tissue, with numerous compartments. To scatter the spores the stalk elongates enormously, while it increases in thickness at the same time; the peridium bursts at the apex, and the glebe is separated from the inner peridial membrane and elevated on the summit of the stalk. When the spores are scattered, the conical membrane (so-called *pileus*) remains with the honeycomb-like structure on its outer surface attached to the apex of the spongy stalk.



FIG. 331.—*Aseroe rubra* Berk. Mature specimen. The peridium is attached below; the glebe is in the middle of the radiating expansion (half natural size). (From de Bary, after Berkeley.)

In Clathrus the development of peridium and glebe agrees with Phallus, but instead of a stalk a net-like structure serves to burst the peridium and elevate the glebe.

Such forms as Mitremyces (Nees ab Esenb.), Tulostoma (Pers.), Polysaccum (DC.), and Sphærobolus (Tode) exhibit other and remarkable types of development. Though they do not properly fall under any of the sections dealt with, they may be regarded as more or less divergent from the Lycoperdaceæ.

LITERATURE.

De Bary—Zur Kenntniss einiger Agaricinen (Bot. Zeit., 1859).

Bonorden—Beobacht. ü. d. Bau d. Agaricinen (Bot. Zeit., 1858).

Bonorden—Mycologische Beobacht. (Phallus, Sphærobolus) (*ibid.*, 1851).

Bonorden—Die Gattungen Lycoperdon u. Bovista, &c. (*ibid.*, 1857).

Brefeld—Bot. Untersuch. ü. Schimmelpilze, iii. (Leipzig, 1877).

Brefeld—For further development of Brefeld's views on the classification, &c., of Basidiomycetes (with much research on the lower forms) see Unters. aus dem Gesammtgebiete der Mykologie, Heft vii., 1888.

Corda—Icones Fungorum Prag (1837–1854).

Eidam—Die Keimung der Sporen u. d. Entstehung d. Fruchtkörper bei d. Nidularieen (Cohn's Beiträge, ii.).

Fischer—Zur Entwick. d. Gasteromyceten (Sphærobolus, Mitremyces) (Bot. Zeit., 1884).

Hartig—Wichtige Krankh. d. Waldbäume (Berlin, 1874).

Hartig—Die Zersetzungserscheinungen d. Holzes d. Nadelholzbäume u. d. Eiche (Berlin, 1878).

Hesse—Mikroskop. Unterscheidungsmerkmale d. typischen Lycoperdaceengenera (Pringsh., Jahrb. x.).

Hesse—Keimung d. Sporen von Cyathus striatus (*ibid.*, x.).

Murray—On the outer peridium of Broomeia (Journ. Linn. Soc. Bot., xx., 1884).
Murray—On two new species of Lentinus, one of them growing on a large sclerotium (Trans. Linn. Soc., 1886).
Nees v. Esenbeck—Plant. mycetoid., &c., Evolutio (Nova Acta Acad. Leop.-Carol., xvi.).
Pitra—Zur Kenntniss d. Sphærobolus stellatus (Bot. Zeit., 1870).
Rossmann—Beitr. zur Entwickl. d. Phallus impudicus (Bot. Zeit., 1853).
Sachs—Morph. d. Cucibulum vulgare Tul. (Bot. Zeit., 1855, p. 833). (Omitted from index to vol. of Bot. Zeit.)
Schlechtendal—Eine neue Phalloidee, nebst Bemerkungen ü. d. ganze Familie (Linnæa, 1862).
Schlechtendal u. Müller—Mitremyces Junghuhnii (Bot. Zeit., 1844).
Schmitz—Mycologische Beobachtungen, &c. (Linnæa, 1842).
Schmitz—Ueber Entw., Bau u. Wachstum von Thelephora sericea u. hirsuta (*ibid.*, 1843).
Schröter—Ueber d. Entwickl. u. d. systematische Stellung von Tulostoma (Cohn's Beiträge, ii.).
De Seynes—L'organisation des Champignons supérieurs (Ann. Sc. Nat., sér. 5, i.).
De Seynes—Rech. sur l. végétaux inférieurs, i. Des Fistulines (Paris, 1874).
Sorokin—Développement du Scleroderma verrucosum (Ann. Sc. Nat., sér. 6, iii.).
Tulasne—Carpologia, i. (Paris, 1861).
Tulasne—Fungi Hypogæi (Paris, 1851).
Tulasne—Obs. sur l'organisation des Tremellinées (Ann. Sc. Nat., sér. 3, xix.).
Tulasne—Nouvelles notes sur les fungi Tremellini et leurs alliés (*ibid.*, sér. 5, xv.).
Tulasne—De la fructification des Scleroderma comparée à celle des Lycoperdon et des Bovista (*ibid.*, sér. 2, xvii.).
Tulasne—Sur les genres Polysaccum et Geaster (*ibid.*, sér. 2, xviii.).
Tulasne—Rech. sur l'organ. d. Nidulariées (*ibid.*, sér. 3, i.).
Tulasne—Description d'une espèce nouvelle du genre Secotium (*ibid.*, sér. 3, iv.).
Tulasne—See Explor. Sc. d'Algérie, p. 434 (Clathrus).
Van Tieghem—Sur le développement du fruit, &c., des Basidiomycètes, &c. (Bull. Soc. Bot. France, 1876). (See also Bot. Zeit., 1876.)
Vittadini—Monographia Lycoperdineorum (Mem. Acad. Torino, v., 1842).
Woronin—Exobasidium Vaccinii (Ber. d. Naturf. Gesellsch. Freiburg, 1867).

Of historical interest is—

Micheli—Nova Plant. Genera, 1729 (Phallus, Clathrus).

The student is also referred for both morphological and systematic treatises to the numerous papers of Berkeley—to be found especially in the Ann. and Mag. of Nat. Hist. and Hooker's Journal of Botany—and to his separate books; for systematic information particularly to the works of Fries, Persoon, Hoffmann (Icones Analyticæ Fungorum, Giessen, 1861–65), Saccardo, Sylloge, vols. v.-vii.; and for British forms to Cooke's Handbook of British Fungi, 1871, and to Stevenson's Hymenomycetes Britannici, 1886.

SIXTH SUBDIVISION.

MYCETOZOA.

The Mycetozoa are a group of organisms separated by a great gulf from the Thallophytes, but presenting certain points of resemblance to the Fungi which may here be indicated, while the amount of that resemblance and the degree of their divergence will be more fittingly estimated at the end of this chapter. Their nutrition is saprophytic, and the organs of reproduction are sufficiently like those of the Fungi to justify the use of the terms *sporange*, *spore*, *swarm-spore*. The vegetative body, on the other hand, differs in structure *toto cœlo* from any form of thallus. It consists of a naked protoplasmic body, either a *plasmode* formed by the coalescence of peculiar swarm-spores, or an aggregation of such swarm-spores. The first case is characteristic of one class, the Myxomycetes; the second case of the other smaller class, the Acrasieæ.

Class XXIII.—Myxomycetes.

The ripe spores of Myxomycetes are capable of germination at once, and many of them retain this power for considerable periods, some for as long as several years. Most germinate at the ordinary spring or summer temperature, and in pure water, while others require a nutrient solution. The germination of the spores of Cribrarieæ and Tubulinæ has not been observed, and the failure of the attempts to procure it may be owing either to the supply of unsuitable media or to a necessity for a period of rest—more likely the former. In structure the spores resemble those of fungi, as has been said. The wall is either smooth or sculptured on the outer surface, and the protoplasm contains one, sometimes two, nuclei. The act of germination consists of the emission of a swarm-spore. The membrane opens and the protoplasm escapes with a creeping motion. This naked protoplasmic body or swarm-spore then exhibits amœboid movements, protruding and withdrawing irregular processes, becoming more or less elongated, and

acquiring a cilium at the end of a finely pointed process. Its movements are of two kinds : a hopping movement, during which it commonly rotates round its longitudinal axis, while the outline undulates ; and a creeping movement, which takes place on a firm substratum with the cilium in advance. The creeping is also sometimes accomplished by the protrusion and retraction of *pseudopodes*. The same swarm-spore often moves both by hopping and by creeping alternately. After passing through this stage, during which swarm-spores multiply by simple division into two (such division taking place in some cases even before leaving the spore), the formation of plasmodes begins. The swarm-spores taking part in this process are such as have withdrawn their cilia and exhibit creeping amœboid movements. Several come into contact and coalesce, thus forming the beginning of the plasmode. Others are drawn towards it—how, no one has ever found out—and successively coalesce with it, until a comparatively large plasmode is formed with the appearance and movements of a huge amœboid swarm-spore without cilia. This plasmode



FIG. 332.—*Chondrioderma difforme* Rost. 1, a ripe spore; 2, the same germinating; 3–5, swarm-spores; 6, 7, the same in amœboid state; 8, two in close contact; 9, the same coalesced; 10, three in contact; 11, two after coalescence, the third still free; 12, young plasmode which has taken up two spores into its substance (× 350). (From Sachs, after Cienkowski.)

FIG. 333.—*Didymium serpula* Fr. *A* and *B*, plasmodes (natural size). *C*, margin of a moving plasmode (× about 200). (After Reinke.)

nourishes itself and grows, acquiring, in the case of some Physareæ, great dimensions, and forming reticulated masses which may be measured by inches. Fuligo varians (Somm.) (or as it is more commonly called Æthalium septicum (Fr.) or 'flowers of tan,' from its appearing during summer on tan) is such a body, but the plasmodes of other families of Myxomycetes, as well as of some Physareæ, generally remain very small in comparison with this. The appearance of the strands or branches of the plasmode (under the microscope) is that of a turbid granular mass bordered by a clearer hyaloplasm. The surface of the plasmode of Physareæ is invested with a soft shiny envelope of a substance different from protoplasm. The plasmodes of certain other forms are similarly invested with envelopes, as to the nature of which not much is known. The larger portion of the granules contained in the plasmodes of Physareæ are of calcium carbonate; granules contained in other plasmodes require investigation. Nuclei are abundantly present. Many foreign bodies such as spores, diatoms, &c., are often found included in plasmodes. Constant movement is maintained, and the most characteristic is that of the protrusion and retraction of pseudopodes. Since protrusion is commonly more active on one side than on the other, an advancing movement of the whole is thus brought about. Internal streamings, more or less copious, answer to the amœboid movements. The external causes of movements are: with reference to (1) *illumination*, they are negatively heliotropic; (2) *water*—they are positively hydrotropic, i.e., when not about to form spores they leave comparatively dry spots and move towards moist places; (3) *food*—they are positively trophotropic, i.e., they move towards nutrient substances (as might be expected); (4) *heat*—within certain limits they move towards the warmer side of a surface unequally warmed. These movements are without reference to the direction in space in which they may have to be made. It may be stated here that the process of nutrition takes place only in the amœboid states—the swarm-cell and plasmode.



FIG. 334.—*Stemonitis fusca* Roth. *A*, sporange (natural size). *B*, capillitium (× about 100). (After Reinke.)

Resting states may occur at all motile stages of the life-history. *Microcysts* are the resting states of swarm-spores. They round themselves off, and are invested with a delicate membrane or only with a firm border. Young plasmodes similarly form thicker-walled cysts, and mature plasmodes form multicellular bodies—*sclerotes*.

The *spores* of Myxomycetes are formed either endogenously within *sporanges*, or on the free surface of *sporophores* (Ceratieæ). Sporanges are formed either by the whole plasmode becoming one, or the plasmode divides into portions, each of which becomes a sporange. Such as are situated on stalks begin as small swellings on a strand of the plasmode, and by degrees acquire their mature form as the protoplasm ascends into



FIG. 335.—*a*, *Ceratium hydnoides* Alb. and Sch. Piece of sporophore in act of forming. *b*, *Ceratium porioides* Alb. and Sch. Piece of the margin of a sporophore; spore-formation beginning; two spores which subsequently become slightly ellipsoid on their stalks. (*a* × about 68, *b* × 120.) (After Famintzin and Woronin.)

them. While this process of formation goes on the solid contents of the plasmode are expelled. The interior of the mature sporange is either filled with spores only, or more commonly there is also present a *capillitium* consisting of numerous filaments traversing the cavity in all directions. They probably serve as supports to the wall of the sporange in the first instance, and may further be connected with its rupture and the dispersal of the spores.

There are only two known species of Ceratium (Link), a genus which forms free spores, i.e. not within a sporange. In this case the plasmode before spore-formation consists of a network of innumerable branches from which cylindrical processes arise. The whole protoplasm

flows into these processes and finally breaks up into numerous polyhedral portions. Each of these portions grows outward into the form of a ball connected with the surface by a short narrow stalk. This sphere acquires a wall, and the process of spore-formation is completed.

Class XXIV.—Acrasieæ.

So far as is known the spores of Acrasieæ germinate only in nutrient solutions. The swarm-spores are never ciliated, and move only by creeping in amœboid fashion. Under unfavourable conditions they encyst themselves, and form temporary resting states. They unite in great numbers for the purpose of forming spores again, but the union never amounts to *coalescence* into plasmodes. They are heaped together as it were, and compose bodies of more or less definite form. In this condition each swarm-spore becomes invested with a thin membrane, though no common sporangial wall is formed. Guttulina (Cienk.) forms simple spore-heaps, but in Dictyostelium (Bref.) and Acrasis (Van Tiegh.) a stalk is formed by the swarm-spores in the centre of the mass becoming transformed into series of cells with firm walls, and up it the rest of the swarm-spores climb and form spores at the top.

Doubtful Mycetozoa.

De Bary repudiates the attempt made by Zopf to bring together under this group an ill-assorted assemblage of lower organisms exhibiting amœboid movements. He considers such forms as Bursulla (Sorok.), Protomyxa (Hæck.), Vampyrella (Cienk.), Nuclearia (Cienk.), Monas amyli (Cienk.), Monadopsis (Klein), Pseudospora (Cienk.), Colpodella (Cienk.), and Plasmodiophora (Woron.) to be doubtful Mycetozoa. Plasmodiophora Brassicæ (Woron.), which is parasitic on the roots of Cruciferæ, on which it produces large swellings, is common. The ciliated swarm-spores penetrate into the parenchymatous tissue of such roots. The cells affected swell to a great size, and large amœboids appear in them, but it is not certain whether these are single swarm-spores or small plasmodes formed by the coalescence of several. The whole protoplasmic contents of a cell then break up into spores.

Affinities.

De Bary, to whose remarkable investigations we owe the bulk of our knowledge of the Mycetozoa, considers that the group differs

distinctly from the Fungi (and still more from other plants) 'in all such characteristics as do not belong to all organisms alike. . . . The difference would not be less decided, if the Mycetozoa were without their remarkable amœboid movements, for such movements are observed in other vegetable cells which have not a firm membrane. The characteristic mark of separation lies in the formation of plasmodes or aggregation of swarm-cells' ('Comp. Morph.,' p. 443). He farther remarks that the highest forms of the group give no evidence of close affinity with yet higher organisms, and seeks for their relationship with Amœba. Guttulina, he points out, differs from such forms only by the aggregation of its spores. Guttulina protea (Fay.) (Copromyxa protea, Zopf) even forms solitary spores. This form then links the Amœbæ with the more highly differentiated Acrasieæ, and these connect with the Myxomycetes. Taking together this connection with the animal kingdom, and the want of connection on the other hand with Fungi (to which they have a merely superficial resemblance) or other plants, we are justified in placing them, as de Bary does, 'outside the limits of the vegetable kingdom.'

Literature of Mycetozoa.

Baranetzki—Influence de la lumière sur les Plasmodia des Myxomycètes (Mém. Sc. Nat. Cherbourg, xix., p. 321).

De Bary—Die Mycetozoen (Zeitsch. f. wissensch. Zoologie, Bd. x., 1859; and 2nd edition, Leipzig, 1864).

Brefeld—Dictyostelium mucoroides (Abh. d. Senckenberg. Naturf. Gesellsch., vii.).

Brefeld—Untersuch. aus der Gesammtgebiete der Mykologie (Leipzig, 1884).

Cienkowski—Zur Entwickelungsgesch. d. Myxomyceten (Pringsh. Jahrb. wiss. Bot., iii.).

Cienkowski—Das Plasmodium (*ibid.*, iii.).

Cienkowski—Ueber einige protoplasmatische Organismen (Guttulina). See Just's Jahresber. for 1873, p. 61.

Cienkowski—Beitr. zur Kennt. der Monaden (Schultze's Arch. f. micros. Anat., i.).

Lister—Plasmode of Badhamia and Brefeldia (Ann. of Bot., ii., 1888, p. 1).

Rostafinski—Versuch eines Systems der Mycetozoen (Dissertat. Strassburg, 1873).

Rostafinski—Slucowce (Mycetozoa) (Paris, 1875). A monograph admirably illustrated, but written in Polish. The system in Cooke's Myxomycetes of Great Britain (London, 1877) is adapted from Rostafinski's monograph.

Stahl—Zur Biologie der Myxomyceten (Bot. Zeit., 1884).

Strasburger—Wirkung d. Lichtes und d. Wärme auf Schwärmsporen (Jena, 1878, p. 69).

Strasburger—Zur Entwickgesch. d. Sporangien v. Trichia fallax (Bot. Zeit., 1884).

Van Tieghem—Sur quelques Myxomycètes à plasmode agrégé (Bull. Soc. Bot. de France, 1880, p. 317).

Zopf—Die Schleimpilze, in Schenk's Handbuch der Botanik, iii. (1887).

For further literature see De Bary's Comp. Morph., p. 453, and Rostafinski's monograph.

SEVENTH SUBDIVISION.

PROTOPHYTA.

Whether the Protophyta should be reckoned as a distinct subdivision from the Algæ, or only as the lowest members of that great series, is a question rather of convenience than of principle. In an ideal system of classification founded exclusively on genetic affinities, those organisms would be regarded as 'protophytes' which were the earliest heralds of the appearance of vegetable life on the surface of the globe. But, from the structure and conditions of life of such organisms, it is impossible that they can have been preserved to us in the fossil state, and it is only from the comparative simplicity or complexity in the structure of an organism that we can conjecture whether it is an archaic or a derivative form. And here, as was remarked in the Introduction, we are extremely liable to be misled if we neglect to take into account the phenomenon of the constant appearance of degeneration or retrogression in the vegetable kingdom. An organism may be simple in its structure either because it has never risen, through countless ages, above the simplicity of its primeval ancestors, or because it has fallen back from a more complicated condition. The object of the scientific systematist should be to separate, so far as possible, between these two sets of organisms, to include the former among his lowest class of protophytes, and to relegate the latter in each case to the class from which they have degenerated. But this task is attended with great difficulties, and is often well-nigh impossible. An organism may display degeneration of one set of organs, while another set manifest no such degeneration and have even continued to develop. We may take it indeed as a general law that wherever you have either the vegetative or the reproductive organs strongly developed, while the other set are very feeble or altogether wanting, you have *primâ facie* evidence of retrogression. But, on the other hand, degeneration may take effect in all the organs of a plant, leading to retrogression in all lines towards, it may be, the archaic form. As knowledge advances, the constant tendency will be to transfer to this class of retrogressive members of higher families forms previously regarded as protophytal.

While the Schizophyceæ or chlorophyllous Protophyta approach very closely to the lowest forms of Algæ, the Schizomycetes or non-chlorophyllous Protophyta exhibit greater affinities, as de Bary has shown, with the chlorophyllous forms than with any family of Fungi. Grouping, therefore, all these lowest forms of vegetable life, whether containing chlorophyll or not, into a single subdivision of Cryptogams, it will be most convenient to discuss them under two heads, as distinct and to a certain extent parallel series.

GROUP I.—SCHIZOPHYCEÆ.

An attempt is here made to bring together those chlorophyllous forms which, in the present state of our knowledge, we must regard as primordial; while others, almost equally simple in structure, have been referred to the classes of which they appear to be retrogressive members. The group now under consideration comprises the greater number of the forms of vegetable life which are unicellular, which display no true process of sexual reproduction, and which contain chlorophyll.

Limited in this sense, the Schizophyceæ may be divided into three well-marked classes, the *Protococcoideæ*, the *Diatomaceæ*, and the *Cyanophyceæ*. In the Cyanophyceæ are included those forms in which the pure green colour of the chlorophyll is masked by a blue-green pigment dissolved in the cell-sap, an arrangement not found except in plants of the very simplest structure. The position of the Diatoms has been a subject of much controversy among systematists. They display in some respects a similarity to the Desmids; but, for reasons given below, we are disposed to consider this resemblance as apparent rather than real, and to regard the Diatomaceæ, not as a family derived from the Desmidiaceæ by retrogression, but as a primordial type of great simplicity of structure. In the Protococcoideæ are included those forms in which the pure-green of the chlorophyll is not concealed by the blue-green colouring matter of the Cyanophyceæ, nor by the brown colouring matter of the Diatoms. It is unquestionably from them that all the higher forms of vegetable life have been derived, and the boundary line between the Protococcoideæ and the lower forms of Algæ is one that cannot be accurately laid down.

Literature.

The literature of the Schizophyceæ is included under that of Algæ, or in the works specially named when treating of the separate classes and orders.

Class XXV.—Protococcoideæ.

In this class, the Chlorophyllophyceæ of some writers, are included those simplest forms of vegetable life in which the endochrome consists of pure chlorophyll of its natural green colour, sometimes replaced, to a larger or smaller extent, by a red pigment, but the cell-sap never pervaded, as in the Cyanophyceæ, by a soluble blue colouring-matter. The individuals are of microscopic size, and may be either motile or resting, and very commonly the same species occurs in both conditions. The motile or *protococcus* form is, in the lower members, strictly unicellular, consisting of chlorophyllous protoplasm either naked or invested with a very delicate coat of cellulose or of a carbohydrate nearly allied to cellulose, usually developing but little or no mucilage, and moving freely through the water by means of a pulsating vacuole and two vibratile cilia. In the resting condition the individuals are invested by a much thicker cell-wall, and have a tendency to congregate or coalesce into *palmelloid* families, and to enclose themselves in a common gelatinous envelope. In this state they multiply rapidly by repeated bipartition. The palmelloid form may be derived directly from the protococcoid, the protococcus-cells coming to rest, losing their cilia, and investing themselves with a thicker cell-wall of cellulose; or, in the higher members, the individual consists of a number of *gonids*, chlorophyllous masses of protoplasm, enclosed in a common watery hyaline envelope of mucilage, and propagation takes place by the escape of these gonids from the envelope in the form of naked biciliated *zoospores* or swarm-spores, closely resembling protococcus-cells, which, after going through a motile period, come to rest, lose their cilia, invest themselves with a coat of cellulose, and multiply by repeated bipartition in the palmelloid form. In some cases these swarm-spores are of two kinds, the smaller ones being conjugating *zoogametes*. In no case is the individual filiform and divided by transverse septa, as in the higher families of the Cyanophyceæ.

It cannot be too strongly insisted on that this class is a purely provisional one. Many of the forms at present included in it are, in all probability, nothing but stages in the development of algæ of considerably greater complexity of structure belonging to widely separated families. The external resemblance between the Protococcaceæ and the Chroococcaceæ, and the parallel series of forms in these two families, does not probably represent any genetic affinity. There is, on the other hand, an undoubted alliance with the Pandorineæ, through Chlamydococcus and Chlamydomonas, as well as with the Hydrodictyeæ and

Siphoneæ through intermediate forms. The Protococcaceæ converge also on the boundary line between the vegetable and animal kingdoms: and, since it has been demonstrated that the power of forming chlorophyll and starch is not of itself sufficient to determine an organism to belong to the vegetable kingdom, it is impossible to draw a hard and fast line between the Protococcaceæ and the Flagellate Infusoria, with which they are connected by such forms as Euglena and the Peridinieæ.

The Protococcoideæ are divided into two orders, the boundaries of which are very ill-defined: the *Eremobiæ* and the *Protococcaceæ*.

LITERATURE.

Ehrenberg—Die Infusionsthierchen, 1838.

Nägeli—Neuern Algensysteme, 1847, pp. 123–132; and Gattungen einzelliger Algen, 1849.

Braun—Verjüngung in der Natur, 1851 (Ray Soc. Bot. and Phys. Memoirs, 1853); and Algarum unicellularum genera, 1855.

(Also the Memoirs referred to under the separate genera, and the literature of Algæ generally.)

ORDER I.—EREMOBIÆ (including SCIADIACEÆ).

In this ill-defined family, known by some writers as Characiaceæ, the limits of which are very difficult to assign, are included a number of genera distinguished from the Protococcaceæ by their greater complexity of structure. They are mostly fresh-water, but comprise also a few marine organisms, free-swimming or attached to algæ. In the larger number of genera each individual consists of a number of green protoplasmic bodies, *pseudocysts* or *gonids*—that is, masses of chlorophyllous protoplasm of defined outline but not clothed with a definite cell-wall of cellulose—sometimes of considerable size, enclosed in a common transparent hyaline envelope, which may be simple or may branch in an arborescent manner. In some genera the hyaline envelope is wanting. Multiplication takes place by simple division, or by the transformation of the gonids into zoospores, which sometimes display a differentiation into larger *megazoospores* and smaller *microzoospores* or *zoogametes*. Although conjugation of these gametes has hitherto been observed only in a few cases, this appears to be the earliest indication among chlorophyllous organisms of a differentiation of sexual elements; and the Eremobiæ clearly approach those algæ which multiply by conjugation through Botrydium, or through such forms as Endosphæra, Chlorochytrium, and Phyllobium, or again through Hydrodictyon. Lagerheim (Ber. Deutsch. Bot. Ges., 1884, p. 302) asserts the presence of chromatophores in Glaucocystis (Itz.).

In the following paragraphs only the more remarkable or better known genera are described.

In *Sciadium* A. Br., made by some writers the type of a distinct family SCIADIACEÆ, the peculiar mode of germination of the zoospores gives rise to a remarkably complicated structure. Each individual consists at first of a single elongated cylindrical cell. The green protoplasmic contents of this cell break up ultimately into a number of biciliated zoospores, which are set free by the upper portion of the cell-wall becoming detached in the form of a cap. The zoospores do not, however, escape, but germinate while still attached to the mother-cell, giving rise to a cluster of smaller cylindrical cells springing from the apex of the mother-cell. This may go on until the colony consists of as many as four generations, giving the appearance of a minute branching shrub. The zoospores of each generation are smaller than those of the preceding one, and it is probable that those of the last generation, which escape altogether from the parent-cell, are conjugating zoogametes. It is possible that a form allied to Sciadium may have been the starting-point of the Siphonocladaceæ, with which family it shows a certain affinity, as, for example, with Valonia.



FIG. 336.—*Sciadium arbuscula* A. Br. (magnified).

Chlorothecium Bzi. (Malpighia, 1888, p. 250) occurs in the form of palmelloid colonies with a thick and firm cell-wall on aquatic plants. From the cells of these colonies are developed zoosporanges, or rather *gametanges*, without any alteration of their primitive form; from each gametange there escape from two to four swarm-spores, or occasionally only one, each provided with a single cilium and a conspicuous red pigment-spot. These swarm-spores are *zoogametes*, conjugating by gradual fusion. After hibernating the contents of the *zygosperm* break up into two masses, each of which escapes as a non-sexual *zoospore*, so that the zygosperm is itself a *zoosporange*. From these zoospores are again formed the palmelloid colonies, in which form Chlorothecium may multiply itself non-sexually without producing zoogametes.

The position of *Halosphæra* (Schmitz, Mittheil. Zool. Stat. Neapel, 1878, p. 61) is very doubtful. Each individual is a minute green globe, just visible to the naked eye, as much as 0·5 mm. in diameter, floating on the surface of the sea, and bearing an external resemblance to Volvox.

Each cell contains a nucleus and a vacuole; the green protoplasmic contents break up ultimately into zoospores of a very peculiar form, conical, with two cilia attached to the nearly flat base, recalling those of Hydrurus (p. 256).



FIG. 337. *Haplosphæra viridis* Schm. Globe (× 80), and zoospore (× 150). (After Schmitz.)

In *Dictyosphærium* Näg., which ought possibly to be placed under the Cœnobieæ, the free-swimming colony is composed of globular or kidney-shaped green gonids connected together by delicate threads of mucilage. New colonies are formed by repeated bipartition of the gonids, which frequently exist for a time without any enclosing cell-wall. *Mischococcus* Näg. consists of minute globular gonids connected together in an arborescent manner and enclosed in a hyaline envelope, the whole colony attached to fresh-water algæ. Borzì (Malpighia, 1888, p. 133) describes also a palmelloid form of Mischococcus, the cells of which give birth to megazoospores with only a single cilium. The dendroidal form may spring either from these zoospores or directly from the palmella-cells; its cells also produce uniciliated swarm-spores, similar to the zoospores but smaller. They are apparently zoogametes conjugating to produce a biciliated zoosperm. *Botrydina* Bréb., found on moist ground, trunks of trees, &c., is composed of a number of minute gonids enclosed in a pear-shaped or globular hyaline envelope, as much as 0·1 mm. in diameter, and resembling Aphanocapsa among the Chroococcaceæ. It may possibly be allied to Botrydium.

FIG. 338.—*Dictyosphærium reniforme* Buln. (× 400). (After Cooke.)



FIG. 339.—*Mischococcus confervicola* Näg. (× 400). (After Cooke.)

Characium A. Br. is a minute green organism attached by a gelatinous stalk to algæ or other fresh-water plants, often in groups. It is

ovate or pear-shaped, 0·02-0·025 mm. in diameter in the larger species, often apiculate or spinous at the apex. The cell-contents divide, by successive bipartitions, into zoospores, which commence swarming while still within the mother-cell, indicating an approach to Hydrodictyon. They escape through a lateral or terminal fissure. Nearly allied to Characium are *Hydrocytium* A. Br., also met with in fresh water, and *Hydrianum* Rabh., found in similar localities. In the last genus the zoospores also escape at the apex. In *Apiocystis* Näg. a large number of gonids are sparsely scattered through a stalked pear-shaped gelatinous envelope attached to fresh-water algæ. They occur chiefly in the periphery, and are ultimately converted into zoospores.

FIG. 340.—*Characium ornithocephalum* A. Br. (× 600). (After A. Braun.)

Codiolum A. Br. is a club-shaped marine organism, about 0·04 mm. in diameter, and four to six times the length, attached to rocks or sea-weeds. It is propagated by zoospores, or, according to some observers, also by resting hypnospores. *Hauckia* Bzì. (Nuov. Giorn. Bot. Ital., 1880, p. 290) grows on rocks exposed to the sea. The gonids are placed in pairs on a long hyaline stalk; it produces zoospores of two different sizes, but no process of conjugation has at present been observed.

FIG. 341.—*Apiocystis Brauniana* Näg. (× 100). (From nature.)

FIG. 342.—*Codiolum gregarium* A. Br. (magnified). (After Hauck.)

Sykidion Wright (Trans. Irish Acad., 1881, p. 27) is also a marine genus, allied to Characium and Hydrocytium; but the zoospores escape through a terminal instead of a lateral fissure.

Of free-swimming forms occurring in fresh water, *Nephrocytium* Näg. consists of kidney-shaped gonids enclosed in a hyaline envelope. Although the production of zoospores has not been detected in this genus, its position is probably here, though its true place may possibly be among the Sorastreæ. Dangeard (Bull. Soc. Linn. Normandie, i., 1888, p. 196) has observed a mode of propagation by the formation of daughter-colonies within the membrane of the parent-colony. In *Ophiocytium* Näg., the originally cylindrical individual becomes curved in a serpentine manner, and produced at one extremity into a hyaline spine. The zoospores escape by the detachment of the cap-like apex of the hyaline envelope.

FIG. 343.—*Nephrocytium Nägelii* Grün. (× 300). (After Cooke.)

In *Hormospora* Bréb. the free-swimming individual or colony consists of a very elongated straight or bent cylinder, sometimes branching, the gonids arranged in a single or double row within a dense hyaline envelope. No formation of zoospores has been observed. Cylindrocapsa (Reinsch) (see p. 227) is placed here by some authorities.

FIG. 344.—*Hormospora mutabilis* Bréb. (× 200). (From nature.)

Although in the majority of the genera named above only one kind of swarm-spore has hitherto been observed, it is highly probable that some or all of them produce both megazoospores and zoogametes with a sexual function.

LITERATURE.

Fresenius—Abhandl. Senckenberg. Naturf. Gesell., iii., 1856–8, p. 237.
Archer—Microscop. Journ., 1866.
Zukal—Oesterr. Bot. Zeitschr., 1880, p. 11.
Holmes—Journ. Linn. Soc., xviii., 1881, p. 132.
Borzì—Nuov. Giorn. Bot. Ital., 1882, p. 272; and Studi Algologici, 1883.
Lagerheim—Bot. Centralbl., xii., 1882, p. 33; and Oefv. Vetensk. Akad. Förhandl., Stockholm, 1885, p. 21.
Klebs—Unters. Bot. Inst. Tübingen, i., 1883, p. 233.
Bennett—Journ. Micr. Soc., 1887, p. 9; and 1888, p. 2.

Order 2.—Protococcaceæ (including Palmellaceæ).

In this family are included a number of organisms of very simple structure, many of which occur both in the free-swimming (*protococcus*) and in the resting (*palmella*) condition. In the former state they bear a very close resemblance to the zoospores of the higher algæ. Other forms are known in one condition only, in which they have a free swimming motion without the aid of cilia.

Protococcus Ag. is one of the commonest objects in fresh water, especially stagnant rain-water, forming masses of a bright green colour, either floating free or attached to a submerged or floating object, but destitute in this state of any spontaneous power of motion. In this palmella-condition each individual consists of a nearly spherical cell, varying between forty and fifty microns (= ·04–·05 mm.) in diameter, which multiplies rapidly by repeated bipartition of its contents. The bright green endochrome has usually intermixed with it a larger or smaller quantity of a red pigment, the proportion varying according to the conditions of life, &c. The change to the active condition takes place in the following way. The protoplasm withdraws itself from the cell-wall, and escapes in the form of an ovoid mass provided with two very long and slender vibratile cilia and a pulsating vacuole, by the agency of which it is driven rapidly through the water. The pulsation of this vacuole has been explained by the alternate absorption from the water, through the agency of the chlorophyll, of carbon dioxide, and the expulsion of free oxygen resulting from the process of assimilation. In some cases the contents of the mother-cell do not escape as a single zoospore, but break up before escaping into eight or more smaller zoospores. The motile protococcus may be either entirely without cell-wall of cellulose, or may have a very delicate one, through orifices in which the protoplasmic cilia protrude. Some observers state that there are two kinds of zoospore in Protococcus—microzoospores and megazoospores, and that conjugation takes place between the latter; but this last statement at all events requires confirmation. After swimming about rapidly for a time in all directions with an apparently spontaneous movement, the motile protococcus comes to rest, loses its cilia, becomes *encysted*, or invested with a thick cell-wall of cellulose, and again enters the palmella-



Fig. 345. *Protococcus pluvialis* Ktz. *A*, motile condition; *B*, palmella condition (× 250). (After Cohn.)

condition in the form of *resting-spores*, which may become dried up and retain their vitality for years as a dry powder, resuming their activity when again placed in water. McNab (Ann. & Mag. Nat. Hist., 1883, p. 124) has, by the use of osmic acid and carmine, detected a nucleus in the free ciliated state of Protococcus, and also in individuals in which cell-division is going on.

When in an active condition in sunlight, Protococcus gives off into the surrounding water large quantities of oxygen, the result of the activity of its chlorophyll, thus contributing to render it habitable for animal life. The amount of the red pigment varies greatly. It is often confined to a small spot near to the point of attachment of the cilia, the 'pigment-spot,' bearing a close resemblance to the 'eye-spot' of the Flagellate Infusoria. If present in larger quantities, so as to give a red tint to the entire organism, this is known as Hæmatococcus (Ag.). In the palmella-condition this form frequently presents the structure and appearance of a blood-red incrustation on rocks and stones, when it has been described as Palmella cruenta (Ag.) and Porphyridium cruentum (Näg.). Very closely allied are the Palmella prodigiosa (Mont.) (Monas prodigiosa, Ehrb.), which forms blood-red spots on bread, potatoes, &c., and the Palmella nivalis (Hook.) (Protococcus nivalis, Ag., Chlamydococcus nivalis, A. Br.), which, under the name of 'red snow,' frequently covers large tracts of snow in arctic and alpine regions in a very short time. Phipson (Compt. Rend., lxxxix., 1879, pp 316, 1078) has examined the red colouring-matter of Palmella cruenta, and finds it to consist of minute globules about four microns (·004 mm.) in diameter, closely resembling those of the hæmaglobin of blood, but somewhat smaller. He proposes for the pigment the name *palmellin*. It is soluble in water, but insoluble in alcohol, ether, and carbon bisulphide. Like hæmaglobin it contains traces of iron. Other lowly-organised snow and ice plants besides Palmella nivalis are brightly coloured, and appear to perform an important function in melting the snow by their strong absorption of the rays of heat. In addition to palmellin, Palmella contains also xanthophyll, and a small quantity of another substance of the nature of camphor and possessing a marshy odour, which Phipson calls *characin*, and which is present in other terrestrial and fresh-water algæ, and especially in Chara. In the hæmatococcus-condition it is sometimes impossible to detect directly the presence of chlorophyll ; but experiments by Engelmann (Rev. Internat. Sci. Biol., 1882, p. 468, and Bot. Zeit., 1882, p. 663) seem to show that there is always a certain amount of chlorophyll present, though it is possible that the power which Hæmatococcus undoubtedly has of decomposing carbon dioxide may be due to the presence of other substances allied to chlorophyll, but differing from it in colour.

It is impossible to distinguish between the genera Protococcus (Ag.), Pleurococcus (Meneg.), and Palmella (Lyngb.); but it is doubtful whether Chlamydococcus (A. Br.) and Chlamydomonas (Ehrb.), which undergo much more complicated changes of form, and in some conditions very closely resemble Protococcus, have been rightly identified with it (see p. 299). Hæmatococcus Butschlii (Blockmann, Ber. Heidelberg Naturh. Ver., 1886) probably belongs to Chlamydomonas. Schnetzler



FIG. 346.—*Glochiococcus anglicus* Benn. (× 200). (From nature.)



A B

FIG. 347.—*Chlorococcum gigas* Grün. (× 300). (After Cooke.)

(Bull. Soc. Vaud. Sc. Nat., 1882, p. 115) regards Palmella uvæformis (Ktz.) as a stage in the development of a Stigeoclonium; while Andersson identifies it with Draparnaldia.

Glochiococcus (Lagerh.) (Acanthococcus, Reinsch, Ber. Deutsch. Bot. Ges., 1886, p. 237) differs from Palmella in the cell-wall, which is thick and lamellated, being in most of the species furnished with warts, spines, or other prominences. The cells, which closely resemble the zygosperms of desmids, divide into eight or sixteen daughter-cells, which remain but a short time in connection, being set free by the deliquescence of the outer membrane.

Chlorococcum (Fr.) is analogous to Chroococcus among the Chroococcaceæ. Several species are common in pools or on moist walls or rocks. In C. gigas (Grün.) the cells are as much as 0·01–0·015 mm. in diameter, and either a single cell or a colony of cells is enclosed in a very thick lamellated hyaline envelope. In *Glœocystis* (Näg.), corresponding to Glœocapsa among the Cyanophyceæ, the cells are associated in families of two, four, or eight, each family being enclosed in a lamellated gelatinous envelope, in addition to the similar envelope which encloses each cell. In *Schizochlamys* (A. Br.) the cells escape from the surrounding envelope by the latter splitting into two or four equal parts. *Eremosphæra* (de By.) is a beautiful bright green globe, 0·1–0·15 mm. in diameter, floating free in bog-pools, and enclosed in a thin hyaline envelope.

FIG. 348.—*Schizochlamys gelatinosa* A. Br. (× 600).

Botryococcus Ktz. consists of mulberry-like masses of thick-walled cells united together into colonies, with no investing membrane, or only a very slight one; it is found in bog-pools, and is endowed with a rotating as well as a free-swimming motion. It has possibly a genetic affinity with the Cœnobieæ.



FIG. 349.—*Botryococcus Braunii* Ktz. (× 400).



FIG. 350.—*Urococcus insignis* Hass. (× 400). (From nature.)

In *Urococcus* Hass. the endochrome is bright red, and the cell-walls throw off successive layers of mucilage, which form together a cylindrical or fusiform stalk, composed, in some species, of a large number of distinct annular segments.

Tetraspora Lk. is composed of cells associated together in large numbers in a single layer imbedded in a copious gelatinous envelope. It has no spontaneous motion, and is possibly allied to Merismopedia, and also appears to have affinities with the Ulvaceæ. Gay (Bull. Soc. Bot. France, 1886, Sess. Extraord., p. 41) records in T. gelatinosa (Desv.) the formation of biciliated zoospores, one being produced from the contents of each cell, and afterwards becoming encysted into a resting-spore. In *Palmodictyon* Ktz. the gelatinous envelope is filiform and branched, and cell-division takes place chiefly in two directions only.

The position of the following genera is very uncertain. Very little is known of their mode of reproduction, and they lack the copious gelatinous envelope which is characteristic of the family generally. They are mostly but feebly endowed with spontaneous movements, and may probably be a resting condition of algæ or protophytes classed under entirely different groups.

FIG. 351.—*Raphidium falcatum* Ktz. (× 800). (From nature.)

FIG. 352.—*Scenedesmus obtusus* Mey. (× 400). (From nature.)

Raphidium Ktz. includes several species very common in fresh water, and consisting of very narrow fusiform acuminate cells, usually curved, solitary or joined together in bundles, the cells being in the latter case united by their middle. Cell-division takes place in one direction only.

Under the class Palmellaceæ are usually placed also the genera *Scenedesmus* Mey. and *Polyedrium* Näg., but their rank as independent organisms is exceedingly doubtful. Their probable position has already been discussed under the heads of the Sorastreæ and the Pediastreæ respectively (see pp. 303 and 299).

Reinsch unites Polyedrium with three other genera to make up a separate family, POLYEDRIACEÆ, belonging to Palmellaceæ.

Richter connects Glœocystis with the Chroococcaceæ, and hence genetically with higher forms of algæ. Cienkowski regards Pleurococcus, Glœocystis, and probably other genera of Protococcaceæ, as resting conditions of Chlamydomonas, or of similar organisms classed among the Cœnobieæ which multiply by conjugation. Under suitable conditions he states that they can all be made to produce biciliated zoospores with two contracting vacuoles and a nucleus. The part taken by some Protococcaceæ in the development of lichens has already been discussed on p. 318.

LITERATURE.

Cohn—(Protococcus) Nov. Act. Akad. Cæs. Leop.-Carol., xxii., 1850, p. 605 (see Ray Soc., Bot. and Phys. Mem., 1853, p. 515).
Cienkowski—Bot. Zeit., 1865, p. 21.
Rostafinski—(Hæmatococcus) Mém. Soc. Sc. Nat. Cherbourg, 1875, p. 142.
Lagerheim—Oefv. Svensk. Vetensk. Akad. Förh., Stockholm, 1882, p. 47; and 1883, p. 37 (Bot. Centralbl., xii., 1882, p. 33).
Richter—Hedwigia, 1880, pp. 154, 169, 191; 1884, p. 65; and 1886, p. 249.
Dangeard—(Chlamydococcus) Ann. Sc. Nat., vii., 1888, p. 105.
Reinsch—(Polyedrium) Notarisia, 1888, p. 493.

Class XXVI.—Diatomaceæ.

The family of Diatoms—called by the older writers Bacillariaceæ—includes a very large number of genera and species, all microscopic, some of them extremely abundant in running, stagnant (but not putrid), and salt water. The individuals are strictly unicellular, and are either free-swimming and isolated, or attached to one another in a linear series or in zigzag chains, adhering to one another by means of small annular cushions, or fixed to some solid object by a simple or compound gelatinous stalk. They are, with very few exceptions, characterised by the presence in the cell-wall of a deposit of silica, by which it becomes converted into a hard but thin and perfectly transparent shell; and this is always invested in a thin gelatinous envelope. Some species are closely adherent to submerged plants by the whole of one side; in other cases whole colonies are enclosed in a common gelatinous envelope, which assumes the form of a simple or compound tube, flattened plate, or globular mass. This is especially the case with the marine species.

Each individual or *frustule* consists of two more or less symmetrical

halves known as *valves*; the silicified cell-wall of the older of these halves is slightly the larger of the two, fitting on to the younger one like the lid of a cardboard box. The cell-wall is composed of an organic matrix closely allied in composition to cellulose, impregnated with silica or a compound of silica; either of these two ingredients can be removed and the other left behind, the former by calcination, the latter by the action of hydrofluoric acid. In those species which are fixed by a gelatinous stalk, this stalk is also composed of a substance allied to cellulose. The overlapping edge of one of the two valves over the other is called the *girdle* or *hoop*; this girdle may be simple, or there may be several. In many species—and probably in all, if examined with a sufficiently high power—each valve is marked with a number of rows of very fine perforations, which, except under the very highest microscopic powers, appear as if confluent into striæ or furrows. There may be two or three sets of these apparent striæ, but they do not, as a rule, reach to the centre of the valve. So constant is the arrangement and the fineness of these striations in some of the more abundant species, that they furnish an admirable test for the definition and angular aperture of microscopic lenses. Some species of Navicula (Bory) and Pleurosigma (Sm.) are especially used for this purpose. Some marine genera in particular (Triceratium, Ehrb., Coscinodiscus, Ehrb., &c.) are characterised by the beautiful honeycomb-like *areolation* of the cell-wall, due to the presence in it of actual chambers, which may or may not be covered by a thin membrane. The membrane at the bottom of these chambers is also most minutely perforated, constituting what is known as the *secondary markings*. In describing diatoms, the aspect in which the girdle is turned towards the observer is spoken of as the *front*, *girdle*,



FIG. 353.—*Pinnularia viridis* Sm. *A*, valve-view; *B*, girdle-view (diagrammatic). *r*, furrows; *m*, raphe; *g*, central node; *k*, terminal nodules; *a*, outer and older valve; *i*, inner valve; *n*, secondary lines (× 800). (After Pfitzer.)

or *zonal view*; the aspect in which the surface of the valve is turned towards the observer is the *side* or *valve view*. In many diatoms the central space on the valve view not occupied by transverse striæ shows at its middle and at each end a strongly refractive thickening known as a node or *nodule*; and these nodules are connected with one another by a longitudinal line or rib—the *raphe* or suture. The primary classification of the genera of diatoms usually adopted depends on the presence or absence of this raphe.



FIG. 354 A.—*Anomœneis spherophora*. *a*, *c*, girdle-view; *b*, valve-view. The endochrome-plates are shaded (× 900). (After Pfitzer.)

Each diatom-cell contains a nucleus and a nucleole. The chlorophyll occurs in the form of plates or bands arranged with more or less symmetry, and there are usually also drops of oil, especially when conjugation is about to take place. A very few diatoms are green; but in the great majority of cases the colour of the chlorophyll is obscured by a characteristic brown pigment known as *diatomin*, readily soluble in alcohol, forming a brownish-yellow solution which is only slightly or not at all fluorescent. With concentrated sulphuric acid it assumes a beautiful blue-green colour. Petit (Brebissonia, 1879–80, p. 81) has very carefully investigated the chemical and physical properties of the colouring matter of diatoms. He regards diatomin as a compound of chlorophyll and phycoxanthin, and as having a great analogy with the chlorophyll of the higher plants, the two spectra being very similar.



FIG. 354 B.—*Gomphonema constrictum* Ehrb. *s*, valves, side view, showing nucleus; g_I, g_{II}, girdle-views; *q*, transverse section through middle of cell, showing silicified cell-wall, one half overlapping the other; *k*, nucleus; *p*, dense protoplasm; g_I, g_{II}, girdle surfaces (magnified). (After Pfitzer.)

Many of the solitary species of diatom, such as those belonging to

the genus Navicula, possess the power of propelling themselves through the water with considerable rapidity backwards and forwards in the direction of their longer axis, often with a jerking motion, or of creeping along the bottom on some submerged substance. The cause of this motion is a subject on which a large amount of attention has been bestowed. Nägeli attributed it to osmotic currents passing through the cell-wall. Ehrenberg believed that he had actually seen, in some cases, the extrusion through the raphe of vibratile cilia, in other cases of a 'foot' or pseudopode; but his observations have not been confirmed by others. The explanation of the motion now generally accepted is that of Schultze—that it is due to the contractility of the protoplasm which is exuded outside the cell-wall. Mereschkowsky (Bot. Zeit., 1880, p. 529) states the arguments in support of the various views with regard to the causes of the motion, and sums up in favour of the theory that it is the result of osmotic currents within the siliceous cell-wall. Hallier, again (Unters. über Diatomeen, 1880), considers it due to a contractile layer of protoplasm, and asserts that at an early stage diatoms have no true cell-wall of cellulose. Onderdonk (Microscope, 1885, p. 205) also attributes it to 'external cyclosis.'



FIG. 355.—Stages in the formation of the auxospore of *Frustulia saxonica* Ag. *s*, valves; *m*, gelatinous envelope; *c*, endochrome-plates; *a*, auxospore (× 1,200). (After Pfitzer.)

Diatoms have three modes of multiplication:—by simple division, by auxospores, and by a kind of conjugation which is regarded by some as sexual; but the three modes pass gradually one into another. Simple division always commences with the bipartition of the nucleus. When it is about to commence the two valves separate from one another, the contents divide into two daughter-cells, and new siliceous valves are formed inside the old ones, and therefore necessarily smaller than they. The valves of the new individual are formed necessarily one after the other, the one formed later being smaller. The individuals produced in

this way constantly diminish in size, until the original size is restored by the formation of an *auxospore*, resulting from the contents leaving the siliceous valves, which fall away from one another, and increasing in size, either by simple growth or by the coalescence of two auxospores produced in the same mother-cell. In other cases two distinct auxospores appear to be produced from the contents of a single mother-cell. The auxospore finally becomes invested in a new siliceous cell-wall. In those cases in which the process has been most carefully followed out, the auxospore does not appear to owe its origin to any process of true sexual union.

In some genera what is regarded by some as a true process of conjugation has been observed, a *zygosperm* being produced as the result of the coalescence of the protoplasmic contents of two different individuals. The conjugating diatoms are here placed side by side enclosed in a common gelatinous sheath; the contents of each escape by the falling apart of the two valves, and unite into a single zygosperm. In other cases two zygosperms result from the conjugation of a pair of cells. The protoplasm of each cell, as it escapes from its siliceous wall, puts out two protuberances; these meet in pairs, and the whole contents of the pair of mother-cells finally pass into the two zygosperms, which complete their development in precisely the same way as the auxospores. Buffham states (Journ. Quek. Micr. Club, 1885, p. 131) that in the conjugation of Rhabdomena (Ktz.) the 'male' frustule is always smaller than the 'female' frustule, and that the union is effected by the 'male' frustules attaching themselves in numbers to any part of the girdle of the 'female' frustule. De Bary and Pfitzer do not regard the fusion of the cell-contents of diatoms as in any sense a true process of sexual conjugation. De Bary (Bot. Zeit., 1858, Supplement, p. 61) thus summarises the four modes in which diatoms are reproduced by means of auxospores or zygosperms: —(1) Two products of conjugation are formed by the union of the contents of two distinct individuals; (2) a similar process results in the formation of a single product of the same nature; (3) a single act of conjugation (production of auxospore) takes place between two portions of the contents of the same individual; (4) two such acts of conjugation take place simultaneously between different portions of the contents of the same individual. In all cases the formation of a new individual is completed by the simple division of the product of union



FIG. 356.—*Gomphonema constrictum* Ehrb. attached by gelatinous stalks to a fresh-water alga (greatly magnified).

(auxospore or zygosperm) into two symmetrical halves with or without the intervention of a period of rest.

Still another mode of reproduction is described by Count Castracane and by some other observers, in Mastogloia (Thw.) and a few other genera, in the production of endogenous *spores* within the frustules.

It will be seen that, notwithstanding the great abundance of diatoms, some important points in their life-history still remain unsettled. On the minuter details of the modes of reproduction, the spontaneous motion of diatoms and its causes, the structure of the siliceous cell-wall, and the chemical and physical properties of diatomin, the reader is referred to the very extensive literature of the subject; only the most important memoirs are referred to below. The number of described species certainly exceeds 10,000; but this has been unduly increased by want of attention to the necessary variations in size in the same species. Not unfrequently diatoms form a gelatinous yellow scum on the surface of the water, or completely encrust submerged algæ and other water-plants; they abound on the surface of wet walls and rocks, and are not unfrequently present in the air. Some species are cosmopolitan; the marine forms are especially remarkable for their size and beauty. Various deposits found on the surface of the globe, often of very considerable thickness, known as tripoli, 'Kieselguhr,' &c., consist almost entirely of the fossilised siliceous shells of diatoms, and they enter largely into the composition of a variety of earths used for manufacturing purposes. In some countries, such as China, Japan, Siberia, Lapland, &c., they form, cemented together by salts of lime, the edible earths which are mixed with meal to make a kind of flour. They occur also in large quantities in guano.

As has already been stated, the position of the Diatomaceæ in the natural system is a point on which there has been much controversy. Those who regard the mode of reproduction already described as a true process of conjugation place them in the class of Conjugatæ, near to the Desmidiaceæ, with which family they present many points of resemblance in external form, phenomena of spontaneous movement, &c.; and it is possible that the diatoms may be derived from the desmids by retrogressive metamorphosis. But we are, on the whole, disposed to the conclusion that they have a totally different origin; their very wide distribution in time and space, the sharp differentiation of the family, and the enormous number of species, favouring the view that they represent a comparatively small ascent from an archaic type which has never attained any higher degree of development.

Illustrative genera:—Eunotia (Ehrb.), Diatoma (DC.), Melosira (Ag.), Gomphonema (Ag.), Navicula (Bory), Rhabdonema (Ktz.),



FIG. 357—Diatomaceæ: *A*, *Eunotia monodon* Ehrb.; *B*, *Triceratium Favus* Ehrb.; *C*, *Surirella splendida* Ktz.; *D*, *Synedra Arcus* Ktz.; *E*, *Navicula rhomboides* Ehrb.; *F*, *Pleurosigma lacustre* Sm.; *G*, *Cocconema lanceolatum* Ehrb.; *H*, *Meridion constrictum* Rlfs.; *I*, *Achnanthes brevipes* Ag.; *K*, *Diatoma elongatum* Ag. (variously magnified). (After W. Smith.)

Nitzschia (Hass.), Pleurosigma (W. Sm.), Achnanthes (Bory), Meridion (Leibl.), Biddulphia (Gray), Amphora (Ehrb.), Campylodiscus (Ehrb.), Cymbella (Ag.), Epithemia (Bréb.), Pinnularia (Ehrb)., Stauroneis (Ehrb.), Surirella (Turp.), Synedra (Ehrb.), Mastogloia (Thw.), Amphipleura (Ktz.), Fragillaria (Lyng.), Tabellaria (Ehrb.), Aulacodiscus (Ehrb.), Coscinodiscus (Ehrb.), Licmophora (Ag.).

LITERATURE.

Agardh—Conspectus Diatomacearum, 1830.
Ehrenberg—Die Infusionsthierchen, 1838.
Kützing—Die kieselschaligen Bacillarien, 1844.
Rabenhorst—Die Süsswasser-Diatomaceen Deutschlands, 1853.
Rylands—(Marking of diatoms) Quart. Journ. Micr. Sc., 1860, p. 25.
Grünow—Oesterreichische Diatomaceen, 1862.
Heiberg—Kritisk Oversigt Danske Diatomaceer, 1863.
Cleve—Svenska och Norska Diatomaceer, 1868.
W. Smith—Synopsis of British Diatomaceæ, 1872.
Hamilton L. Smith—Conspectus of the Diatomaceæ, Lens, 1873, p. 63.
Habirshaw—Catalogue of the Diatomaceæ, 1877.
Mereschkowsky—(Movements of diatoms) Bot. Zeit., 1880, p. 520.
Hallier—Untersuchungen über Diatomeen, 1880.
Brun—Diatomacées des Alpes, 1880.
Van Heurck—Synopsis des Diatomacées de Belgique, 1881.
Pfitzer—Die Bacillariaceen, in Schenk's Handbuch der Botanik, ii., 1882.
Castracane—Diatomaceæ of the *Challenger* Expedition, 1886.

Also numerous papers by Wallich, Thwaites, Carter, Lüders, Millardet and Kraus, Schultze, Schmitz, Askenasy, Borzcow, Cox, Schaarschmidt, O. Müller, Flögel, Schütt, Nelson and Karop, Deby, Brun, Cleve, Grünow, Janisch, F. W. Lewis, Greville, O'Meara, Petit, Gregory, Donkin, Ralfs, Walker-Arnott, Castracane, Brébisson, Kitton, Leuduger Fortmorel, Pfitzer, Rattray, Van Heurck, Grove and Sturt, and others.

For the preparation of diatoms as microscopic objects see H. L. Smith, Lens, 1873, p. 209; Kitton, Science Gossip, 1877. For detailed bibliography see Deby and Kitton, Bibliography of the Microscope, pt. 3, the Diatomaceæ, 1882.

Class XXVII.—Cyanophyceæ.

In the Cyanophyceæ or Phycochromaceæ are included all those unicellular chlorophyllous organisms in which the cell-sap is uniformly pervaded by a blue-green pigment which masks the pure green of the chlorophyll. This pigment can be separated, giving, in its aqueous solution, a beautiful blue colour in transmitted, a blood-red in reflected

light, and is known as *phycocyanin*. This colour is, in a few cases, replaced by a red, and, in some Scytonemaceæ, by a brown endochrome, the *scytonemin* of Nägeli.

Although the Cyanophyceæ are described as unicellular, this term must be used with some limitation in the sense employed with respect to the Protococcoideæ, in comparison with which some of the blue-green algæ display greater differentiation in their vegetative structure. In several families, notably the Oscillariaceæ and Rivulariaceæ, the protoplasm of the very long filament is broken up transversely into a large number of imperfect cells known as *pseudocysts*, which are, however, never invested with a true cellulose cell-wall. In the Nostocaccæ, the breaking up of the filament into cells is much more complete ; but even here it is doubtful whether the very thin membrane in which they are invested is composed of true cellulose. It displays the strongest tendency to deliquesce into a hyaline jelly, and, not unfrequently, as in the Scytonemaceæ and some Nostocaceæ, this gelatinous envelope becomes strongly coloured. Chromatophores with enclosed pyrenoids and nuclei are stated to have been detected in some species ; while other authorities assert that neither starch nor a true nucleus is found in the Cyanophyceæ.

Multiplication by ciliated zoospores is, with a few doubtful exceptions, unknown in the Cyanophyceæ. The ordinary mode of propagation in the lower families is by simple cell-division. Quiescent *resting-spores* or *cysts*, with thicker cell-walls, are also produced in several families. In the filiform orders, portions of the filament known as *hormogones*, consisting of a varying number of pseudocysts, become detached ; these hormogones are endowed for a time with a power of spontaneous motion, after which they come to rest, and develop into new filaments. In the Oscillariaceæ this power of motion extends to the entire individual. In the Rivulariaceæ, Scytonemaceæ, and Nostocaceæ, a further differentiation is exhibited, of special cells within the filament known as *heterocysts* ; but their function is unknown.

The life-history of many of the Cyanophyceæ is still but imperfectly known, and their classification into families is very obscure. Some of those who have most closely investigated their structure and development are of opinion that a large number of the apparently distinct forms are but stages in the development of the same pleomorphic organism, often really belonging to a higher type. This will be further discussed under the separate families. They may be conveniently divided into two groups : the *Chroococcaceæ*, which are strictly unicellular, and the *Nostochineæ*, embracing those families in which the aggregation of cells or pseudocysts assumes more or less of a filiform character.

Sub-class 1.—Nostochineæ.

It is convenient to group together under this head those families of Cyanophyceæ, the *Nostocaceæ*, *Rivulariaceæ*, *Scytonemaceæ*, and *Oscillariaceæ*, which are characterised by the individual consisting of a cellular or pseudocellular filament, reproduced by motile hormogones, and, in some of the families, forming heterocysts.

Borzì, who has closely examined the structure and life-history of the Nostochineæ, considers that they display only a rudimentary cell-structure. The cell-wall is extremely thin, is inseparable from the contained protoplasm, and not sharply differentiated from it in its chemical reactions. The cells do not contain true starch, its place being taken, in spores and in the constituent elements of the hormogones, by a substance to which he gives the name *cyanophycin*—of a granular nature, apparently formed from the substance of the cell-wall, possibly a carbohydrate, and identical with the gelatinous substance of which the sheath which encloses the filaments is composed. The only indication of a nucleus is a local fragmentation and concentration of proteinaceous matter. Borzì states that in all the families of Nostochineæ the cells of a filament are in communication with one another by pores in their transverse walls, through which pass strings of protoplasm, or of a substance allied to cellulose. When heterocysts are about to be formed, this intercellular communication is suspended, and the pores are closed by a thickening of the cyanophycin or other cell-contents. In the Nostocaceæ, Scytonemaceæ, and Rivulariaceæ, the filaments are enclosed in a gelatinous sheath, which is much thinner in the Oscillariaceæ, and altogether wanting in Borzia (Cohn). This mucilaginous sheath may be continuous or septated, and may be composed of a single layer or of several ; in the latter case it not unfrequently assumes the form of a funnel at the end of the filament. Gomont differs to a certain extent from these conclusions, previous observers having, according to him, confounded the envelope proper of the cell with the mucilaginous sheath of the trichome. By the action of chromic acid he was able to separate the one from the other, and finds the very thin perfectly transparent envelope proper of the cell to possess properties intermediate between those of fungus-cellulose and of vegetable cutin. Bornet and Flahault call the string of cells or pseudocysts the *trichome*, the trichome with its envelope the *filament*.

When in a purely vegetative condition, all the cells of a filament are alike in form and size, and multiply by repeated bipartition. Scott and Wille have been able to demonstrate the presence of a nucleus in

several species of Oscillaria (Bosc.) and Tolypothrix (Ktz.). The *hormogones* consist of a number of cells, often considerable, but usually variable in the same species. Only in a few species is the number constant, and then usually two, three, or four. Their contents are generally of a yellowish-green colour. According to Borzì, the hormogones of the Nostochineæ are of two kinds, straight and spiral. The latter kind occur only in the Oscillariaceæ, and are confined to the terrestrial species of Oscillaria, Microcoleus (Desm.), and Spirulina (Lk.). They are invested in a thin gelatinous sheath, and their movement is of a spiral nature, lasting much longer than that of the straight hormogones. These latter occur in all the families ; they are not invested in a gelatinous sheath ; their motion is rectilinear and of short duration. The cells have excessively thin cell-walls, and contain abundance of cyanophycin, slightly coloured by phycocyanin. The cells of the hormogones are in communication with one another in the same way as the ordinary vegetative cells.

The *spores* of Nostoc and other Nostochineæ are, according to Borzì, formed as follows. The cells which become transformed into spores cease dividing transversely and increase somewhat in size. The gelification of the outer layer of the cell-walls ceases at the same time and a new cell is formed inside the old one by a process of rejuvenescence, the wall of the mother-cell ultimately disappearing altogether. The membrane of the spores thus constituted is formed out of their protoplasmic contents, and is homogeneous, without differentiation into exospore and endospore. The spore is, therefore, not homological with those of the higher Cryptogams, but is regarded by Borzì as partaking more of the nature of a cyst. Gomont, on the other hand, finds that the spores of the Nostochineæ, which are always the result of the encysting of ordinary vegetative cells, possess a distinct exospore and endospore, the former being again composed of two distinct layers, the outer of which is frequently warty or otherwise marked.

Literature.

Thuret—Mém. Soc. Sc. Nat. Cherbourg, 1857, p. 29 ; and Classification des Nostochinées, 1875.

De Bary Beitr. zur Kenntniss d. Nostocaceen, Flora, 1863, pp. 553 and 577.

Bornet and Thuret—Notes Algologiques, 1876–1880 (especially Fasc. ii.).

Borzì—Nuov. Giorn. Bot. Ital., 1882, pp. 272 and 384 ; and Malpighia, 1886.

Bornet and Flahault—Ann. Sc. Nat., iii.–vii., 1886–1888 ; and Mém. Soc. Sc. Nat. Cherbourg, 1887, p. 195.

Gomont—Morot's Journ. de Bot., 1888, p. 43 ; and Bull. Soc. Bot. France, 1888, p. 204.

Order 1.—Nostocaceæ.

The Nostocaceæ are distinguished from all the other families of filamentous Cyanophyceæ by the less close connection with one another of the pseudocysts of which the filament is composed, giving it always more or less of a moniliform or necklace-like appearance. In most of the genera these pseudocysts are spherical or elliptical; but in Nodularia (Mert.) they are disc-shaped, and more closely connected with one another than in the other genera. The extent to which the filaments are enveloped in mucilage varies greatly. In some genera (Nodularia) each filament is enclosed in a distinct hyaline sheath. In others (Anabæna, Bory, Aphanizomenon, Morr., Sphærozyga, Ag., Cylindrospermum, Rlfs.) this sheath is obscure or wanting; in Aulosira (Kirch.) the filaments are either naked, or enclosed in a dry membranous sheath. In most species of Nostoc (Vauch.) a single filament or a number of filaments are enclosed in a hyaline jelly, often of considerable size and definite outline, formed from the more or less complete coalescence of the separate sheaths, which can still sometimes be indistinctly made out. Such a jelly-like mass, which sometimes floats freely in the water of bog-pools, but is more often found on damp soil, is sometimes called a 'thallus' or 'frond.' It may vary in size from 0·2 mm. to that of a small plum; is often of a green, violet, or blue colour, and, in the larger terrestrial species, the outer layers of the integument are more or less hardened, forming what is known as a 'periderm.'



Fig. 358.—*Nostoc commune* L. (natural size). (After Cooke.)

The Nostocaceæ display no differentiation of the two extremities of filament. In nearly all the genera some of the cells are here and there replaced by *heterocysts*—cells incapable of further division, of a slightly larger diameter, and with a somewhat thicker cell-wall, which is often yellow, the green protoplasmic endochrome being replaced by a watery colourless cell-sap. These heterocysts, the function of which is unknown, may be terminal, basal, or intercalary. Others of the cells, always in this case intercalary, are, in most species, replaced by *resting-spores*, also distinguished from the ordinary cells by their larger size and thicker cell-wall, but containing a green endochrome. They are often formed more or less in connection with the heterocysts. In Cylindrospermum the terminal cells of the filaments become heterocysts, and the spores, some-

times associated in chains, are formed immediately beneath them. The cell-wall of the resting-spores is often coloured yellow or brown, and is sometimes warty. After a period of rest they germinate by the bursting of the cell-wall and protrusion of the protoplasm. The *hormogones* are strings of from four to eight or twelve ordinary cells, situated between two heterocysts, which detach themselves from the rest of the filament, escape from their mucilaginous envelope, move about with a creeping motion, then come to rest, and develop into new individuals.

Bornet describes the motion of the hormogones of Nostoc as a creeping movement along a solid substratum at a rate of 1 μ (·001 mm.) per second. After some hours they come to rest, large refringent globules which had previously been formed in the cells disappear, and they assume the appearance of ordinary filaments. Sometimes they



FIG. 350.—*N. hyalinum* Benn. *A*, 'frond' (× 200); *B*, portion of trichome with heterocyst, *c* (× 600). (From nature.)

FIG. 360.—*Anabæna flos-aquæ* Fr. (× 400). (After Cooke.)

invest themselves with a mucilaginous sheath, and are transformed directly into spores, but usually the filament lengthens, displaying at the same time more or less sinuosity. The heterocysts do not appear at regular intervals. A new filament is thus formed altogether resembling those which spring directly from the germination of the spores. Once formed, it is subject to intermittent periods of growth, a second generation of spores, with thinner cell-walls, being sometimes formed after the first. In other cases the filament springing directly from a hormogone assumes a zigzag form, in consequence of some of the cells dividing in the transverse, others in the longitudinal direction. In N. muscorum (Ag.) all the cells except the heterocysts are sometimes transformed into spores. The spores will occasionally germinate while still within the 'frond.' The filaments of Nostoc are ultimately serpentine, and often

intricately interwoven, owing to the heterocysts maintaining their position when once formed, while the portions of the filament between them continue to lengthen by cell-division.

Most of the Nostocaceæ inhabit fresh, usually stagnant, water, bog-pools, &c.; a very few grow in salt water. Several species of *Anabæna* Bory, *Aphanizomenon* Morr., and some other genera, frequently collect as a bright bluish-green scum on the surface of bog-pools; Aphanizomenon flos-aquæ (Rlfs.) is one of the organisms which contribute to the phenomenon know as the 'breaking of the meres.' The decay of species of Anabæna has a large share in causing the foul odour and injurious properties of stagnant water. Of the numerous species of *Nostoc* Vauch., a few only swim freely in water; these are usually minute with a colourless 'thallus' more or less nearly spherical and distinctly lamellated, with a spontaneous power of active motion in the water. The larger number are terrestrial, growing on wet rocks, among moss on damp soil, or even on the comparatively dry ground, forming conspicuous jelly-like masses, often as much as from one to two inches in diameter, usually of a green, yellow, brown, or sometimes red colour, the 'frond' enclosing a large number of bright blue-green filaments (fig. 358). It is not uncommon for other chlorophyllous protophytes or algæ, the mycele of fungi, and the protoneme of mosses, to become enclosed, and even to continue growing, within their gelatinous envelope. Species of Nostoc and Anabæna are found in hot springs, and several species of the same genera are frequently to be met with within the cells of living aquatic or land plants, where they carry on an epiphytic existence (see pp. 28, 165, 171). They occur with especial frequency in Lemna, causing yellow spots in the leaves.



FIG. 361.—*Aphanizomenon flos-aquæ* Morr., with spore (× 400). (After Cooke.)

FIG. 362.—*Cylindrospermum macrospermum* Ktz., with heterocysts and spores (× 400). (After Cooke.)

Isocystis Bzi. is probably the simplest form of the Nostocaceæ. It consists of delicate moniliform filaments, which do not form colonies, but

float freely on the surface of the water, solitary or in interwoven masses, forming but little mucilage, and destitute of heterocysts.

Hansgirg (Bot. Centralbl., xxii. and xxiii., 1885) regards the genera Nostoc, Anabæna, Cylindrospermum, and Sphærozyga as stages of development—analogous to certain zooglœa-conditions of the Schizomycetes—of various species belonging to the Oscillariaceæ, Rivulariaceæ, and Scytonemaceæ. The relationship of Nostoc with Drilosiphon will be alluded to under this last family (see p. 440). The Nostocaceæ appear also to approach the Schizomycetes through Leuconostoc (Van Tiegh.).

Among the Nostocaceæ are included, in addition to the genera already mentioned, Nodularia (Mert)., Cylindrospermum (Ktz.), Sphærozyga (Ag.), and Aulosira (Kirch.).

LITERATURE.

Meneghini – Monogr. Nostoc. Ital., 1843.
Thuret Mém. Soc. Sc. Nat. Cherbourg, 1857, p. 23; Ann. Sc. Nat., 1875, p. 372.
Janczewski—Ann. Sc. Nat., 1874, p. 123.
Bornet and Thuret—Notes Algol., fasc. i., pp. 2–3; fasc. ii., pp. 78–132.
Borzì—Flora, 1878, p. 465; and Nuov. Giorn. Bot. Ital., 1878, p. 236.
Fischer—Beitr. zur Kenntniss der Nostocaceen, 1853.

ORDER 2.—RIVULARIACEÆ.

In the Rivulariaceæ – including also the Calotrichaceæ—the cells or pseudocysts of which each filament is composed are not connected together in a necklace-like form, but constitute a continuous thread divided by transverse septa which are exceedingly thin and often scarcely perceptible. Many of the species are extremely minute, the individual filaments being quite microscopic, and grow attached in tufts to a solid substratum, some water-plant or floating body, from which they radiate in a star-shaped manner, forming small green discs or cushions, often imbedded in copious mucilage. Each filament displays a distinct differentiation of the two apices, the distal extremity being elongated and attenuated into a hyaline hair, while at the base a portion is marked off into a more or less globular colourless *basilar cell*, the rudimentary cell-division being exhibited by the portion intermediate between the basilar cell and the terminal hyaline hair. The terminal hair is, according to Gomont, in perfect continuity with the rest of the true membrane of the filament, and is distinguished only by having fewer transverse septa, and by the entire absence of granular protoplasm. The outer layers of the walls of the filaments have a very strong tendency to become transformed into mucilage, from which is formed not only the copious jelly in which

the colony is usually invested, but in most species also a separate delicate membranous sheath to each individual, often of a yellowish colour, which ultimately becomes open at the apex. In Microchæte diplosiphon (Gomont, Bull. Soc. Bot. France, 1885, p. 209) this sheath is double. The common gelatinous envelope of the colony sometimes becomes at length calcareous. In other species, which float on the surface of the water, the development of mucilage is much feebler.

The filaments of the Rivulariaceæ branch in a peculiar way. A small portion of the green tract between the basilar cell and the terminal bristle becomes rounded off, loses its endochrome, and forms a *heterocyst* resembling the basilar cell. This heterocyst now assumes the part of a basilar cell ; the terminal portion of the filament beyond it detaches itself, and applies itself laterally to the heterocyst,



FIG. 363. —*Rivularia polyotis* Hauck. *a*, invested in its gelatinous envelope (natural size). *b*, vertical section (× 12). *c*, filament (× 150). (After Hauck.)

forming a 'pseudo-ramulus' or false branch ; the original filament then develops a new apical portion in a direct line above the heterocyst. Heterocysts have not yet been observed in all the genera ; they sometimes occur interstitially in the filament without giving rise to a pseudo-ramulus ; their function is obscure. The terminal hyaline bristle is of

only temporary duration ; when it disappears it leaves the membranous sheath open at the extremity. This is especially well seen in Calothrix (Ag.).

The ordinary mode of multiplication of the Rivulariaceæ is by means of *hormogones*, fragments of the green portion which become detached from the rest of the filament, escape from the gelatinous envelope, move about with a creeping motion, eventually come to rest, invest themselves with a gelatinous sheath, and develop into a new filament in which the differentiation of the basal and apical extremities is soon manifested. The formation of hormogones is confined to the lower and central portions of the filament, and commences only after the disappearance of the terminal hyaline hair. They vary greatly in length, being composed of from two to fifty pseudocysts. When fully formed, they glide slowly out of the sheath, several often attached to one another. At the period of detachment of the hormogones the whole filament displays a slow movement ; otherwise it is quiescent, the power of motion which in the Oscillariaceæ belongs to the entire filament being in the Rivulariaceæ restricted to the hormogones. Beck (Verhandl. zool.-bot. Gesell. Wien, 1886, p. 47) describes a peculiar mode of formation of the hormogones in Glœotrichia natans

FIG. 364.—*Calothrix crustacea* Thur. (× 160). (After Bornet.)

(Thur.) while still within the sheath, the cell-contents passing from a heterocyst into the basal cell of a hormogone.

Multiplication by quiescent *resting-spores* has been observed in some species of Rivulariaceæ. The lower portion of the green part of a filament immediately above the basilar cell is transformed into an elliptical thick-walled spore, which escapes from its investing membrane, and, after a period of rest, either develops directly into a new filament, or breaks up into a number of hormogones. The spores of Glœotrichia punctulata (Thur.) are rough. Under the name 'conids' Bornet and Flahault also describe special propagative cells which become detached from the lower part of the filament.

FIG. 365.—*Isactis plana* Thur. (× 160). (After Bornet.)

By far the larger number of species of Rivulariaceæ grow in fresh water, especially stagnant, also on damp soil and on wet rocks. The species of the typical genus *Rivularia* Roth are especially abundant in both running and standing water; R. fluitans (Cohn, Hedwigia, 1878, pp. 1 and 33) floats free on the water, forming a blue-green scum which enters largely into the composition of what is known as 'flos aquæ.' Some species have a red tinge. *Calothrix* Ag., though placed by Rabenhorst among the Scytonemaceæ, has all the characteristics of the Rivulariaceæ. The filaments of some species are comparatively thick, as much as 0·01 mm. in diameter, and are invested in a copious, often coloured, mucilaginous sheath. *Microchæte* Thur. is nearly allied to Calothrix, as is *Glœotrichia* Ag. to Rivularia. Several species of Rivularia and Calothrix grow in salt or brackish water, and *Isactis* Thur. is exclusively marine. In *Hormactis* Thur., which is also marine, the filaments are curved in a serpentine manner, and this character, together with the interstitial heterocysts, appears to indicate an approach to the Nostocaceæ. Other genera included in the Rivulariaceæ are Leptochæte

(Bzi.), Amphithrix (Ktz.), Dichothrix (Zanard.), Sacconema (Bzi.), Brachytrichia (Zan.), and Polythrix (Zan.), the last being marine.

Hansgirg (Bot. Centralblatt, xxii. and xxiii., 1885) considers the genera ordinarily placed under Rivulariaceæ as being higher developments of organisms belonging to the Oscillariaceæ.

LITERATURE.

De Bary—Flora, 1863, p. 577.
Bornet and Thuret—Notes Algol., fasc. i., 1876, pp. v.–viii.; and fasc. ii., 1880, pp. 157–175.
Bornet and Thuret—Etud. Phycol., 1878, pp. 1–6.
Bornet and Flahault—Ann. Sc. Nat., iii. (1886), p. 337; and iv. (1886), p. 341.

ORDER 3.—SCYTONEMACEÆ

(including STIGONEMEÆ and SIROSIPHONEÆ).

The Scytonemaceæ resemble the Rivulariaceæ in consisting of branched filaments, often comparatively stout, enclosed, either singly or in numbers, in a mucilaginous sheath; but differ from that family in displaying no differentiation of the two extremities. The filament terminates at each end in a large thin-walled apical cell, by the repeated division of which the greater part of the growth in length takes place. The filaments display no oscillation or other spontaneous motion. The mucilaginous sheath which invests one or more filaments is of considerable thickness, except over the apical cells, where it is very thin; elsewhere it is generally lamellated, the lamellæ decreasing in number towards the apex, which gives the appearance of a number of funnels inserted one in another. It is often coloured by a deep yellow or brown pigment known as *scytonemin*, and becomes dry and pulverulent with age, but in younger filaments the sheath is sometimes altogether wanting. The filaments are not septated laterally, but the contents are divided into 'gonids' or *pseudocysts*, of a spherical or elliptical form, and arranged in a single row in the thinner, often in two parallel rows in portions of the thicker, filaments. These pseudocysts are at first green, but frequently become subsequently dark brown; and the filament exterior to the pseudocysts is commonly filled by endochrome coloured brown by scytonemin; the entire plant being therefore distinguished by its brown or orange colour. In Tolypothrix (Ktz.) and Plectonema (Thur.) the filaments generally retain permanently a green colour.

The Scytonemaceæ may multiply by the individual filaments, enclosed in a common sheath, which have no genetic connection with one another, escaping separately from their sheath, and then investing themselves in a new mucilaginous envelope. But the ordinary mode of

propagation is either by *resting-spores* or by *hormogones*. In the former case, at the end of the season of growth, the disc-shaped pseudocysts towards the end of a filament assume a roundish or ovate form, the gelatinous sheath disappears, and the spores thus formed remain for a time united together in masses. These resting-spores are capable of resisting a high degree of cold and drought; they germinate after a period of rest, when the membrane bursts transversely. In the formation of hormogones the sheath also becomes absorbed, beginning from the apical cell, and the filament breaks up into a number of hormogones, each consisting of many pseudocysts. In the Stigonemeæ the hormogones are formed only in the lateral branches, which contain only a single row of pseudocysts. The hormogones move slowly in the water in a straight line; in some cases an entire filament may be converted into a motile hormogone. During germination the hormogone, enclosed in a delicate transparent mucilaginous sheath, breaks up into portions of various lengths, or it becomes a new individual without breaking up; and at this period one of the apical pseudocysts usually becomes a *heterocyst*.



FIG. 366. – *Stigonema minutum* Hass. *A*, outline of filament (× 100); *B*, portion of primary filament (× 200); *C*, ditto with branch (× 200). (From nature.)

The branching of the Scytonemaceæ takes place in two different ways, characteristic of the two sub-orders into which they are divided—the *Scytonemeæ* and the *Sirosiphoneæ*. In the Scytonemeæ, which include the genera Scytonema (Ag.), Petalonema (Berk.), Tolypothrix (Ktz.), Plectonema (Thur.), and Drilosiphon (Ktz.), *pseudo-ramuli* are formed somewhat in the same way as in Rivularia, but the branches are much stouter, and stand out at right angles to the main axis. In *Scytonema* Ag. two contiguous pseudocysts separate at indefinite spots on the filament, and each of these then acts as the terminal pseudocyst

to a lateral branch ; the branches therefore spring in pairs at a right angle from the main axis. Thick-walled heterocysts may be formed at any spot in the filament. In *Tolypothrix* Ktz. the false branches spring singly from beneath heterocysts. In the Sirosiphoneæ, including Stigonema (Ag.), Sirosiphon (Ktz.), Fischera (Schw.), Capsosira (Ktz.), Hapalosiphon (Näg.), Mastigocladus (Cohn), and Mastigocoleus (Lagerh.), the formation of a false branch is preceded by a change in



FIG. 367.—*Hapalosiphon byssoideus* Kirch. (× 200). (From nature.)

the direction of cell-division. Two or three contiguous pseudocysts in an older portion of the filament divide in a direction parallel to the axis of growth of the filament, and one of the new pseudocysts thus formed now becomes the basal pseudocyst of a lateral branch, which generally consists of a single row of pseudocysts at right angles to the axis. In *Plectonema* Thur. the branches protrude outside the mucilaginous sheath. Heterocysts are formed in all parts of the filament, but their function is unknown.

Coleodesmium Bzì. appears to be one of the simplest forms of the Scytonemaceæ. No pseudo-ramuli are formed ; the filaments increase by fission only, and a number remain united in a bundle within a common envelope. *Mazæa* (Bornet, Bull. Soc. Bot. France, 1881, p. 287) is, on the other hand, a genus in which the development is carried to its highest point. The gelatinous 'fronds' are about 25 mm. in diameter ; the heterocysts are borne on pedicels consisting of from one to three cells, and the whole appearance is that of a Rivularia, the filaments being immersed in a homogeneous jelly, and spreading from a central spot. No distinct sheath has been observed, nor any resting-spores or hormogones. In *Petalonema* Berk. the mucilaginous sheath forms a kind of broad coloured wing to the filament. *Mastigocoleus* (Lagerheim, Notarisia, 1886, p. 65) is a marine genus growing attached to the shells of molluscs, with both terminal and lateral heterocysts ; the filament sometimes ends in a hair, as in the Rivulariaceæ. *Drilosiphon Julianus* Ktz., frequent on the damp walls of greenhouses, is characterised by an outer calcareous sheath, and is a remarkably pleomorphic organism. According to Zukal (Oesterr. Bot. Zeitschr., 1883, p. 73) it forms two kinds of hormogones, and displays a kind of alternation of generations. The ordinary hormogones produce only more and more slender filaments, which gradually assume a moniliform character, and are then known as Nostoc parietinum (Rabh.). Eventually the cells of these nostoc-filaments separate, and assume the character of an Aphanocapsa, or, in other cases, become the organism known as Glœocapsa fenestralis (Ktz.) ; or very slender filaments are produced, constituting the Leptothrix parasitica and muralis (Ktz.), which forms are distinctly connected genetically with Drilosiphon. These leptothrix-filaments may again break up into vibrio- and bacillus-forms. The second kind of hormogone has a fusiform shape, and consists usually of from four to eight pseudocysts. It may remain dormant for a time, and, on germinating, reproduces the ordinary thick filaments.

Wille (Ber. Deutsch. Bot. Gesell., 1883, p. 243) and Scott (Journ. Linn. Soc., xxiv., 1887, p. 188) have determined the presence of a cell-nucleus in Tolypothrix. Wille states also that in Stigonema compactum (Kirch.) the necklace-like pseudocysts are in direct communication with one another through perforations in their cell-walls. When this species passes into the Glœocapsa-condition, the perforations disappear, in consequence of the gelification of the common sheath, and the separate cells then carry on their existence as distinct individuals. Under the name Tolypothrix amphibia, Zopf (Ber. Deutsch. Bot. Gesell., 1883, p. 319) describes an organism having both an aerial and an aquatic form, the latter being a true Tolypothrix with its filaments

enclosed in sheaths and breaking up into hormogones, from which is derived the aerial form with the nature of a Chroococcus, and dividing in three directions.

Rabenhorst and Cooke regard Stigonema, and the latter authority also Hapalosiphon, as genera of lichens ; and Bornet and Flahault state that several organisms described as species of Sirosiphon and Stigonema are really lichens in a more or less advanced stage of development. Hansgirg (Oesterr. Bot. Zeitschr., 1884, and Bot. Centralbl., xxii. & xxiii., 1885) considers the genera placed under Scytonemeæ to be the highest forms of development of various organisms hitherto mostly placed under Oscillariaceæ. In the same way, from Tolypothrix and Scytonema may arise, by further development, the corresponding forms of Hapalosiphon, Mastigocladus, Sirosiphon, Stigonema, Fischera, and other genera usually placed under Sirosiphoneæ.

With the exception of Mastigocoleus, the Scytonemaceæ are found only in fresh water, in bog-pools, or very commonly on wet rocks or trunks of trees, or among moss. They may form mats of considerable thickness, but the individual filaments, including the sheath, seldom exceed 0·04–0·05 mm. in thickness. Several Scytonemaceæ are known to enter into the composition of lichens (see fig. 279 D.)

LITERATURE.

Bornet—Notes Algol., fasc. 1, 1876, pp. iv.–v. ; fasc. 2, 1880, pp. 135–156.
Bornet and Flahault—Ann. Sc. Nat., v., 1887, p. 51.

ORDER 4.—OSCILLARIACEÆ (including CHAMÆSIPHONACEÆ).

The Oscillariaceæ or Oscillatorieæ, in which the Lyngbyeæ are also included, consist of delicate blue-green threads, occurring singly or in large floating masses in fresh running, or more abundantly in stagnant, less often in salt, water. The filaments are cylindrical and unbranched, straight, or (Oscillaria princeps, Vauch.) with the terminal portion bent at an obtuse angle with the rest of the filament ; in Spirulina (Lk.) the whole filament is coiled in a corkscrew-like manner. The filament is divided by very delicate transverse septa into disc-shaped pseudocysts ; there is no differentiation between the two extremities. The cell-wall has the property of transforming its outer layers into copious mucilage, which forms a gelatinous sheath investing either single filaments, as in Lyngbya (Ag.) and Symploca (Ktz.), or a number of filaments, as in Inactis (Ktz.) and Microcoleus (Desm.). In most species of Oscillaria (Bosc.) and Spirulina a distinct sheath is either wanting or it is extremely thin and delicate, but the filaments are often imbedded in structureless jelly.

The blue-green colour of phycocyanin is sometimes replaced by a red or violet pigment. Scott finds a cell-nucleus in several species of Oscillaria.

The only certainly known mode of multiplication of the Oscillariaceæ is by a filament escaping from its mucilaginous sheath, and breaking up into *hormogones*, each composed of a small number of pseudocysts, which round themselves off at both ends and develop into new filaments.

Fig. 368.—*Oscillaria tenuis* Ag. (× 400). (After Cooke.)

The family derives its name from the oscillating or wavy motion with which the filaments are endowed. This consists in a creeping movement in the direction of the length of the filament, now backwards and now forwards, accompanied by a curvature of the filament and rotation round its own axis; but, according to Borzì, this power of motion is limited to the reproductive period. The filaments of the Oscillariaceæ have a remarkable power of resistance to both cold and desiccation, to which they are adapted by the encysting of the filament and hardening of the mucilaginous sheath.

The movements of the Oscillariaceæ are greatly influenced by temperature and light, being much more active in warmth and sunshine than in cold and shade, but their cause is involved in considerable obscurity. Cohn (Arch. mikr. Anat., 1867, p. 48) observed that the oscillating movements take place only when the filament is in contact with a solid substratum. Siebold (Zeitschr. wiss. Zool., 1849, p. 284) states that if the water in which they grow is coloured by indigo, the particles collect round the filaments of Oscillaria up to their apex, whether they are in motion or not. Sometimes creeping spiral lines of pigment begin to be formed at both ends of the filament and meet in the middle, where the particles become heaped up into little balls; or sometimes this begins in the middle and advances to both ends. The mode in which the particles of indigo adhere to the filament and to one another appears to indicate the excretion of mucilaginous protoplasm.

Engelmann (Bot. Zeit., 1879, p. 49) claims to have detected this external secretion in the case of Oscillaria dubia (Ktz.). Zukal compares the motion of Spirulina to that of a growing tendril, and asserts that it is intimately connected with the growth of the filament. It consists of a slow torsion of the entire helix round its own axis, and is the result of the more rapid growth in length of the filament than of the ideal axis of growth. If the motion is suddenly interrupted, the filaments become for a moment quiescent, and then retreat towards the central point of the movement, forming a dark green lump. Hansgirg, on the other hand (Sitzber. Böhm. Gesell. Wiss., see Bot. Centralbl., xii., 1882, p. 361), considers the twisting and nodding movements to be due, not to the growth of the filament, but to osmotic changes in the cell-contents ; the separate cells exhibiting motion when the envelope itself is at rest. He regards the movements as of the same nature as those of the sarcode in the pseudopodes of Rhizopods and other Protozoa. The same observer states further (Bot. Zeit., 1883, p. 831) that in the protoplasm which had escaped from the broken end of a filament of O. princeps, he has observed



FIG. 369.—*Oscillaria princeps* Vauch. (× 200). (From nature.)

a number of amœboid cells from 9 to 12 μ in diameter, nearly spherical in form, and putting out colourless pseudopodes about twice the length of the central body, and to these he attributes the motile properties of the protoplasm. He believes the cause of the oscillating motion to be that the internal protoplasm takes up water more rapidly, and consequently swells to a greater extent, than the enveloping sheath, causing the filament to move slowly backwards and forwards within the sheath. In those species in which each filament is not invested in a separate sheath, variations in the turgidity are also brought about by variations in the endosmotic and exosmotic currents. Finally Schnetzler (Arch. Sc. Phys. et Nat., 1885, p. 164) describes the movements in O. ærugineo-cœrulea (Ktz.) as of six different kinds, viz. : (1) rotation of the filament or of its segments round its axis ; (2) creeping or gliding over a solid substratum ; (3) a swimming change of position in the water ; (4) rotation or flexion of the entire filament ; (5) sharp tremblings or concussions ; and (6) a radiating arrangement of the entangled filaments.

Most of the species of the typical genus *Oscillaria* Bosc. grow in dense slimy tufts attached to other algæ or floating bodies, the filaments being not more than from 2 to 6 μ in diameter ; in a few species,

such as O. princeps (Vauch.), the diameter is much greater, the separate filaments, just visible to the naked eye, floating on the surface of the water. *Lyngbya* Ag. is distinguished by its property of forming 'persistent cells,' the function of which is not known; they may possibly be propagative spores. The species have a much less active motion than those of Oscillaria, and chiefly inhabit salt or brackish water. *Symploca* Ktz. grows in tufts, frequently among grass with the habit of a Rivularia. In *Microcoleus* Desm, and *Inactis* Ktz. a large number of filaments are enclosed in the same gelatinous sheath.

The genera Chamæsiphon (A. Br.), Clastidium (Kirch.), Cyanocystis (Bzi.), and Dermocapsa (Crouan) (to which Thuret adds Xenococcus and Sphærogonium, Rostaf.) constitute Borzi's family of *Chamæsiphonaceæ*, distinguished, according to that author, by the presence of *coccogones*, propagative cells of the nature of sporanges, in which *conids* are formed by repeated



FIG. 370.—*Spirulina tenuissima* Ktz. (× 400). (After Cooke.)



FIG. 371.—*Lyngbya æstuarii* Liebm. (× 200). (After Hauck.)

division, the usual number in each coccogone being four, eight, or sixteen. Clastidium (Jahrhft. vaterl. Naturk. Würtemberg, 1880, p. 135) is characterised by each filament having a terminal bristle. Dermocapsa and Xenococcus are epiphytic on Catenella, Lyngbya, and other marine algæ. The former genus has been placed among both Florideæ and Fucaceæ, owing to its mode of propagation; D. violacea (Crouan) has a bright red colour.

Plaxonema Tangl (Sitzber. Akad. Wiss. Wien, 1882) is a filamentous protophyte with the habit of an Oscillaria, but characterised by the presence of a disc-shaped chromatophore in the blue-green protoplasm. Under certain conditions the filaments break up into zoogloea-like colonies. *Borzia* Cohn (Jahrber. Schles. Vaterl. Cultur, 1883, p. 226) is a genus of Oscillariaceæ with the habit of a bacillus, consisting of

short oblong rods, which oscillate slowly and are not enclosed in a gelatinous sheath. In B. trilocularis (Cohn) each hormogone is composed uniformly of three pseudocysts only.

Many of the Oscillariaceæ enter largely into the composition of the blue-green scum seen on the surface of stagnant ditches, &c. Together with others of the Nostochineæ they are said to have the power of decomposing vegetable matter, and to this is largely due the foul stench of stagnant water. In addition to many species of Lyngbya a few belonging to other genera grow in salt or brackish water. Several species of Oscillaria are found in thermal springs.

FIG. 372. *Symploca hydnoides* Ktz. (*a*, natural size; *b*, × 200). (After Hauck.)

The relationship of the genera of Oscillariaceæ to one another, and even to genera at present included in other families, is still very obscure. Hansgirg believes that many of the forms described as species of Chroococcus result from the breaking up of filaments of Lyngbya; while, on the other hand, most of Kützing's species of Leptothrix, and many of Oscillaria, may be simply hormogones of species of Rivulariaceæ, Scytonemeæ, and Stigonemeæ, propagating by frequent divisions, and becoming invested in a more or less thick gelatinous sheath. Many species of Lyngbya may be only the young stages of development of those species of Calothrix and Scytonema to which they are found attached. Oscillaria or Lyngbya antliaria (Hansg.) he now regards as in reality an Aphanocapsa. The same author further states that in the



FIG. 373.—*Symploca violacea* Hauck (× 280). (After Hauck.)

Lyngbyeæ and other of the higher families of Cyanophyceæ, nuclei, pyrenoids, and chromatophores occur, but only when they are in a condition of retrogression from the filiform state, and are breaking up into the unicellular condition. Under the name *Chroomonas Nordstedtii* Hansgirg describes (Bot. Centralbl., xxiii., 1885, p. 229) a biciliated

organism with blue-green endochrome which he regards as the swarm-cell condition of a phycochromaceous alga which occurs normally in a filamentous form, probably as Oscillaria tenuis (Ag.) or O. Frölichii (Ktz.).

LITERATURE.

Fresenius—Ueb. d. Bau u. d. Leben d. Oscillarieen, 1845.
Braun—Bot. Zeit., 1852, p. 395.
Bornet and Thuret—Notes Algol., fasc. 1, pp. iii.-iv. ; and fasc. 2, pp. 132–135.
Zukal—Oesterr. Bot. Zeitschr., 1880, p. 11.
Hansgirg – Oesterr. Bot. Zeitschr., 1884, pp. 313 *et seq.* ; and Ber. Deutsch. Bot. Gesell., 1885, p. 14.

Sub-class 2 and ORDER 5.—Chroococcaceæ.

The Chroococcaceæ share with the Schizomycetes the distinction of being among the lowest forms of vegetable life. The separate cells are always microscopic, and are filled with a blue-green or violet endochrome which owes its colour to the phycocyanin dissolved in the cell-sap ; they contain neither distinct chlorophyll-grains nor starch, nor, except in Chroodactylon (Hansg.), a distinct nucleus. The cells are either isolated, or are more often connected together into colonies by a mucus formed from the disintegration of the outer layers of the cell-wall ; they are never united into a filament. This gelatinous envelope is either colourless and hyaline, or of a blue, brown, or olive colour, and is often strongly lamellated. In Chroococcus (Näg.) it is homogeneous and capable of swelling greatly ; in Glœocapsa (Ktz.) it is composed of two successive layers, and becomes eventually, in some species, crustaceous, and of a very dark brown or even black colour. The internal *pseudocysts* or gonids are never endowed with cilia, as in some Protococcaceæ, and are usually quiescent ; but in Microcystis (Ktz.) they have a constant 'swarming' motion within the hyaline envelope. The entire organism has usually a power of slow spontaneous motion. Multiplication by swarm-spores or zoospores is unknown except in the doubtful case of Merismopedia (Mey.) (Goebel, 'Outlines of Classification,' p. 22). *Resting-spores* or *cysts* (akinetes) are formed in Glœocapsa by the cells of which a colony is composed investing themselves, while still within their common gelatinous envelope, in a rough or spiny coat of



FIG. 374.—Stages in the development of *Chroococcus turgidus* Näg. (greatly magnified). (After Reinke.)

cellulose ; the spiny resting-spores thus formed reproduce the colony by division after a period of quiescence.

With the above exceptions the only mode of propagation in the Chroococcaceæ is by division—repeated bipartition of the cell, which may take place in one, two, or three directions. This is usually accompanied by the disappearance of the separate gelatinous envelope of each individual cell ; but in *Glœocapsa* these still remain, and as many as three or four generations of families may be enclosed within the original envelope, each surrounded by its own investment. In *Chroococcus* Näg. it is not unusual for the individual cells to be entirely isolated within the common envelope. In *Synechococcus* Näg. division takes place in one direction only, and the derivative cells remain attached to one another



FIG. 375.—*Aphanothece microscopica* Näg. (× 70). (From nature.)



FIG. 376.—*Glœothece granosa* Rabh. *A*, gelatinous colony (magnified); *B*, cells (× 250). (After Cooke.)

in a string ; but the attachment is very loose, and soon ceases. In *Merismopedia*, *Tetrapedia* Reinsch., and *Glœochæte* Lagerh. division takes place in two directions, the result being the formation of a plate of cells, often of great regularity. In *Chroococcus*, *Glœocapsa*, *Glœothece* Näg., *Aphanocapsa* Näg., *Aphanothece* Näg., *Microcystis*, and most other genera, division takes place in all three directions. In *Clathrocystis* Henf. the gelatinous envelope, which is of great extent, is broken up into clathrate segments. In *Cœlosphærium* Näg., a common organism in bog-pools moving about with considerable rapidity, it is lobed at the margin, the pseudocysts appearing like blue-green projections on the surface of the globe. *Chroodactylon* Hansg. (Ber. Deutsch. Bot. Gesell., 1885, p. 14) is distinguished by the formation of cell-families branching in an arborescent manner, by its distinct cell-nucleus, and

by the star-shaped chromatophores which enclose moderately large pyrenoids. It has possibly been erroneously referred to this family.

Most of the genera and species of Chroococcaceæ grow in moist situations, as on damp rocks, where they frequently form large shining blue-green mucilaginous masses ; others swim freely on the surface of bog-pools ; a few are found in salt water, attached to sea-weeds. The gonids or algal constituents of many lichens have been shown to be protophytal organisms belonging to the Chroococcaceæ. As it is highly probable that many forms at present included in the family are stages in the history of development of more highly organised protophytes, or even of algæ, their place in a final system of classification is altogether uncertain until their life-history has been more thoroughly investigated. Many are closely analogous to corresponding forms among the Protococcaceæ, as Chroococcus to Chlorococcum, Glœocapsa to Glœocystis,



FIG. 377.—*Microcystis marginata* Men. (× 400). (After Cooke.)



FIG. 378.—*Cœlosphærium Kützingianum* Näg. (× 400). (After Cooke.)

Aphanocapsa to Protococcus, Cœlosphærium to Botryococcus, and Merismopedia to Tetraspora ; but they are probably merely parallel series of forms without any direct genetic connection. Richter, however, identifies Glœocapsa and Glœocystis.

The same observer suggests also a genetic affinity between the various genera usually included under the Chroococcaceæ of the following nature. The lowest form is the naked Aphanocapsa-condition, corresponding to Palmella among the Protococcaceæ. From this naked or only slightly encysted condition is developed the Glœocapsa- or Glœocystis-form, with several gelatinous envelopes, the Chroococcus-condition, where the investment is altogether wanting, and the cœnobe- or Cœlosphærium-condition, where there is only a slight vesicular envelope. The Glœocapsa-form is especially adapted for exposure to air and growth upon a comparatively dry substratum ; the cœnobe-type is developed in water or on a moist substratum in the air. With this is connected the cylindrical form, a higher stage, because it displays differentiation in the direction of growth, and a development towards

the filiform condition. This cylindrical condition, when attained, is usually unstable, but becomes stable in Synechococcus. Glœocapsa may also pass into an encysted filiform condition in Sirosiphon (see p. 439). Zopf insists on the close affinity between the blue-green Schizophyceæ and the Schizomycetes. Hansgirg regards many of the forms included under Chroococcus as resulting from the breaking up of filaments of the higher Cyanophyceæ such as Lyngbya, while Glœocapsa may be derived in the same way from Stigonema, and Synechococcus from Calothrix. He believes, in fact, most if not all of the organisms hitherto included in this family to be connected, by retrogressive metamorphosis, with other more highly developed forms, and even possibly in some cases (Flora, 1886, p. 291) with the protoneme of a moss. Microcystis is regarded by Richter (Hedwigia, 1885, p. 18) as a resting-form of Euglena. The Chroococcaceæ, like the other Cyanophyceæ and the Protococcaceæ, enter largely into the composition of Lichens. The reader will find this subject amply treated by Bornet in his 'Recherches sur les gonidies des Lichens' (Ann. Sci. Nat., 5 Sér., xvii. and xix.).

LITERATURE.

Nägeli – Gattungen einzelliger Algen, 1849.
Borzì—Nuov. Giorn. Bot. Ital., 1878, p. 369; and 1879, p. 47.
Richter—Hedwigia, 1880, pp. 154, 169, 191.
Zopf—Bot. Centralbl., x., 1882, p. 32; and Ber. Deutsch. Bot. Gesell., 1883, p. 319.
Tangl—Anzeiger Akad. Wiss. Wien, 1883, p. 87.
Hansgirg – Oesterr. Bot. Zeitschr., 1884, pp. 313, 351, 389; and Bot. Centralbl. xxii. and xxiii., 1885.

GROUP II. AND CLASS XXVIII.—SCHIZOMYCETES (BACTERIA).

Though this group has been the subject of a great deal of investigation and much speculation, it cannot be said that our knowledge of it is in due proportion to the literature. The minute size of the cells precludes an exact study of their structure. This leads to errors of determination and to confusion of forms in culture experiments, and thus renders difficult the study of the course of development. A large number of the investigators have been and are unequipped with a knowledge of natural history, and are unfitted to appreciate the significance of phenomena observed, or, in other cases, are incapable of observation of

the kind at all. Records of the successive occurrence of different forms in the same situation have been substituted for direct observations of continuity, while errors of even grosser kind abound in the vast literature of the subject. Bacteria are the present refuge of those who believe in 'spontaneous generation,' just as higher forms of organised beings were the subject of their speculations in former times, when the instruments of investigation were less perfect.

Bacteria are either single minute cells of roundish form, or cylindrical and rod-like cells or rows of cells. As has been said, their very minute size has prevented our attaining an exact knowledge of the cell structure. The cell-cavity is ordinarily filled with homogeneous protoplasmic contents. Chlorophyll has been discovered tingeing the protoplasm in three forms—Bacterium viride and Bacillus virens of Van Tieghem, and more faintly in Bacterium chlorinum of W. Engelmann. A red colouring matter discovered by Lankester, and named by him *bacteriopurpurin*, tinges the protoplasm of Beggiatoa roseo-persicina (Zopf), but though colours are often associated with masses of Bacteria, it is difficult to discern in the magnified view the exact seat of it, whether it occur in cell-contents, cell-wall, or substratum. In some forms (which do not contain chlorophyll) a substance resembling starch is found. No one has yet detected nuclei with absolute certainty. The cell-membranes are very delicate, and in such cases as Spirillum (Ehrenb.) highly elastic, but it happens to most Bacteria at one stage of their development that gelatinous outer layers are formed, which either invest single cells and cell-groups, or unite into masses large numbers of cells. A great number of Bacteria have the power of free movement. During such movement rotation takes place round the longitudinal axis, while movements of oscillation also occur in other forms. Cilia or flagella are found in some, but not all, of these moving Bacteria, and it has not been proved that they are motile organs as one might too readily infer. As a matter of fact it is not known, in many cases, whether these flagella are parts of the membrane or of the protoplasm protruded through it; and since they are not always present in motile Bacteria, they need not be regarded as essential organs of locomotion at all events.

Various growth-forms occur which were at first associated with different Bacteria and received generic names. Individual Bacteria are either roundish or in the form of straight rods, or of twisted rods. As de Bary has remarked, 'a billiard ball, a lead pencil, and a corkscrew so exactly illustrate these three chief forms,' that there is no need of models to convey instruction in this respect. The round growth-forms are termed *Coccus* (*Micrococcus* Cohn &c.); the rod-like forms include those

which have been termed *Bacillus* Cohn (long rods), and specially *Bacterium* Cohn (short rods); the shortly coiled forms are known as *Vibrio* Cohn; and the spiral forms have received the names of *Spirillum* Ehrenb., *Spirochæta* Ehrenb., &c., and the very elongated filiform forms are *Leptothrix* Ktz., *Beggiatoa* Trev. &c. *Involution forms* are swollen bladder-like structures of irregular outline probably produced by malnutrition. These growth-forms of cells or of individuals either occur free, or in the form of filaments, or more seldom of flat surfaces or cube-like packets. Large gelatinous masses called *Zoogloea*, composed of numerous individuals of these growth-forms, occur in various situations such as the surfaces of fluids and solids, or they may be found suspended in fluids.



FIG. 379.—*Bacillus Fitzianus* Zopf. Transition forms from round cocci to rods. *d*, with spores (× 4,000). (After Buchner.)

Such forms of Bacteria are grouped into two divisions, viz. those which form their spores endogenously, the *Endosporous Bacteria*, and those which have no such mode of forming spores, the *Arthrosporous Bacteria*. This classification, which can hardly be regarded as finally satisfactory, corresponds, at all events, with the state of our knowledge of the course of development of Bacteria.

Endosporous multiplication is accomplished by the formation within a cell of a minute, granule-like body, which gradually enlarges, while the surrounding protoplasm disappears until it reaches its mature form

within the wall of the mother-cell. The mature *spore* is usually a highly refringent body with definite outline, and of globular or ellipsoidal form. The formation of spores commonly takes place when the substratum yields no more nourishment, or vegetation is otherwise interrupted, and it usually occurs in most of the cells, others remaining sterile. *Arthrosporous* reproduction is effected by the simple separation of members which form the starting point of new growths.

The spores of Bacteria are capable of germination from maturity onwards, often for considerable periods. They withstand the operation of external agencies, such as extreme degrees of temperature and the like, with varying success, many of them exhibiting astonishing endurance. Such arthrosporous forms as Beggiatoa, which vegetate in water, are probably incapable of withstanding desiccation at a very high temperature, but the spores of endosporous Bacteria possess, many of them, remarkable powers of endurance. The spores of Bacillus Anthracis (Cohn) (the cause of splenic fever) remain alive in absolute alcohol. They may be kept for at least three years in an air-dry state, and for at least one year in water, and probably for longer either in air or in water. They were found by Brefeld to withstand boiling in a nutrient solution for a quarter of an hour, the greater part of them for half an hour, a smaller number for one hour, but none for three hours. And so with the spores of other forms. The temperatures at which germination takes place, the minimum, the optimum, and the maximum, vary with different forms; but for the most part the minimum and certainly the optimum may be said to be above the ordinary temperature of a room. Similarly the optimum temperature for vegetation is usually high, being about 30°C., speaking very generally.



FIG. 380.—*Bacillus Megaterium* de By. *a*, outline of a motile chain of rods; *b*, a pair of same; *b*, a quadricellular rod after treatment with alcoholic solution of iodine; *c*, a five-celled rod before spore-formation; *d*—*f*, successive stages of pair of rods while forming spores, about an hour interval between each—state *d* was about two o'clock afternoon, and the spores in *f* were ripe towards evening; *r*, a quadricellular rod with ripe spores; g^1 a five-celled rod with three ripe spores placed in a nutrient solution after several days' desiccation at 12.30; g^2, same about 1.30; g^3, the same about 4 o'clock; h^1, two spores with the walls of the mother-cells dried, and then placed in a nutrient solution about 11.45; h^2, the same about 12.30; *i*, *k*, *l*, later stages of germination; *m*, a rod formed from a spore placed eight hours before in a nutrient fluid, and in the act of splitting transversely. (*a* × 250, the other figures × 600.) (After de Bary.)

With reference to their behaviour towards the supply or exclusion of

FIG. 381.—*Beggiatoa alba* Trev. 1, group of filaments. 2—5, filaments of varying thickness, 5 breaking up; small dark circles in all cases grains of sulphur; where such are abundant transverse segmentation is indistinct. 6 8, filaments rich in sulphur showing transverse segmentation after treatment with solution of methyl violet; in 8 longitudinal division is shown (formation of cocci or spores). 9, filaments which have broken up into spores. 10, motile spores. (1 × 540, other figures × 900.) (After Zopf.)

oxygen, there is great variation among Bacteria. This variation extends from those forms, called *aerobiotic* by Pasteur, which require a plentiful supply of free oxygen for the purpose of vegetation (e.g. Bacillus subtilis, Cohn), to others (*anaerobiotic*) in which vegetation is promoted by its exclusion (e.g. Bacillus Amylobacter, Van Tiegh.). Intermediate forms occur between such extremes; and Nägeli has shown that aerobiotic forms continue to vegetate when the supply of free oxygen ceases.

Speaking in general terms, what has been said already of the mode of life of Fungi in other respects, e.g. nutritive adaptation, holds good of such Bacteria as contain no chlorophyll. They are saprophytes exciting fermentations and producing combustions of the substratum, and putrefactive processes; or parasites, though very rarely on living plants, it may very well be on account of the acid reaction. As parasites in living animals they obtain the greatest share of our interest, since, as everyone knows, it is sought to connect them with a large number of diverse diseases. That this attempt is made with the greatest rashness in many cases, on utterly insufficient data, often on the mere presence of some vaguely determined form in diseased tissues, is a scandal of medical literature. On the other hand it has been thoroughly proved in certain cases that their presence and action have the character of exciting causes of disease, and it cannot be doubted that painstaking research will bring to light numerous other instances of equal weight. Slipshod research will only retard progress in this direction. It has already done much in obscuring results, and in placing obstacles in the path. The most noteworthy feature, as de Bary has pointed out, in the parasitism of Bacteria in the living bodies of animals is their *facultative* parasitism—(as illustrated for example in the well-established case of Bacillus Anthracis in anthrax or splenic fever), a matter of grave importance from the medical point of view.

Among saprophytes may be mentioned Bacterium Termo (Duj.), an exceedingly abundant agent of putrefaction; Bacillus Megaterium (de By.) (fig. 380), and Beggiatoa alba (Trev.) (figs. 381, 382), the 'sewage-fungus' of engineers, found in sulphuretted waters, the effluents from manufactories and sewage-works, which has a remarkable power of extracting sulphur from the water, and storing it up in the form of minute refringent globules.

As regards the position of Bacteria, 'to say that they are offshoots of the Fungi is to "contradict all trustworthy observations" so flatly, that the view need not be seriously discussed in this place' (de Bary, 'Comp. Morph.,' &c., p. 474). They are only fungi in the very limited sense of their being 'thallophytes which contain no chlorophyll,' and indeed it

has been seen that certain Bacteria do contain chlorophyll. Looking at their morphological characters, so far as these are at present known to us, it cannot be doubted that the nearest allies of the Arthrosporous Bacteria are those Protophyta, Nostocaceæ, Oscillariaceæ, Chroococcaceæ, &c., which contain chlorophyll. Leuconostoc (Van Tiegh.) has already been mentioned (p. 433) as an intermediate form. A gap certainly exists between Arthrosporous and Endosporous forms ; but so far as



FIG. 382.—*Beggiatoa alba* Trev. Curved and spiral forms. *A*, attached filaments. *B—H*, portions of spiral filaments ; *H* showing separate cells, *E* swarming ('spirillum') with a cilium at each end. Small dark circles are sulphur granules (× 540). (After Zopf.)

can be seen the latter stand nearer to the former than to any other group, and the interval which separates them may become narrower with farther knowledge. On the other side a connection appears to be indicated between Bacteria and the Flagellata ; but more than this one

can hardly say. In conclusion, and as summing the matter up, the words of de Bary ('Comp. Morph.,' &c., p. 475) may be quoted. 'If we assume for a moment a connection between the Bacteria and the Flagellata, it is evident that as a consequence the following series of forms converge to the Flagellata : *firstly*, the series of Bacteria and the Nostocaceæ ; *secondly*, that of the Mycetozoa ; *thirdly*, that of the chlorophyllaceous Algæ, with which are connected in ascending line the main series of the vegetable kingdom and of the Fungi as one or more lateral branches. . . . *fourthly*, and lastly, the Rhizopoda and the Protozoa with the animal kingdom, which connects with these in an ascending line.'

LITERATURE.

De Bary -Vorlesungen über Bacterien (Leipzig, 1885 and 1886). (See English translation by Garnsey and Balfour, Oxford, 1887.)

The above contains an admirable guide to the literature of the subject. (See also the same author's Comparative Morphology, &c.)

Cornil et Babes—Les Bactéries, &c., 2nd ed. (Paris, 1886).

Crookshank—Introduction to Practical Bacteriology (London, 1886).

Duclaux—Chimie biologique (Paris, 1883).

Grove—A Synopsis of the Bacteria and Yeast Fungi (London, 1884).

Klein—Micro-organisms and Disease, 3rd ed. (London, 1886).

Hueppe—Die Formen der Bacterien (Wiesbaden, 1886).

Hueppe—Die Methoden der Bacterienforschung (Wiesbaden, 1885).

Zopf—Die Spaltpilze (Breslau, 1884).

The references to the vast literature of the subject in the above books will be a sufficient guide to the most ardent student. At the same time it would be unpardonable to abstain from a special reference to the labours of Cohn, Pasteur, Koch, Lankester, Brefeld, Van Tieghem, Prazmowski, Naegeli, and Lister, since these are of fundamental importance.

INDEX.

(*The figures in large type refer to Illustrations.*)

ABI

Abies, 3
Abjunction, 312, 275
Absidia (Van Tiegh.), 338, 339
Absorbing system, 138
Acanthococcus (Lagerh.), 417
Acetabularia (Lmx.), 286, 251, 252
— mediterranea (Lmx.), 251, 252
Achlya (Nees ab Esenb.), 335, 346, 291, 292
— polyandra (Hildebr.), 291
— racemosa (Hildebr.), 291
Achnanthes (Bory), 426
— brevipes (Ag.), 457
Acrasieæ, 401, 405, 406
Acrasis (Van Tiegh.), 405
Acroblaste (Reinsch), 280
Acrocarpi, 149
Acrogynous Jungermanniaceæ, 161
Acrospore, 330, 343, 352, 361, 362, 363, 364, 365, 367, 368, 369, 370, 372, 373, 374, 376
Acrosticheæ, 80, 83
Acrostichum (L.), 85
Adder's Tongue, 100
Adiantum (L.), 85, 44, 50, 58
— capillus-Veneris (L.), 44, 50
Adventitious bud, 77, 81, 100, 49
Æcidia, 384, 385, 386, 314
Æcidiospore, 6, 384, 385, 386, 314
Æcidium (Pers.), 383
Ægopodium Podagraria, 348, 349
Aerial leaf, 28, 7, 8
Aerobiotic Bacteria, 454
Æthalium septicum (Fr.), 403
Agar-agar, 210
Agaricini, 393, 394, 270, 273, 318–321
Agaricus (L.), 4, 392, 393, 270, 273, 319, 320
— campestris (L.), 320
— cryophilus (Bull.), 273
— Emerici (Berk.), 316
— Gardneri (Berk.), 316
— igneus (Tul.), 316
— lampas (Berk.), 316
— melleus (L.), 309, 310, 316, 392, 270, 319
— noctilucens (Lév.), 316
— olearius (DC.), 316
Agarum (Grev.), 244
Aglaozonia (Zan.), 251
— reptans (Ktz.), 252
Air-bladder, 232, 236, 241, 244, 206, 207, 211

ANO

Air-cavity, 102, 105, 109, 147, 167, 169, 24, 110, 112, 153, 154
Akinete, 185, 258, 274, 446
Alaria (Grev.), 239, 244
— esculenta (L.), 244
Alcoholic ferment, 380
Algæ, 2, 3, 4, 184, 169–265
Algal cells of Lichens, 318, 319, 370, 279–282, 307, 308
Alsophila aculeata (Klotzsch), 57
Alternaria (Nees ab Esenb.), 374
Alternation of generations, 16, 182, 189, 214, 263, 325, 440
Amanita, 393
Amansia (Lmx.), 194, 209
Amœba, 406
Amœboid motion, 196, 199, 217, 274, 284, 344, 401, 402, 403, 405, 406
Amphigaster, 160, 138, 139
Amphipleura (Ktz.), 426
Amphiroa (Lmx.), 206
Amphithecium, 145
Amphithrix (Ktz.), 437
Amphora (Ehrb.), 426
Amylum-star, 176
Anabæna (Bory), 430, 432, 433, 360
— flos-aquæ (Fr.), 360
Anacrogynous Jungermanniaceæ, 161
Anadyomene (Lmx.), 289
Anaerobiotic Bacteria, 454
Ancylisteæ, 4, 330, 344, 345
Ancylistes (Pfitz.), 331
— Closterii (Pfitz.), 330
Andreæa (Ehrh.), 145, 150, 123, 124
— alpestris (Schmp.), 123, 124
Andreæaceæ, 136, 150, 123, 24
Androspore, 6, 225, 202
Aneimia (Sw.), 78, 90, 91, 43
— Phyllitidis (Sw.), 43
Aneura (Dum.), 164
Angiopteris (Hoffm.), 91, 94, 95, 71
— caudata (De Vriese), 71
— evecta (Hoffm.), 93
Angiosperms, 12, 13
Annularia (Brongn.), 126, 101
Annulus, 74, 78, 86, 89, 110, 122, 146, 393, 55, 56, 83, 110, 112, 319, 320
Anomœneis spherophora (Pfitz.), 354A

ANT

Antherid, 7 (see also under Vasc. Crypt., Musc., Char., Algæ, and Fungi), 11, 30, 36, 43, 44, 65, 74, 78, 106, 108, 130, 141, 157, 163-165, 175-177, 192, 199, 201-203, 208, 210, 248, 259, 286, 290, 291, 303-306, 314
Antheridial tube, 291
Antherozoid 7 (see also under Vasc. Crypt., Musc., Char., Algæ, and Fungi), 11, 15, 17, 20, 30, 44, 78, 108, 130, 157, 163, 199, 201, 210, 248, 259, 290
Anthoceros (L.), 157, 165, 144, 145
— lævis (L.), 144, 145
Anthoceroteæ, 159, 164, 144, 145
Antipodal cells, 15
Aphanizomenon (Morr.), 430, 432, 361
— flos-aquæ (Morr.), 432, 361
Aphanocapsa (Näg), 412, 440, 445, 447 448
Aphanomyces (de By.), 334, 335
Aphanothece (Näg.), 447, 375
— microscopica (Näg.), 375
Aphlebia, 120, 92
Apical cavity, 179
— cell, 23, 41, 74, 90, 98, 102, 109, 156, 173, 174, 176, 191, 214, 228, 241, 437, 52, 162
— papilla, 179
Apiocystis (Näg.), 413, 341
— Brauniana (Näg.), 341
Aplanes Braunii (de By.), 334
Aplanospore, 185, 258, 274
Aplolepidæ, 148
Apogamy, 11, 52, 69, 86, 47
Apophyse, 148, 112, 124
Apospory, 11, 69, 86, 48
Apothece, 355, 356, 379, 372, 373, 308
Appendage, 378, 379, 312
Appendiculæ, 363
Aquilegia, 364
Archæocalamites (Stur), 126, 99
— radiatus (Stur), 99
Archegone, 8 (see also under Vasc. Crypt., Musc., Char., and Algæ), 4, 10, 14, 20, 29, 43, 45, 46, 61, 74, 79, 80, 109, 131, 141, 142, 147, 148, 158, 164, 165, 166
Archegoniophore, 86, 61
Archespore, 13, 20, 36, 60, 80, 99, 111, 134, 135, 144, 146, 159, 33, 56, 110
Archidium (Brid.), 145, 150
Areolation, 420
Arthrocladia (Duby), 245
Arthrodonteæ, 148
Arthropitys (Göpp.), 126
Arthrosporous Bacteria, 451, 452, 455
Artotrogus (Mont.), 325
Ascobolus (Pers.), 356, 359, 360, 361, 368, 369, 370, 372, 378, 305
— furfuraceus (Pers.), 305
Ascogenous hyphæ, 353, 355, 360, 364, 366, 367, 368, 369, 370, 372, 373, 374, 305, 306, 308
Ascogone, 360
Ascomycetes, 4, 312, 353, 383, 386, 267, 268, 271, 276, 278, 300-313
— course of development, 361
— homologies of the organs, 377
— (doubtful), 4, 378, 312, 313
Ascophylla (Stackh.), 235
Ascospore, 6 (see also under Ascomycetes), 268, 276, 278, 300, 304, 311, 312, 313
Ascotricha (Berk.), 370
Ascus, 353 (see also under Ascomycetes), 276, 300, 303, 305, 308, 311, 312
Aseroe (La Bill.), 398, 331
— rubra (Berk.), 331
Aspergillus glaucus (Lk.), 366
Asperococcus (Lmx.), 241, 245, 218
— bullosus (Lmx.), 218

BAT

Aspidieæ, 83
Aspidium (Sw.), 85, 42, 55, 58
— filix-mas (Sw.), 72, 76, 77, 82, 42, 55
Aspleniеæ, 83
Asplenium (L.), 69, 85, 49, 51, 56, 58
— Adiantum-nigrum (L.), 51
— bulbiferum (Forst.), 81
— decussatum (Sw.), 49
— Trichomanes (L.), 56
Assimilating tissue, 138, 193
Asterophyllites (Rev.), 126, 130, 101
Asterotheca (Presl), 122, 94
— Sternbergii (Stur), 94
Astromyelon (Williams.), 126
Athyrium filix-fœmina (Bernh.), 69, 76
Atrichum undulatum (P. B.), 102, 119
Attachment-disc, 228
Aulacodiscus (Ehrb.), 426
Aulacomnion (Schw.), 149
— androgynum (Schw.), 140
Aulosira (Kirchn.), 430, 433
Auricle, 92, 161, 140
Auxiliary cell, 203
Auxospore, 6, 422, 423, 355
Azolla (Lam.), 19, 26, 31, 115, 12, 13
— caroliniana (Willd.), 13
— filiculoides (Lam.), 12
Azygosperm, 338, 340, 342, 343
Azygites (Fr.), 338

BACILLARIACEÆ, 419
Bacillus (Cohn), 440, 451, 379, 380
— amylobacter (Van Tiegh.), 454
— anthracis (Cohn), 452, 454
— Fitzianus (Zopf), 379
— Megaterium (de By.), 454, 380
— subtilis (Cohn), 454
— virens (Van Tiegh.), 450
Bacteria, 305, 306, 449
— conditions of vegetation, 452
— growth-forms of, 450
— mode of life, 454
— movements of, 450
— spores of, 451, 452
Bacteriopurpurin, 450
Bacterium (Cohn), 451
— chlorinum (Engelm.), 450
— Termo (Duj.), 454
— viride (Van Tiegh.), 450
Bæomyces (Pers.), 361
Balsamia (Vitt.), 358
Bangia (Lyngb.), 190, 191, 216, 193-195
— ceramicola (Chauv.), 193
— fusco-purpurea (Lyngb.), 194, 195
Bangiaceæ, 216
Barberry, 383, 384, 385
Barbula (Hedw.), 140, 147, 149, 105
Barilla, 190
Bartramia (Hedw.), 149
Basal cell, 68
Basid, 312, 384, 385, 388, 389, 390, 393, 394, 275, 317, 321
Basidiomycetes, 4, 306, 312, 388, 266, 270, 273, 316-331
Basidiomycetous Lichens, 319
Basidiospore, 6, 312, 388, 389, 390, 275, 317, 321, 322, 325, 329
Basilar cell, 433, 365
Batarrea (Pers.), 308, 397, 325, 326
— Steveni (Fr.), 325, 326
Batrachospermeæ, 211, 189-191
Batrachospermum (Bory), 191, 211, 212, 213, 214, 189
— moniliforme (Roth), 189

BEA

Beam, 226, 204
Beer-yeast, 268, 313
Beggiatoa (Trev.), 4, 451, 452, 381, 382
— alba (Trev.), 454, 381, 382
— roseo-persicina (Zopf), 450
Biddulphia (Gray), 426
Binuclearia (Witt.), 276
Bladder-wrack, 230, 236
Blasia (Mich.), 156, 157, 164
Blechnum (L.), 85, 58
Blyttia (Endl.), 160
Bog-mosses, 151
Boletus (L.), 392, 396
Bonnemaisonia (Ag.), 201
Bornetia (Thur.), 194, 209
— secundiflora (Ag.), 194
Bornia (Brongn.), 126
Borzia (Cohn), 428, 444
— trilocularis (Cohn), 445
Bothrodendron (L. and H.), 116
Botrychium (Sw.), 97, 98, 99, 100, 73–75
— Lunaria (Sw.), 97, 98, 100, 73–75
Botrydiaceæ, 280, 285, 290
Botrydina (Bréb.), 186, 412
Botrydium (Wallr.) 186, 284, 285, 410, 412, 250
— granulatum (Wallr.), 285, 250
Botryococcus (Ktz.), 186, 418, 448, 349
— Braunii (Ktz.), 349
Botryopteris (Ren.), 122
Botrytis cinerea (Pers.), 374
Brachytrichia (Zan.), 437
Bract, 173, 163–165
Bracteole, 173
Brake, 82
Breaking of the meres, 432
Brittle-worts, 181
Brownian movement, 269
Bruckmannia (Stbg.), 127
Bryaceæ, 136, 146, 102–120
Bryum (L.), 145, 149, 115
— argenteum (L.), 115
Bryopsideæ, 289
Bryopsis (Lmx.), 289, 290
Bud, 133, 181, 196, 214, 250, 252, 162
Bulbil, 61, 67, 176
Bulbochæte (Ag.), 188, 222, 226, 202
— setigera (Ag.), 202
Bundle-sheath, 44, 109
Bursulla (Sorok.), 405
Buxbaumia (Hall.), 149

CABBAGES, 326
Calamarieæ, 125, 130, 96–100
Calamites, 125, 126, 97
Calamitina (Weiss.), 126, 98
Calamocladus (Schmp.), 126
Calamodendreæ, 125
Calamodendron (Brongn.), 125, 126
Calamostachys (Schmp.), 125, 127
Calcareous incrustation, 181, 195, 206, 211, 286, 304, 403, 440, 167
Calcium oxalate, 85, 308
Callithamnion (Lyngb.), 4, 189, 191, 194, 204, 180
— caudatum (Ag.), 194
— seirospermum (Griff.), 204, 80
— versicolor (Drap.), 204
Callus, 240, 244, 217
Calothrix (Ag.), 435, 436, 445, 449, 364
— crustacea (Thur.), 364
Calotrichaceæ, 433
Calypogeia (Radd.), 162, 164, 134
— Trichomanis (Cord.), 134
Calypter, 134, 144, 110, 114, 117, 131,

CHI

Campylodiscus (Ehrb.), 426
Canal, 143, 109
Canal-cell, 69, 133, 45
Capillitium, 358, 396, 397, 404, 326, 334
Capitulum, 177, 163
Capsella bursa-pastoris, 326
Capsosira (Ktz.), 439
Carinal canal, 106, 108
Carpogenous cells, 176, 179
Carpogone, 8, 185, 199, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 385, 179, 303–307
Carpomitra (Ktz.), 241
Carposperm, 185, 201, 203
Carpospore, 185, 201, 222, 263, 179, 199
'Carposporeæ, 3
Carpostome, 201
'Carrageen moss,' 210
Casuarina, 113
Catharinea undulata (W. & M.), 102
Caudex, 71
Caulerpa (Lmx.), 289, 256
— prolifera (Lmx.), 256
Caulerpeæ, 289, 256
Cell-cap, 222, 200
Cellulin, 176
Central bundle, 138
— cavity, 105, 81
— cell, 10, 14, 17, 26, 31, 69, 133, 143, 159, 109, 158
— node, 353
Ceramiaceæ, 196, 204, 172, 179–181
Ceramidium, 201
Ceramium (Lyngb.), 204, 181
— strictum (Grev.), 181
Ceratieæ, 404
Ceratium (Lk.), 404, 335
— hydnoides (Alb. & Sch.), 335
— porioides (Alb. & Sch.), 335
Ceratodon (Brid.), 149
Ceratopteris thalictroides (Brongn.), 77, 81, 52
Ceterach (Adans.), 85
Cetraria islandica (Ach.), 283
Chætangiaceæ, 211
Chætocladieæ, 339
Chætocladium (Fres.), 339, 340, 350
Chætomium (Kze.), 370
— fimeti (Fckl.), 356
Chætomorpha (Ktz.), 276
Chætopeltis (Bert.), 222
Chætophora (Schr.), 273, 275, 276
Chætophoraceæ, 222, 258, 273, 275, 279
Chætopteris (Ktz.), 249
— plumosa (Ktz.), 249
Chætostylum (Van Tiegh.), 339
Chalara Mycoderma (Cienk.), 380
Chamæsiphon (A. Br.), 444
Chamæsiphonaceæ, 441, 444
Chantransia (Fr.), 211, 213, 214, 216, 190, 191
— corymbifera (Thur.), 214, 191
— virgatula (Thur.), 190
Chara (L.), 174, 175, 176, 179, 181, 182, 160–162, 164, 167, 168
— aspera (Willd.), 176
— crinita (Wallr.), 181
— fragilis (Desv.), 176, 160, 162, 164, 168
— hispida (L.), 176, 161, 167
Characeæ, 15, 173, 160–168
Characiaceæ, 410
Characin, 181, 416
Characium (A. Br.), 345, 412, 340
— ornithocephalum (A. Br.), 340
Chareæ, 173, 182, 160–162, 164, 167, 168
Cheilanthes (Sw.), 85, 58
Chiloscyphos (Cord.), 164

CHL

Chlamydococcus (A. Br.), 186, 299, 300, 409, 417
— nivalis (A. Br.), 416
Chlamydomonas (Ehrb.), 186, 299, 300, 409, 417, 419
— Morieri (Dang.), 300
— pulvisculus (Müll.), 300
Chlamydospore, 6, 339
Chlorochytrium (Cohn), 284, 345, 410
Chlorococcum (Fr.), 417, 448, 347
— gigas (Grün.), 417, 347
Chlorogonium (Ehrb.), 300
Chlorophyceæ, 305
Chlorophyllophyceæ, 409
Chlororufin, 279
Chlorothecium (Brzi.), 411
Chlorotylium (Ktz.), 280
Choanephora (Cunn.), 340
Choiromyces (Vitt.), 358
Chondria (Ag.), 209
Chondrioderma difforme (Rost.), 332
Chondrites (Sternb.), 304
Chondrus (Grev.), 208, 185
— crispus (Stackh.), 210, 185
Chorda (Stackh.), 241, 216
— filum (Stackh.), 242, 216
— tomentosa (Lyngb.), 244
Chordaria (Ag.), 247, 220
— flagelliformis (Ag.). 220
Chordariaceæ, 190, 247
Chromatophore, 194, 410, 427, 444, 445
Chromophyton (Wor.), 188, 257
Chroococcaceæ, 409, 419, 427, 446, 455, 374–378
Chroococcus (Näg.), 417, 441, 445, 446, 447, 448, 449, 374
— turgidus (Näg.), 374
Chroodactylon (Hansg.), 446, 447
Chroolepideæ, 187, 258, 273, 279, 247
Chroolepus (Ag.), 280, 284
— aureum (Ktz.), 280
— Iolithus (Ag.), 280
— umbrinum (Ktz.), 280
Chroomonas Nordstedtii (Hansg.), 445
Chrysochytrium (Schroet.), 347
Chrysomyxa (Ung.), 386
Chylocladia (Grev.), 208
Chytridiaceæ, 4, 250, 344, 352, 297
— (doubtful). 347
Chytridium (A. Br.), 4, 249, 346
Cilia (of peristome), 135, 147, 111
Cinclidium stygicum (Sw.), 120
Cingularia (Weiss), 128, 100
— typica (Weiss), 100
Circinate vernation, 72, 51
Circinella (Van Tiegh.), 339
Cladochytrieæ, 344, 345, 346
Cladochytrium (Nowak.), 346, 352
Cladonia (Hill), 361, 279
— furcata (Hoffm.), 279
Cladophora (Ktz.), 275, 276, 330, 243
— gracilis (Ktz.), 243
Cladosporium herbarum (Lk.), 374
Cladostephus (Ag.), 249, 250
Clamp-connection, 306, 266
Clastidium (Kirchn.), 444
Clathrocystis (Henf.), 447
Clathrus (Mich.), 398, 399
Clavarieæ, 391
Claviceps (Tul.), 361, 375, 309–311
— purpurea (Tul.), 375, 309–311
Cleistocarp, 355, 356
Closterium (Nitzsch), 259, 268, 269, 330, 239, 241
— Dianæ (Ehrb.), 239
— rostratum (Ehrb.), 241

COR

Clover, 364
Club-moss, 61
Coccidium, 201
Coccocarpia molybdia (Pers.), 280
Coccogone, 444
Cocconema lanceolatum (Ehrb.), 357
Coccus, 450, 379, 381
Codiolum (A. Br.), 413, 342
— gregarium (A. Br.), 342
Codium (Stackh.), 289, 290
Cœlastrum (Näg.), 186, 303, 265
— cubicum (Näg.), 265
Cœloblastæ, 186, 281
Cœlosphærium (Näg.), 447, 448, 378
— Kützingianum (Näg.), 378
Coemansia (Van Tiegh. and Le Mon.), 341
Cœnobe, 184, 186, 291, 292, 259–265
Cœnobieæ, 186, 291, 412, 418, 419, 259–265
Coleochætaceæ, 188, 190, 219, 220, 199
Coleochæte (Bréb.), 189, 220, 199
— divergens (Prings.), 220
— pulvinata (A. Br.), 220, 199
— scutata (Bréb.), 220
— soluta (Prings.), 220
Coleodesmium (Brzi.), 440
Collateral vascular bundles, 75, 89, 98
Collema (Ach.), 321, 322, 372, 385 307
— pulposum (Bernh.), 307
Collemaceæ, 322, 359, 360, 361, 370, 372, 307 308
Collenchymatous tissue, 93
Colpodella (Cienk.), 405
Columel, 29, 86, 135, 145, 165, 339, 351, 110, 112
Commissure, 92
Completoria complens (Lohde), 343
Compositæ, 364
Concentric vascular bundles, 18, 75, 98, 109
Conceptacle, 11, 196, 199, 207, 232, 290, 182–184, 188, 208, 209
Conducting tissue, 138, 193
Conferva (L.), 185, 273, 274, 276
— bombycina (Ktz.). 273
Confervaceæ, 185, 187, 190, 259, 273, 277, 278, 242–244
Confervites (Sch.), 304
Confervoideæ, 237
— Heterogamæ, 185, 188, 219, 199–204
— Isogamæ, 185, 186, 272, 242–247
'Conid,' 327, 436, 444
Conidiobolus utriculosus (Bref.), 343
Conidiospore, 325, 327, 309
Coniferæ, 13, 310, 3
Conjugatæ, 185, 187, 258, 233–241
Conjugation, 185 (see also under Algæ, Fungi, and Schizophyceæ), 198, 222, 233–238, 241, 246, 247, 252, 293–297, 299
Conjugation-tube, 346, 297
'Connubium,' 263
Consortism, 318
Continuity of protoplasm, 66, 76, 192, 194, 230, 240, 428, 440
Contractile protoplasm, 422
— (pulsating) vacuole, 257, 292, 300, 409, 415
Convolvulus, 364
Coprinus stercorarius (Fr.), 392, 318, 321
Copromyxa protea (Zopf), 406
'Copulation,' 263
Cora (Fr.), 319
Corallina (L.), 189, 194, 206, 183, 184
— officinalis (L.), 206, 183
— rubens (L.), 184
Corallinaceæ, 196, 199, 203, 206, 182–184
Coralline, 195, 206
Cordyceps (Fr.), 376

COR

Coremium (Lk.), 312
— glaucum (Lk.), 367
Cork, 98
Cormophytes, 135, 156, 173
Corn-mildew, 383
Corpusculum, 14, 3
Corsinia (Radd.), 160, 166
Cortex, 173, 174, 221, 247, 39
Cortical lacuna, 81
— tissue, 76, 228, 242, 249, 321, 216, 271, 311
Corticium (Pers.), 391
Coscinodiscus (Ehrb.), 420, 426
Cosmarium (Cord.), 268, 269, 270, 239
— cœlatum (Ralfs), 239
Cotyledon, 17, 28, 41, 55, 71, 105, 8, 9, 14
Craterospermum (A. Br.), 263
Cribrarieæ, 401
Cronartium (Pers.), 386
Crouania (Ag.), 204, 172
— attenuata (Ag.), 172
Crown, 179, 164, 166
Cruciate (tetraspores), 195
Crucibulum (Tul.), 397, 327, 328
— vulgare (Tul.), 327, 328
Cruciferæ, 326, 327
Cruoria (Fr.), 191, 193, 210
Crustaceous Lichens, 321
Cryptogramme crispa (R. Br.), 72
Cryptonemia (Ag.), 208
Cryptonemiaceæ, 208
Crystalloid, 276, 288, 308
Ctenomyces (Eidam), 367
Cupule, 140, 157, 170, 155
Cupuliferæ, 310
Cushion, 93, 115, 87
Cutleria (Grev.), 239, 241, 251, 26, 227
— adspersa (de Not.), 252
— multifida (Grev.), 252, 226, 227
Cutleriaceæ, 187, 239, 251, 254, 226–229
Cyanocystis (Brzi.), 444
Cyanophyceæ, 408, 409, 426, 358–378
Cyanophycin, 428
Cyathea (Sm.), 85, 124, 59
Cyatheaceæ, 71, 73, 78, 80, 81, 85, 123, 124, 57, 59
Cycadeæ, 13, 1
Cyclomyces (Kze.), 392
Cyclopteris hibernica (Forbes), 121
Cyclosis, 175, 422, 163
Cylindrocapsa (Reinsch), 227, 414
Cylindrocapsaceæ, 228
Cylindrospermum (Ralfs), 430, 433, 362
— macrospermum (Ktz.), 362
Cymbella (Ag.), 426
Cymopolia (Lmx.), 288
Cyst, 185, 258, 264, 272, 274, 278, 427, 429, 446
Cystid, 394, 321
Cystocarp, 201, 213, 169–171, 173, 174, 176–178, 183, 184, 188, 189, 191
Cystopteris (Bernh.), 69, 85, 58
— bulbifera (Bernh.), 81
— fragilis (Bernh.), 67
Cystopus (Lév.), 312, 323, 326, 327, 328, 335, 269, 275, 285, 287
— candidus (Lév.), 326, 269, 285
— cubicus (Lév.), 326
— Portulacæ (Lév.), 275, 287
Cystosira (Ag.), 235

Dacryomyces (Nees ab Esenb.) 389
Danæa (Sw.), 94, 95
Danæites (Göpp.), 122
Dasya (Ag.), 196, 209, 174
— elegans (Ag.), 174

ECT

Dasycladaceæ, 280, 281, 286, 251–255
Dasycladus (Ag.), 287, 288, 253
— clavæformis (Ag.), 253
Davallia (Sm.), 85
Davallieæ, 83
Delesseria (Grev.), 193, 208
Delesseriaceæ, 208
Delphinium, 364
Derbesia (Sol.), 289
Dermocapsa (Crouan), 444
— violacea (Crouan), 444
Desiccation of spores, 315, 452
Desmarestia (Lmx.), 241
Desmidiaceæ, 187, 190, 258, 259, 268, 239–241
Desmidium (Ag.), 187, 268, 239, 241
— Swartzii (Ralfs), 239, 241
Desmids, movements of, 269
Diaphragm, 31, 39, 105, 124, 125, 179, 9, 14
Diatoma (DC.), 424, 357
— elongatum (Ag.), 357
Diatomaceæ, 187, 258, 408, 419, 353–357
Diatomin, 257, 421
Diatoms, movements of, 422
Diatrype (Fr.), 373
Dichothrix (Zanard.), 437
Dicksonia (L'Hérit.), 71, 84, 85, 42
— antarctica (Lab.), 42
Dicranum (Hedw.), 138, 149
Dictyonema (Mont.), 319
Dictyopteris (Lmx.), 254
Dictyosiphon (Grev.), 239, 241, 245
— hippuroides (Lyngb.), 247
Dictyosphærium (Näg.), 412, 338
— reniforme (Buln.), 338
Dictyostelium (Bref.), 405
Dictyota (Lmx.), 254
Dictyotaceæ, 188, 237, 254, 230, 231
Dictyuchus (Leitg.), 334, 335
Didymium (Schrad.), 316, 333
— serpula (Fr.), 333
Dimargaris (Van Tiegh.), 341
Diplolepidæ, 148
Diploxylous bundles, 117
Dipsacus, 364
Disc, 170, 184, 220
Discocarp, 355, 369
Discomycetes, 308, 319, 355, 356, 370
Dispira (Van Tiegh.), 341
Diverticula, 209, 187
Docidium (Bréb.), 268, 269, 239
— baculum (Bréb.), 239
Doodia (R. Br.), 85
Dothidea (Fr.), 354
Draparnaldia (Ag.), 273, 275, 276, 417, 244
— glomerata (Ag.), 244
Drilosiphon (Ktz.), 433, 438, 440
— Julianus (Ktz.), 440
Dry-rot, 309
Dudresnaya (Born.), 189, 202, 204, 179
— coccinea (Crouan), 179
Dulse, 210
Dumontia (Lmx.), 208
Dumontiera (Nees ab Esenb.), 171
Duriæa (Bor.), 166
Durvillæa (Bory), 188, 228, 230, 235, 236
Dutch rushes, 113
Dwarf male, 225, 201, 202

Ecklonia (Horn.), 244
Ectocarpaceæ, 187, 239, 247, 212, 221, 222
Ectocarpus (Lyngb.), 187, 239, 241, 247, 221, 222
— investiens (Hauck), 221
— pusillus (Griff.), 249

ECT

Ectocarpus siliculosus (Ktz.), 249, 222
Edible earths, 424
Elachista (Duby), 247
Elaphomyceæ, 358
Elaphomyces (Nees ab Esenb.), 357, 358, 359, 367
Elater, 102, 111, 128, 134, 159, 163, 171, 84, 85, 158, 159
Elk's-horn-fern, 72
Embryo, 11, 13, 17, 27, 40, 56, 105, 134, 180, 271, 21, 45, 158
Embryo-sac, 14, 3
Embryonic vesicle, 11, 14, 3
Emericella (Berk.), 319
Empusa (Cohn), 342, 343
— Grylli (Fres.), 343
— Muscæ (Cohn), 343
Encalypta (Schreb.), 149
Encalypteæ, 148
Encysted condition, 415
Endoclonium (Franke), 284
Endoderm, 76, 109
Endogenous spore, 424, 451, 379-381
Endophyllum (Lév.), 386
Endophytic Algæ, 222
— Fungi, 317
Endosperm, 15, 39, 20
Endosphæra (Klebs), 284, 410
Endospore, 20. 47, 74, 80, 112, 134, 145, 160, 17
Endosporous Bacteria, 451, 452, 455
Endostome, 147
Endothecium, 145
Enteromorphæ (Lk.), 217, 218, 219, 197
— intestinalis (Lk.), 197
Enterosora (Baker), 79
Entomophthora (Fres.), 342, 343
— curvispora (Nowak.), 342
— ovispora (Nowak.), 342
— radicans (Bref.), 342
Entomophthoreæ, 4, 342
Entyloma (de By.), 350, 351, 352
Envelope of plasmode, 403
Envelope-tissue, 355, 360, 361, 363, 364, 366, 368, 370, 373, 374, 378, 384, 303-305
Ephebe pubescens (Fr.), 281
Ephedra, 113
Ephemerum (Hampe), 150, 121
— serratum (Hampe), 121
Epichloe (Fr.), 361, 376
Epiderm, 39, 54, 153-155
Epidermal tissue, 10, 138, 193, 242, 216
Epiphragm, 146, 397, 112, 117, 327
Epiphytic Algæ, 237, 239, 248, 257. 444, 184, 221
— Fungi, 317
Epiplasm, 354
Epispore, 20, 22, 25, 36, 47, 14, 16
Epithemia (Bréb.), 426
Equiseta cryptopora, 107
— phaneropora, 107
Equisetaceæ, 19, 20, 100, 124, 2, 77-86, 96-101
Equisetites (Sch.), 124
— arenaceus (Brongn.), 124
Equisetum (L.), 102, 124, 2, 77-86
— arvense (L.), 106, 107, 109, 110, 78, 79
— giganteum (L.), 113
— hyemale (L.), 110, 113, 82
— limosum (L.), 109, 111, 86
— littorale (Kuhl.), 109
— maximum (Lam.), 110, 78, 81, 83
— palustre (L.), 110
— pratense (Ehrh.), 110
— sylvaticum (L.), 77, 80
Eremascus (Eid.), 378, 302
— albus (Eid.), 359, 361, 302
Eremobiæ, 186, 410, 336-344

FON

Eremosphæra (de By.), 417
Ergot, 376, 309-311
Erysiphe (Hedw.), 317, 362, 363, 367, 578
— communis (Lk.), 364
— graminis (Lév.), 364
— lamprocarpa (Lk.), 364
— Martii (Lév.), 364
— (Uncinula) spiralis (Berk. & Curt.), 364
Erysipheæ, 309, 359, 360, 362, 364, 365, 378, 303
Essential air-cavity, 106
Euastrum (Ehrb.), 268, 239, 241
— pectinatum (Bréb.), 241
— rostratum (Ralfs), 239
Eucheuma spinosum (Ag.), 210
Eudorina (Ehrb.), 295
Euglena, 345, 410, 449, 297
Eunotia (Ehrb.), 424, 357
— monodon (Ehrb.), 357
Eurotium (Lk.), 359, 360, 364, 367, 378, 275, 304
— herbariorum (Lk.), 366, 275, 304
— repens (de By.), 304
Eutypa (Tul.), 373
Eusynchytrium (Schroet.), 347
Exciple, 355, 372, 308
Exidia (Fr.), 389, 317
— spiculosa (Sommerf.), 317
Exoascus (Fckl.), 379
— alnitorquus (Sad.), 380
— aureus (Sad.), 380
— deformans (Fckl.), 379
— Pruni (Fckl.), 379
Exobasidium (Woron.), 389, 390
— Vaccinii (Woron.), 388
Exospore, 20, 47, 74, 80, 112, 134, 145, 160, 17, 20, 42
Extine, 166, 325, 337. 340, 347, 349, 351

FACULTATIVE parasites, 317, 454
— saprophytes, 317
False stomate, 107
Favella, 204, 181
Fegatella (Radd.), 168, 170, 171, 152
— conica (Cord.), 152
Female conceptacle, 232, 209
— filament, 266
— inflorescence, 151, 156
— prothallium, 15, 102, 103, 4, 79, 80
Ferment, 316, 380, 454
Ferns, 64, 119, 42-76, 92-95
Fertilising-tube, 202, 211, 179
Filament, 184, 428
Filamentous Lichens, 284
Filices, 15, 16, 17, 19, 20, 21, 64, 119, 42-71, 92-95
Filmy ferns, 71, 72, 88, 62-64
Fimbriaria (Nees ab Esenb.), 171
Fischera (Schw.), 439, 441
Fissidens (Hedw.), 137, 149
Fission, 185, 270, 240
Flagellate Infusoria, 410, 416, 455, 456
Float-corpuscle, 31
Floating apparatus, 31
Florideæ, 185, 188, 191, 444, 169-198
Florideæ-green, 194
'Flos-aquæ,' 436
Flower, 134, 139, 141, 154, 104, 106, 129
'Flowering fern,' 73, 90, 66, 67
'Flowers of tan,' 403
Foliaceous Lichens, 321
Foliose Hepaticæ, 135, 156, 160, 132-134, 137-142
Fontinalis (L.), 145, 149, 111
— antipyretica (L.), 111

FOO

Foot, 17, 41, 71, 105, 144, 146, 154, 422, 14, 131
Fossil Cryptogams, 114, 172, 183, 303, 330, 424, 87–101
Fossombronia (Radd.), 156, 164, 135
— pusilla (Nees ab Esenb.), 135
Fovea, 50
Foveola, 51
Fragillaria (Lyngb.), 426
Fragmentation of nucleus, 176
Frond, 72, 191, 228, 239, 241, 242, 430, 432, 359
Frondose Hepaticæ, 135, 156, 135, 136, 143–159
Front view (diatoms), 420. 353, 354
Fructification, 102, 110, 5, 6, 77, 94, 96, 100, 101
Fruit, 180
Frullania (Radd.), 164, 139, 140
— dilatata (Nees ab Esenb.), 139
— Tamarisci (Dum.), 139
Frustule, 268, 419, 423, 353–355
Frustulia saxonica (Ag.), 355
Fruticose Lichens, 321, 283
Fucaceæ, 185, 188, 190, 228, 304, 444, 205–211
Fucoideæ, 235, 237
Fucoxanthin, 230
Fucus (L.), 188, 235, 304, 207–210
— amylaceus,' 210
— furcatus (Ag.), 228
— serratus (L.), 235, 207
— vesiculosus (L.), 232, 235, 236, 207–210
Fuligo varians (Somm.), 403
Funaria (L.), 149, 103, 106, 108–110
— hygrometrica (L.), 103, 106, 108–110
Fundamental tissue, 10, 76, 39, 53, 54
Fungi, 3, 4, 305, 266–331
— cell-contents, 307
— cell-membrane, 308
— differentiation of thallus, 309
— histological characteristics, 306
— luminosity, 316
— nutrition, 316
— spores, 312
— — conditions of germination, 314
— conditions of vegetation, 316
Fungus-cellulose, 308
Funiculus, 328
Funnel, 31, 36, 15
Furrow, 100, 106, 124, 129, 420, 353

GALAXAURA (Lmx.), 211
Galls, 284, 347
Gametange, 272, 277, 279, 296, 349, 411, 246, 247, 260, 298
Gamete, 259, 335, 337, 338, 340, 346, 349, 350, 351, 293, 298, 299
Gasteromycetes, 309, 388, 395, 324–331
Gasteromycetous Lichen, 319
Gautieria (Vitt.), 395
Geaster (Mich.), 396
Gelatine, 190, 210
Gelatinous Lichens, 321
Gelidiaceæ, 209
Gelidium (Lmx.), 209, 210
Geminella (Schroet.), 350
Gemma, 61, 86, 133, 140, 157, 170, 185, 196, 208, 219, 250, 106, 114
Genabea (Tul.), 357, 358
Genea (Vitt.), 358
General bundle-sheath, 109
Genuflexion, 265
Geocalyceæ, 162
Geocalyx (Nees ab Esenb.), 164
Germ-cell, 179
Germ-filament, 20
Germ-tube, 314, 325, 326, 327, 328, 329, 333, 351, 352, 373, 385, 389, 390, 293, 299

HAL

Gigartina (Lmx.), 208, 169
— mamillosa (Ag.), 169
Gigartinaceæ, 203, 208, 169, 185
Gills, 393, 319–321
Giraudia (D. & S.), 239, 247, 212
— sphacelarioides (D. & S.), 249, 212
Girdle, 420, 353, 354
Girdle-band, 268
Girdle-view (diatoms), 420, 354
Gland, 76, 67
Glaucocystis (Itzig.), 410
Glebe, 395, 397, 398, 399, 325, 329–331
Gleichenia (Sw.), 77, 85, 60
Gleicheniaceæ, 85, 60
Glochid, 30, 13
Glochiococcus (Lagerh.), 417, 346
— anglicus (Benn.), 346
Glœocapsa (Ktz.), 417, 440, 446, 447, 448, 449, 279
— fenestralis (Ktz.), 440
Glœochæte (Lagerh.), 447
Glœocystis (Näg.), 300, 417, 419, 448
Glœothece (Näg.), 447, 376
— granosa (Rabh.), 376
Glœotrichia (Ag.), 436
— natans (Thur.), 435
— punctulata (Thur.), 436
Glomerule, 213
Glossopode, 50
Glycogen, 308, 355, 357
Gomphonema (Ag.), 424, 354, 356
— constrictum (Ehrb.), 354B, 356
Gonatonema (Wittr.), 261, 263
— notabile (Wittr.), 263
Gonatozygon (de By.), 271
Gongrosira (Ktz.), 284
— de Baryana (Rabenh.), 280
Gonid, 6, 322, 409, 410, 413, 414, 437, 446, 448, 341, 343, 344
Gonium (Müll.), 186, 299, 302
Gonoplasm, 325, 328
Gottschea appendiculata (Nees ab Esenb.), 133
Gracilaria (Grev.), 199, 201, 208, 178, 187
— compressa (Ag.), 178
— confervoides (Grev.), 209, 187
— lichenoides (L.), 210
Graphiola (Port.), 350
Grasses, 350, 364, 376, 385
Grateloupia (Ag.), 208
Griffithsia (Ag.), 199, 204
Grimmia (Ehrh.), 149
Gulf-weed, 232, 236, 211
Gum-cell, 58, 76, 123
Guttulina (Cienk.), 405, 406
— protea (Fay.), 406
Gymnoascus (Baranet.), 359, 360, 361, 367, 368, 369, 370, 378
Gymnocarpous Lichens, 356
Gymnogramme (Desv.), 16, 67, 83, 85, 58
— leptophylla (Desv.), 65
Gymnomitrium (Nees ab Esenb.), 164
Gymnopodal shoot, 176
Gymnosperms, 12, 13, 14, 15, 52, 1, 3
Gymnosporangium (DC.), 386
Gymnostomous, 135, 147
Gyrolith, 183

HADROME, 19
Hæmatococcus (Ag.), 416
— Butschlii (Blockm.), 417
Halidrys (Grev.), 232, 235, 206
— siliquosa (Lyngb.), 206
Halimeda (Lmx.), 289, 258
— Tuna (Lmx.), 258

HAL

Halosphæra (Schm.), 411, 337
— viridis (Schm.), 337
Halymenia (Ag.), 208
Hapalocystis mirabilis (Sorok.), 347
Hapalosiphon (Näg.), 439, 441, 367
— byssoideus (Kirchn.), 367
Haplomitrium (Nees ab Esenb.), 156, 161
Haplospora (Kjellm.), 249
Hard Fern, 72
Hauckia (Brzi.), 413
Haustorium, 309, 323, 326, 340, 350, 362, 378, 269, 285, 296
Hawlea Miltoni (Stur), 94
Hedwigia (Ehrh.), 149
Helicostylum (Cord.), 339
Helminthocladia (Ag.), 211
Helminthocladiaceæ, 189, 211, 175, 189–191
Helminthostachys (Kaulf.), 98, 99, 100
Helvella esculenta (Pers.), 278
Hepaticæ, 135, 156, 132–159
Heracleum sphondylium, 348
Hermonitis (L.), 85
Heterocyst, 427, 430, 431, 434, 438, 359, 362
Heterœcism, 383
Heteromerous thallus, 320
Heterophyadic Equisetaceæ, 113
Heterosporous Vascular Cryptogams, 13, 15, 20, 21, 5–33
Hibernating spores (Fungi), 315
Hibiscus, 340
Hildenbrandtia (Nard.), 190, 191, 193, 210, 211, 188
— prototypus (Nard.), 188
— rivularis (Ag.), 211
Himanthalia (Lyngb.), 228, 232, 235, 205
— lorea (Lyngb.), 205
Homoiomerous thallus, 321
Homophyadic Equisetaceæ, 113
'Honey-dew,' 376
Hookeria (Sm.), 149
Hoop, 420
Hops, 364
Hormactis (Thur.), 436
Hormidium (Ktz.), 279
Hormiscia (Aresch.), 277
Hormogone, 427, 429, 431, 435, 438, 442, 445
Hormospora (Bréb.), 414, 344
— mutabilis (Bréb.), 344
Horse-tails, 100, 113, 124
House-fly, 343, 378
Hyaline hair, 226, 275, 433, 199
Hyaloplasm, 403
Hyalotheca (Ehrb.), 187, 268
Hybridism, 145, 235
Hydneæ, 393
Hydnobolites (Tul.), 357, 358
Hydnocystis (Tul.), 358
Hydnotria (B. and Br.), 358
Hydnum (L.), 392
Hydrianum (Rabh.), 413
Hydroclathrus (Bory), 245
Hydrocytium (A. Br.), 413
Hydrodictyeæ, 277, 291, 296, 409 260
Hydrodictyon (Roth), 186, 296, 410, 413, 260
— utriculatum (Roth), 260
Hydrolapathum (Rupr.), 193, 196, 208, 71
— sanguineum (Stackh.), 171
Hydropterideæ, 21
Hydrurus (Ag.), 188, 190, 256, 257, 232
— penicillatus (Ag.), 232
Hygroscopic properties, 171
Hymenium, 355, 357, 368, 373, 376, 384, 386, 389, 391, 393, 394, 395, 396, 276, 300, 301, 305, 317, 322
Hymenogaster (Vitt.), 395

KIE

Hymenogastreæ, 395, 396, 398
Hymenomycetes, 388, 395, 266, 270, 273, 319–323
Hymenophallus (Nees ab Esenb.), 398
Hymenophyllaceæ, 16, 17, 18, 19, 64, 66, 67, 70, 71, 73, 76, 80, 81, 86, 121, 122, 61–64, 93
Hymenophyllum (L.), 86, 87, 88, 62–64
— tunbridgense (Sm.), 62
Hypertrophy, 324, 326, 327
Hypha, 228, 306, 208–210, 267, 273, 281, 282, 285, 300, 307, 308, 317, 322
Hypnea (Lmx.), 209
Hypneaceæ, 208
Hypnosperm, 225, 227, 266, 283, 295, 296
Hypnosporange, 285
Hypnospore, 262, 281, 284, 300
Hypnum (Dill.), 149
— populeum (Sw.), 113, 118
Hypochnus (Fr.), 389, 266
— centrifugus (Tul.), 266
Hypocopra (Fckl.), 361
Hypodermal tissue, 107
Hypothece, 355, 360, 361, 370, 308
Hypoxylon (Bull.), 373
Hysterineæ, 356

Ileodictyon (Tul.), 398
Impotent antherids, 332
Impregnating tube, 324, 332, 286
Inactis (Ktz.), 441, 444
Indusium, 24, 29, 50, 73, 79, 86, 121, 19, 32, 59, 63, 93
Inflorescence, 134, 158, 170, 104, 106, 109, 131, 150, 151, 156, 157
Infusoria, agents in fertilisation, 199
Innovation, 85, 133, 139, 151, 157
Intercalary growth, 241
— surface-growth, 222, 200
Internode, 102, 174, 81, 165, 168
Intine, 325, 337, 339, 349, 351
Involucre, 85, 94, 110, 159, 165, 171, 59
Involution-forms (Bacteria), 451
Iodine, 190, 236, 244
'Irish moss,' 210
Isactis (Thur.), 436, 365
— plana (Thur.), 365
Isinglass, 210
Isocystis (Brzi.), 432
Isoëteæ, 21, 47, 119, 28–33
Isoëtes (L.), 19, 38, 47, 52, 119, 28–33
— lacustris (L.), 28–33
Isoëtites (Schmp.), 119
Isogamous reproduction, 272
'Isospore,' 285
Isosporous Vascular Cryptogams, 12, 15, 20, 53, 34–85
Isthmus, 270

Japanese isinglass, 210
Jungermannia (L.), 164, 132, 138, 142
— barbata (Schreb.), 138
— bicuspidata (L.), 142
— nemorosa (L.), 132
Jungermanniaceæ, 159, 160, 172, 132–142

Kallymenia (Ag.), 208
Kaulfussia (Bl.), 78, 92, 93, 94, 95
Kelp, 190, 236, 244
Kickxella (Coem.), 341
Kieselguhr, 424

LAB

Labiatæ, 364
Labium, 51
Laboulbenia flagellata (Peyr.), 312
Laboulbenieæ, 378, 312
Lactarius (Fr.), 394
Lacuna, 105, 81
Lagenidium (Schenk), 330, 331
Lamella, 393, 394, 153, 273, 319–321
Lamina, 50, 230, 242
Laminaria (Lmx.), 239, 304, 215
— digitata (Lmx.), 244
— saccharina (Lmx.), 215
Laminariaceæ, 190, 230, 236, 239, 241, 242, 304, 213–217
Lamprothamnus (A. Br.), 182
— alopecuroides (A. Br.), 176
Lastrea (Presl), 85
Lateral conjugation, 260, 265, 235, 238
Lathyrus, 364
Laticiferous hyphæ, 394
Laudatea (Johow), 319
Laurencia (Lmx.), 209
Laver, green, 219
— purple, 217
Leaf-sheath, 102, 105, 124, 81, 83
Leathesia (Gray), 241
Leiodermaria (Ren.), 117, 89
Leiodermarieæ, 117, 89
Lejeunia (G. & L.), 164
Lejolisia (Born.), 201, 209, 177
— mediterranea (Born.), 177
Lemanea (Bory), 191, 196, 214, 192
— fluviatilis (Ag.), 216
— nodosa (Ktz.), 192
Lemaneaceæ, 189, 195, 214, 192
Lemna, 432
Lenticel. 93
Lepidodendreæ, 115, 87, 88
Lepidodendron (Sternb.), 115, 87, 88
Lepidophloios (Sternb.), 116
Lepidophyllum (Brongn.), 88
Lepidostrobus (Brongn.), 115, 116, 88
— Brownii (Schmp.), 116, 88
— dabadianus (Schmp.), 116
— ornatus (Hook.), 88
Lepidozia (Dum.), 164
Leptochæte (Brzi.), 436
Leptochrysomyxa (de By.), 386
— Abietis (Ung.), 386
Leptogium microphyllum (Ach.), 307, 308
Leptome, 19
Leptophloem, 146
Leptopuccinia Dianthi (Schroet.), 386
— Malvacearum (Schroet.), 386
Leptopuccinieæ, 386
Leptosira (Brzi.), 280
Leptothrix (Ktz.), 4, 440, 445, 451
— muralis (Ktz.), 440
— parasitica (Ktz.), 440
Leptoxylem, 146
Lessonia (Bory), 244, 214
— fuscescens (Bory), 214
Leucobryum (Hampe), 138, 139, 149
Leucochytrium (Schroet.), 347
Leucodon (Schw.), 149
Leuconostoc (Van Tiegh.), 433, 455
Liagora (Lmx.), 194, 211
Lichen-fungi, 317, 318, 361
Lichens, 222, 279, 318, 356, 419, 441, 448, 449, 279–284, 307, 308
— discocarpous, 370
— gymnocarpous, 356
Licmophora (Ag.), 426
Lid-cell, 17, 158
Ligulatæ, 38, 44

MAS

Ligule, 38, 44, 51, 31, 32
Lindsaya (Dry.), 85, 58
Lip-cell, 79, 56
Lithoderma (Aresch.), 190, 251, 225
— fatiscens (Aresch.), 225
Lithophyllum (Phil.), 206
Lithothamnion (Phil.), 206
Liverworts, 135, 156, 160
Lobospira (Thur.), 254
Lomaria (Willd.), 83, 85
— spicant (Desv.), 72
Lomentaria (Gaill.), 209
Lomentariaceæ, 209
Lophoclea (Dum.), 164
Loxsoma (R. Br.), 86, 87, 121
Lucern, 364
Luminosity of fungi, 316
Lunularia (Mich.), 157, 170, 171
Lupins, 364
Lychnothamnus (Leon.), 182
— stelliger (A. Br.), 176
Lycoperdaceæ, 395, 396, 398, 399
Lycoperdon (Tourn.), 308, 395
Lycopodiaceæ, 11, 15, 16, 18, 19, 53, 34–41
Lycopodieæ, 53, 118, 34–39
Lycopodites (Brongn.), 118
— Stockii (Kidst.), 118
Lycopodium (L.), 19, 20, 53, 56, 61, 34–37, 39
— albidum (Bak.), 59
— annotinum (L.), 54, 34, 39
— cernuum (L.), 54, 61, 35
— clavatum (L.), 56, 61, 37
— inundatum (L.), 53
— Phlegmaria (L.), 55, 118, 36
Lygodium (Sw.), 71, 77, 90, 91, 69
— palmatum (Sw.), 69
Lyngbya (Ag.), 441, 444, 445, 449, 371
— æstuarii (Liebm.), 371
— antliaria (Hansg.), 445
Lyngbyeæ, 441, 445

Macrocystis (Ag.), 239, 240, 244, 213, 217
— pyrifera (Ag.), 213, 217
'Macrosporange,' 7
'Macrospore,' 7
Macrosporium Sarcinula (Berk.), 374
'Macrozoospore,' 7
Madotheca (Dum.), 157, 164
Male conceptacle, 233, 208
'— fern,' 72
— filament, 266
inflorescence, 104, 150, 152, 157
— prothallium, 15, 103, 78
Mantle-cells, 36, 80, 99, 104
Manubrium, 175, 177, 163
Marattia (Sm.), 91, 94, 95, 70, 71
— cicutæfolia (Kaulf.), 93, 95
Marattiaceæ, 20, 21, 64, 65, 74, 77, 78, 79, 81, 91, 121, 122, 70, 71, 94
Marchantia (L.), 157, 170, 171, 149–151, 153–159
— polymorpha (L.), 157, 149–151, 153–159
Marchantiaceæ, 133, 157, 160, 167, 149–159
Marchesettia spongioides (Hauck), 210
Marsilea (L.), 33, 37, 114, 4, 6, 14, 15, 17, 19
— Drummondii (R. Br.), 38
— quadrifolia (L.), 34, 6
— salvatrix (L.), 14, 15, 17, 19
Marsileaceæ, 12, 13, 25, 31, 4–6, 14–19
Marsilidium (Schenk), 114
Martensella (Coem.), 341
Massaria (De Not.), 356
Massula, 25, 30, 13
Mastigobryum (Nees ab Esenb.), 164

MAS

Mastigocladus (Cohn), 439, 441
Mastigocoleus (Lagerh.), 439, 440, 441
Mastogloia (Thw.), 424, 426
Mazæa (Born.), 440
Mechanical system, 138, 171
'Medulla,' 118, 124, 280
Medullary system, 76, 228, 242, 249, 321, 271
Megasporange, 7, 11, 22, 277, 7, 18, 19, 26, 27
Megaspore, 7, 11, 15, 20, 22, 118, 155, 7–9, 14–16, 18, 20, 29, 125
Megazoosporange, 260
Megazoospore, 7, 218, 273, 296, 410, 196, 260
Melampsora populina (Jacq.), 386
Melanospermeæ, 235, 237
Melanospora (Cord.), 359, 360, 370
Melobesia (Lmx.), 193, 194, 196, 206, 182, 184
— membranacea (Lmx.), 182
— Thureti (Born.), 184
Melobesiaceæ, 191
Melosira (Ag.), 424
Meridion (Leibl.), 426, 357
— constrictum (Ralfs), 357
Merismopedia (Mey.), 418, 446, 447, 448
Merispore, 6
Mertensia (Willd.), 77, 85
Merulius lacrymans (Fr.), 309
Mesocarpaceæ, 187, 258, 259, 260, 233–235
Mesocarpus (Hass.), 260, 262, 263, 330, 233, 235
— neaumensis (Benn.), 261
— parvulus (Hass.), 233
— pleurocarpus (de By.), 235
Mesoglœaceæ, 239, 247, 220
Metzgeria (Cord.), 164
Meum athamanticum, 348
Micrandres, 225, 201, 202
Micrasterias (Ag.), 268, 239
— rotata (Grev.), 239
Microchæte (Thur.), 436
— diplosiphon (Gom.), 434
Micrococcus (Cohn), 450, 379
Microcoleus (Desm.), 429, 441, 444
Microcyst, 404
Microcystis (Ktz.), 446, 447, 449, 377
— marginata (Meneg.), 377
Microspora (Thur.), 276, 242
— floccosa (Thur.), 242
Microsporange, 7, 11, 22, 277, 7, 11, 18, 19, 26, 32, 33
Microspore, 7, 11, 12, 20, 22, 40, 115, 116, 155, 7, 11, 18, 20, 30, 125
Microthamnion (Näg.), 280
Microzoospore, 218, 277, 410
Mildew of corn, 383
Mischococcus (Näg.), 412, 339
— confervicola (Näg.), 339
Mitremyces (Nees ab Esenb.), 399
Mnium (L.), 138, 140, 149
Mohria (Sw.), 90, 91
Monadopsis (Klein), 405
Monas amyli (Cienk.), 405
Monoblepharideæ, 331, 290
Monoblepharis (Cornu), 4, 331, 290
— sphærica (Cornu), 290
Monoclea (Hook.), 164, 143
— Forsteri (Hook.), 143
Monocleaceæ, 164, 143
Monosiphonous, 192, 195
Monospora (Sol.), 196
Monostroma (Thur.), 217, 218, 219, 198
— bullosum (Thur.), 198
Moonwort, 100
Morchella (Dill.), 356
Mortierella (Coem.), 338, 339
— nigrescens (Van Tiegh.), 338
— Rostafinskii (Bref.), 338

NOS

Mosses, 135, 145
Mougeotia (de By.), 262, 264, 267, 330
— calcarea (Wittr.), 262
Moulds, 366
Mucilage, 69, 220, 221, 238, 256, 257, 263, 269, 283, 298, 398, 248, 330
Mucilage-cell, 58, 76, 89, 92, 168, 67, 154
Mucilaginous sheath, 259, 267, 268, 274, 298, 428, 430, 434, 437, 440, 441
Mucor (Mich.), 4, 307, 339, 340, 350, 381, 293, 296
— Mucedo (L.), 293, 296
— racemosus (Fres.), 339
— tenuis (Lk.), 338
Mucoreæ, 337, 339, 340
Muccrini, 4, 308, 315, 316, 335, 341, 342, 344, 345, 377, 378, 293–296
Multilocular zoosporange, 187, 237, 212, 227
Multinucleatæ, 186, 280, 248–258
Musci, 135, 136, 102–131
Muscineæ, 2, 132, 102–159
Mushroom, 311, 320
Mutinus caninus (Fr.), 308, 329
Mycele, 309, 337, 363, 366, 369, 375, 389, 275, 293, 296, 303–305, 318, 330
Mycetozoa, 305, 401, 456, 332–335
— doubtful, 405
Mycoidea (Cunn.), 222
— parasitica (Cunn.), 222
Mycorhiza, 310
Mylitta (Fr.), 309
Myrionema (Grev.), 241
Myxomycetes, 401, 406, 332–335
Myzocytium (Schenk), 330

NARDOO, 38
Navicula (Bory), 420, 422, 424, 357
— rhomboides (Ehrb.), 357
Neck, 16, 26, 39, 86, 68, 143, 179, 109, 148, 158
Neck-canal-cell, 16, 27, 39, 69, 159, 10
Neck-cell, 26, 68, 10
Neckera (Hedw.), 149
Nectria cinnabarina (Fr.), 287
Nemalieæ, 192, 195, 199, 211, 175
Nemalion (Ag.), 211, 213, 175
— multifidum (Ag.), 175
Nemastoma (Ag.), 208
Nemathece, 196, 199, 202
Nematodonteæ, 148
Nematophycus (Carruth.), 304
Neomeris (Lmx.), 287, 254
— Kelleri (Cram.), 287, 254
Nephrocytium (Näg.), 414, 343
— Nägelii (Grün), 343
Nephrodium (Rich.), 85
Nephrolepis (Sch.), 77, 85
Nereocystis (Post.), 240, 242, 244
Neutral zone, 175, 163
Nidularieæ, 397, 327, 328
Nitella (Ag.), 175, 176, 179, 182, 163, 165, 166
— flexilis (Ag.), 163, 165, 166
Nitelleæ, 173, 182, 163, 165, 166
Nitophyllum (Grev.), 196, 208, 173
— punctatum (Harv.), 173
Nitzschia (Hass.), 426
Node, 102, 173, 421, 165, 168
Nodularia (Mert.), 430, 433
Nodule, 421
Nostoc (Vauch.), 28, 165, 171, 429, 430, 431, 432, 433, 358, 359
— commune (L.), 358
— hyalinum (Benn.), 359
— muscorum (Ag.), 431
— parietinum (Rabenh.), 440

NOS

Nostocaceæ, 185, 427, 428, 430, 455, 456, 358–362
Nostochineæ, 186, 427, 428, 358–373
Nothoclæna (R. Br.), 85
Notommata, 284
Nuclei, plurality of, 186, 187, 188, 194, 272, 275, 281, 284
Nuclearia (Cienk.), 405
Nullipore, 206
Nummularia (Tul.), 373

Obelidium (Nowak.), 346
Octaviana asterosperma (Vitt.), 324
'Octospore,' 217
Œdogoniaceæ, 188, 220, 222, 200–202
Œdogonium (Lk.), 222, 223, 226, 200, 201
— ciliatum (Hass.), 201
— gemelliparum (Hass.), 201
Oidium Tuckeri (Berk.), 364
Olfersia (Radd.), 80
Oligocarpia Lindsæoides (Stur), 94
Olpidieæ, 344, 345, 346, 347
Olpidiopsis fusiformis (Cornu), 346
— Saprolegniæ (Fisch.), 346
Olpidium (A. Br.), 347
Onions, 327
Onoclea (L.), 80, 85
Onygena (Pers.), 358, 359
Ooblastema-filament, 202, 211, 215
Oogamous reproduction, 188
Oogone, 185 (see also under Algæ and Fungi), 199, 201, 202, 210, 228, 248, 249, 259, 286, 290, 291
Oomycetes, 323, 332, 285–292
Oophyte, 10, 16, 132, 135
Oosperm, 11 (see also under Vasc. Crypt., Musc., Char., Algæ, and Fungi), 203, 210, 229, 248, 259, 286, 290, 291
Oosphere, 8 (see also under Vasc. Crypt., Musc., Char., Algæ, and Fungi), 4, 10, 15, 45, 109, 148, 158, 201, 203, 210, 229, 259, 286, 291
'Oosporange,' 237
'Oospore,' 8
'Oosporeæ,' 3
Opercule, 135, 112, 117, 131
Ophiocytium (Näg.), 414
Ophioglossaceæ, 13, 16, 20, 21, 81, 95, 123, 72–76
Ophioglossum (L.), 97, 98, 99, 100, 72, 76
— pedunculosum (Desv.), 96
— vulgatum (L.), 98, 100, 72, 76
Orthotrichum (Hedw.), 149
Oscillaria (Bosc.), 429, 441, 442, 443, 445, 368, 369
— ærugineo-cœrulea (Ktz.), 443
— antliaria (Hansg.), 445
— dubia (Ktz.), 443
— Frölichii (Ktz.), 446
— princeps (Vauch.), 441, 443, 444, 369
— tenuis (Ag.), 446, 368
Oscillariaceæ, 427, 428, 429, 433, 437, 441, 455, 368–373
— movements of, 442
Oscillatorieæ, 441
Osmunda (L.), 66, 67, 75, 89, 90, 65–68
— cinnamomea (L.), 90
— regalis (L.), 73, 90, 65–67
Osmundaceæ, 66, 75, 76, 78, 80, 81, 88, 65–68
Osmundites (Carruth.), 123
Ostiole, 207, 356, 182–184, 208, 209, 311
Ovulites (Lam.), 304
Oxymitra (Bisch.), 166

Pachyma (Fr.), 309
Pachyphlœus (Tul.), 358

PER

Padina (Adans.), 254, 230, 231
— Pavonia (Gaill.), 230, 231
Palæopteris hibernica (Schimp.), 121, 93
Palæostachys (Weiss), 128
Pale, 72
Palisade-cells, 321
Palisade-parenchyme, 19
Palmella (Lyngb.), 415, 417, 448
— cruenta (Ag.), 416
— nivalis (Hook.), 416
— prodigiosa (Mont.), 416
— uvæformis (Ktz.), 276, 417
Palmellaceæ, 257, 415, 418, 419
Palmellin, 416
Palmelloid condition, 277, 409, 411, 412, 415, 246, 345
Palmodictyon (Ktz.), 418
Pandorina (Ehrb.), 186, 299, 300, 263
— morum (Ehrb.), 263
Pandorineæ, 186, 291, 299, 409, 263
Paraphyse, 55, 79, 87, 134, 142, 196, 233, 237, 244, 354, 355, 356, 358, 370, 371, 372, 373, 374, 386, 388, 394, 36, 55, 106, 208, 209, 276, 300, 305
Parasites (Fungi), 316, 317, 329, 383
— (Bacteria), 454
Parasitic Algæ, 249, 280, 284
Parsley-fern, 72
Parthenogenesis, 181, 267, 333, 338
Parthenosperm, 261, 262, 267
Parthenospore, 292
Pecopteris arborescens (Schl.), 124
Pediastreæ, 186, 190, 291, 298, 418, 261, 262
Pediastrum (Mey.), 298, 261, 262
— integrum (Näg.), 261
Pedicel, 80
Pedicel-cell, 177, 179, 233, 369
Pellæa (Lk.), 85
Pellia (Radd.), 162, 164, 136
— epiphylla (Cord.), 136
Peltate leaf, 28, 8, 9
— scale, 102, 110, 2, 83
Pelvetia (Dcne.), 235
Penicillium (Lk.), 312, 358, 359, 360, 366, 378
— glaucum (Lk.), 316, 367
Penicillus (Ktz.), 289
Penium (Bréb.), 268, 241
— margaritaceum (Bréb.), 241
Perianth, 134, 141, 142
Pericarp, 180, 201, 221, 263, 199
Pericentral tubes, 192
Perichæte, 134, 141, 162, 109, 129, 131
Periderm, 430
Peridinieæ, 410
Peridiolum, 397, 328
Peridium, 311, 357, 358, 395, 396, 397, 398, 399, 325, 328–331
Perigone, 142, 106
Perigyne, 159, 163, 171, 142, 158
Perinium, 160
Peripheral growth, 241
Periphyse, 356
Periplasm, 324, 325, 327, 331, 332, 286
Peristome, 135, 147, 110–112, 114, 118
Perithece, 355, 356, 362, 365, 370, 373, 374, 375, 378, 379, 311, 312
Peronospora (Cord.), 4, 323, 326, 327, 328, 329, 378
— arborescens (de By.), 286
— calotheca (de By.), 269, 285
— densa (Rab.), 327
— Ficariæ (Tul.), 327
— nivea (de By.), 327
— parasitica (de By.), 326, 327
— pygmæa (Ung.), 327

PER

Peronospora Rumicis (Cord.), 327
— Schachtii (Fckl.), 327
— Schleideniana (Ung.), 327
— Trifoliorum (de By.), 327
— Viciæ (de By.), 327
— viticola (de By.), 327
Peronosporeæ, 4, 309, 312, 315, 323, 331, 332, 333, 377, 378, 269, 272, 275, 277, 285–289
Peronosporites (W. G. S.), 330
— antiquarius (W. G. S.), 330
Persistent cell, 444
Petalonema (Berk.), 438, 440
Peyssonelia (Dcne.), 210
Peziza (Dill.), 356
— (Pyronema) confluens (Pers.), 369, 276, 300, 306
Phacidiaceæ, 356
Phæophyll, 230
Phæosporeæ, 187, 190, 230, 237, 212–232
Phæothamnieæ, 258
Phæothamnion (Lagerh.), 258, 276
Phæozoosporeæ, 237
Phalloideæ, 395, 397, 329–331
Phallus (L.), 398, 399, 330
— impudicus (L.), 398, 330
Phascaceæ, 136, 150, 121, 122
Phascum (L.), 140, 150
Phloem-sheath, 18, 59
Phlyctidium (A. Br.), 346
Phragma, 125
Phragmidium (Lk.), 386, 314
— incrassatum (Lk.), 314
Phycochromaceæ, 426
Phycocyanin, 427, 442
Phyco-erythrin, 194, 217
Phycomyces (Kze.), 338, 339, 293
— nitens (Kze.), 293
Phycomycetes, 4, 5, 323, 352, 377, 381, 285–299
Phycophæin, 230, 240, 256
Phycoseris (Ktz.), 217
Phycoxanthin, 230, 421
Phyllade, 51
Phyllitis (Ktz.), 245
Phyllobium (Klebs), 284, 410
Phylloglossum (Kze.), 19, 53, 56, 59, 60, 61, 38
— Drummondii (Kze.), 61, 38
Phyllosiphon (Kühn), 284
Phyllotaxis, 59, 99, 106, 139, 151, 194
Phyllotheca (Brongn.), 124, 96
— equisetiformis, 96
Physareæ, 403
Physcia parietina (Nyl.), 279
Physiotium (Nees ab Esenb.), 161
Physma (Massal.), 372
Physoderma (Wallr.), 346
Phytophthora (de By.), 323, 327, 328, 332, 335, 272, 277, 289
—infestans (de By.), 314, 326, 327, 328, 272, 277, 289
— omnivora (de By.), 327, 329
Pigment-spot, 223, 292, 300, 416
Pileus, 286, 391, 392, 393, 398, 399, 251, 252, 273, 319, 320, 330
Pilinia (Ktz.), 280
Pilobolus (Tode), 339
Pilularia (L.), 19, 33, 37, 114, 5, 16, 18
— globulifera (L.), 36, 5, 16, 18
Pinnularia (Ehrb.), 426, 353
— viridis (W. Sm.), 353
Piptocephalideæ, 340, 378, 295, 296
Piptocephalis (de By.), 340, 342, 295, 296
— Freseniana (de By. and Wor.), 295, 296
Pithophora Kewensis (Wittr.), 276, 245
Pithophoraceæ, 187, 273, 276, 245
Pits in oogonial wall, 332

POL

Placenta, 25, 29, 35, 80, 86, 201
Placental cells, 209, 287
Plagiochasma (L. & L.), 171
Plagiochila (Dum.), 164
Plantago, 364
Plasmatoparous Peronosporeæ, 327
Plasmode, 401, 402, 403, 404, 405, 406, 332, 333
— movements of, 403
— resting states of, 404
Plasmodiophora (Woron.), 405
— Brassicæ (Woron.), 405
Platycerium (L.), 83, 85
— alcicorne (Desv.), 72, 77
Plaxonema (Tangl), 444
Plectonema (Thur.), 437, 438, 439
Pleomorphy of Uredineæ, 383
Pleospora (Rabenh.), 361, 362, 374
— herbarum (Rabenh.), 374
Plerome-sheath, 18, 109
Pleuridium (Brid.), 150, 122
— subulatum (Rabenh.), 122
Pleurocarpi, 149
Pleurocladia (A. Br.), 247
Pleurococcus, 279, 300, 417, 419
Pleurosigma (W. Sm.), 420, 426, 357
— lacustre (W. Sm.), 357
Plocamium (Lmx.), 196, 208
— coccineum (Huds.), 209
Plumule, 41
Podosphæra (Kze.), 360, 362, 378, 303
— Castagnei (de By. and Wor.), 364, 303
— Kunzei (Lév.), 364
— pannosa (de By. and Wor.), 303
Pollen-grain, 11, 12, 14, 3
Pollen-sac, 13, 1
Pollen-tube, 11, 14, 3
Pollexfenia (Harv.), 194, 209
Pollinoid, 7, 185, 198, 199, 207, 254, 360, 361, 370, 371, 372, 373, 374, 384, 385, 175, 179, 182, 184, 191, 307
Polyedriaceæ, 419
Polyedrium (Näg.), 299, 418, 419
Polygonum, 364
— Hydropiper, 351
Polyhedra, 296, 298, 299, 262
Polyides (Ag.), 202, 209
Polyphagus (Nowak.), 314, 346, 297
— Euglenæ (Nowak.), 345, 297
Polyphysa (Lmx.), 288, 256
— peniculus (R. Br.), 256
Polypodiaceæ, 64, 67, 75, 78, 79, 80, 81, 83, 121, 123, 42, 44–56, 58
Polypodieæ, 83, 84
Polypodium (L.), 77, 84, 85, 53, 58
— leiorhizum (Wall.), 53
Polyporeæ, 393, 396, 322, 323
Polyporus (Mich.), 392, 396, 322, 323
— annosus (Fr.), 316
— igniarius (Fr.), 322, 323
— obvallatus (Berk. and Cooke), 396
— sulphureus (Fr.), 316
— volvatus (Pk.), 396
Polysaccum (DC.), 399
Polysiphonia (Grev.), 192, 193, 194, 201, 209, 170
— opaca (Zan.), 170
Polysiphonous, 192
Polyspore, 6
Polystichum angulare (Willd.), 69, 48
Polystigma (Pers.), 360, 372, 385
Polythrix (Zanard.), 437
Polytrichaceæ, 146
Polytrichum (L.), 138, 139, 142, 148, 149, 104, 107, 112, 117
— commune (L.), 104, 107, 117

POL

Polytrichum piliferum (Schreb.), 112
Polyzonia (Suhr), 194
Pomeæ, 386
Pore, 154, 187
Pores of Polyporeæ, 393, 322
Porphyra (Ag.), 189, 191, 199, 216, 219
— vulgaris (L.), 217
Porphyraceæ, 189, 190, 196, 198, 199, 216, 219, 193–195
Porphyridium cruentum (Näg.), 416
Potato-disease, 328
Pottia (Ehrh.), 147, 149
Prasiola (Ag.), 217, 219
Preissia (Cord.), 169, 170, 171
Primary node, 180
— root, 180, 168
Procarp, 198, 199, 213, 176, 187, 192
'Proembryo,' 177, 180
'Proembryonic branch,' 176
Progametange, 349, 298
Prolific cells, 276
Prolification, 69, 139, 142, 196, 289, 104, 171
Promycele, 325, 328, 329, 337, 338, 350, 351, 352, 362, 373, 385, 386, 391, 299, 314
Propagation, 8
Propagule, 196, 237, 250, 289, 223, 224
Prothallium, 10, 339 (see also under Vasc. Crypt., Musc., & Char.), 4, 8, 9, 14, 15, 20, 29, 34–36, 42, 43, 50, 65, 74, 78–80, 126, 168
Prothalloid branch, 176
— growth, 70, 48
Protococcaceæ, 186, 345, 409, 410, 415, 448, 345–352
Protococcoideæ, 186, 408, 409, 449, 336–352
Protococcus (Ag.), 186, 285, 409, 417, 448, 279, 345
— pluvialis (Ktz.), 345
Protomyces (Ung.), 4, 350, 352, 298
— macrosporus (Ung.), 348, 298
Protomycetaceæ, 348, 298
Protomyxa (Haeck.), 405
Protoneme, 133, 135, 136, 140, 214, 449, 105, 121, 125
Protophloem, 59
Protophyta, 2, 3, 4, 186, 276, 407, 336–382
Protosalvinia (Daws.), 115
Protozoa, 456
Prozoosporange, 346, 297
Prunus, 364, 373
Psaronieæ, 124
Psaronius (Cord.), 124, 95
Pseudo-bulbil, 70
Pseudo-cortex, 192, 212
Pseudocyst, 410, 427, 437, 441, 446
Pseudo-parenchyme, 249, 251, 307, 356, 358, 360, 366, 372, 384, 267
Pseudopode, 151, 154, 402, 403, 422, 443, 131, 332
Pseudo-ramulus, 434, 438, 364, 366
Pseudospora (Cienk.), 405
Psilophyton (Daws.), 119
Psiloteæ, 21, 53, 61, 119, 40, 41
Psilotum (Sw.), 17, 18, 19, 20, 53, 61, 63, 40
— triquetrum (Sw.), 61, 40
Pteris (L.), 85, 46, 47, 54, 58
— aquilina (L.), 75, 76, 77, 82, 54
— serrulata (L. fil.), 69, 46, 47
Ptilophyton (Daws.), 118
Ptilota (Ag.), 194, 204
Puccinia (Pers.), 383, 385, 386
— coronata (Cord.), 314
— graminis (Pers.), 383, 274, 314, 315
— straminis (Fckl.), 314
Puff-ball, 311, 396
Pulvinus, 93

RHI

Punctaria (Grev.), 187, 241, 245
Punctariaceæ, 239, 245
Pycnid, 362, 374
Pycnochytrium (de By.), 347
Pycnophycus (Ktz.), 235
Pycnospore, 362, 374, 375
Pyrenocarp, 355
Pyrenomycetes, 319, 355, 356, 370
Pyronema (Fckl.), 356, 359, 360, 361, 369, 370, 372, 378, 276, 306
— confluens (Tul.), 276, 306
Pythium (Pringsh.), 317, 324, 329, 330, 331, 332, 335, 286
— Chlorococci (Lohde), 329
— circumdans (Lohde), 329
— de Baryanum (Hesse), 329
— entophytum (Pringsh.), 329
— Equiseti (Sad.), 329
— gracile (Schenk), 329, 286
— intermedium (de By.), 329
— proliferum (de By.), 329
— vexans (de By.), 325, 329, 377

QUATERNARIA (Tul.), 373

RADIOLARIANS, yellow cells of, 318
Radula (Dum.), 162, 164, 137, 141
— complanata (Dum.), 137, 141
Ralfsia (Berk.), 241, 251
Ralfsiaceæ, 239, 251, 225
Ramentum, 72
Ranunculus, 364
Raphe, 421, 353
Raphidium (Ktz.), 418, 351
— falcatum (Ktz.), 351
Reboulia (Radd.), 171
Receptacle, 80, 170, 232, 246, 149–151, 154, 156, 157, 205, 207, 219
Receptive spot, 69, 227
'Red snow,' 416
Rejuvenescence, 223, 274, 281
Renaultia (Stur), 122
Reproduction, 8
Reserve-system, 139
Resting-cell, 213, 264, 335, 339, 346, 347
Resting-sporange, 333
Resting-spore, 223, 258, 276, 278, 281, 285, 298, 315, 344, 345, 350, 351, 352, 416, 427, 430, 436, 261, 361, 362
Resting-swarm-cell, 274
Resting-zoosporange, 346
Retrogression, 4, 407, 449
Rhabdonema (Ktz.), 423, 424
Rhacophyllum adnascens (L. & H.), 120, 92
Rhipidonema (Mattir.), 319
Rhizidieæ, 344, 345
Rhizidium (A. Br.), 346
Rhizocarpeæ, 18, 19, 21, 114, 4–19
Rhizoclonium (Ktz.), 276
Rhizoglossum (Presl), 99
Rhizoid, 16, 132, 139, 156, 174, 180, 222, 228, 239, 241, 242, 273, 277, 282, 285, 332, 340, 345, 346, 363, 14, 43, 50, 65, 102, 153, 168, 250, 252, 256, 280, 318
Rhizome, 51, 72, 81, 86
Rhizomorph, 392, 270, 319
Rhizomorpha (Roth), 309
— fragilis (Roth), 319
Rhizophore, 45
Rhizophydium (Schenk), 346
Rhizopoda, 456
Rhizopus (Ehrb.), 339, 294
— nigricans (Ehrb.), 337, 294

RHO

Rhodomela (Ag.), 209
Rhodomelaceæ, 209, 170, 174
Rhodophyll, 194
Rhodospermeæ, 191
Rhodospermin, 194
Rhodosporeæ, 191
Rhodymenia (Grev.), 208, 186
— binda (Ktz.), 196
— palmata (Grev.), 209
— Palmetta (Grev.), 186
Rhodymeniaceæ, 208, 171, 186
Rhytidolepidæ, 117
Riccia (L.), 165, 166, 148
— glauca (L.), 148
Ricciaceæ, 160, 165, 146–148
Ridge, 100, 106, 124, 129, 81
Riella (Mont.), 156, 159, 165, 166, 146
— helicophylla (Mont.), 146
Rivularia (Roth), 436, 363
— fluitans (Cohn), 436
— polyactis (Hauck), 363
Rivulariaceæ, 185, 427, 428, 433, 445, 363–365
Roccella tinctoria (DC.), 284
Roestelia (Reb.), 386
Root-cap, 18, 78, 109
Root-hair, 18, 78
Rotation of protoplasm, 175, 163
'Royal Fern,' 73, 90, 66, 67
Rozella (Cornu), 345, 347
Rumex, 364
Russula adusta (Fr.), 308
Rye, 310
Rytiphlœa (Ag.), 194, 209

SOR

Saccharomyces (Meyen), 4, 380, 381, 268, 313
— albicans (Reess), 380
— cerevisiæ (Meyen), 380, 268, 313
— ellipsoideus (Reess), 380
— Mycoderma (Reess), 380
— Pastorianus (Reess), 380
Sacconema (Brzi.), 437
Sacheria (Sir.), 214
Saddle, 51
Salicaceæ, 310
Salmon-disease, 332
Salvinia (L.), 17, 18, 25, 31, 7–11
— natans (L.), 7–11
Salviniaceæ, 12, 13, 20, 25, 114, 7–13
Saprolegnia (Nees ab Esenb.), 4, 332, 333, 334, 335, 338, 339, 346, 347
Saprolegnieæ, 4, 308, 312, 324, 332, 338, 347, 377, 291, 292
Saprophytes, 315, 316, 317, 329, 366, 454
Sargasso Sea, 232
Sargassum (Ag.), 230, 232, 235, 211
— bacciferum (Ag.), 232, 236, 211
Scalariform conjugation, 260, 265, 233, 234, 236, 237
— tracheide, 18, 76, 98, 123
Scenedesmus (Mey.), 303, 418, 352
— obtusus (Mey.), 352
Schizæa (Sm.), 90, 91
Schizæaceæ, 64, 80, 81, 90, 122, 43, 69
Schizochlamys (A. Br.), 417, 348
— gelatinosa (A. Br.), 348
Schizogonium (Ktz.), 279
Schizomycetes, 433, 449, 379–382
Schizoneura (Schimp.), 124
Schizophyceæ, 408, 336–378
Sciadiaceæ, 410, 411
Sciadium (A. Br.), 411, 336
— arbuscula (A. Br.), 336
Sclerenchyme, 75, 123, 53, 54
Scleroderma (Pers.), 397
Sclerosis, 58, 309, 39
Sclerote, 309, 310, 361, 373, 374, 376, 392, 404, 271, 309, 310, 311, 318
Sclerotinia (Fckl.), 373
— Fückeliana (de By. and Wor.), 361, 374
— sclerotiorum (de By.), 360, 361, 373, 374, 271
Sclerotium (Tode), 310
Scolecopteris (Stur), 122, 94
— polymorpha (Stur), 94
Scolopendrium (Sm.), 81, 85, 58
Scutiform leaf, 28, 8, 9
Scyamina (Van Tiegh.), 295
Scytonema (Ag.), 438, 441, 445, 279
Scytonemaceæ, 303, 427, 428, 433, 437, 366, 367
Scytonemeæ, 438, 441, 445, 366, 367
Scytonemin, 427, 437
Scytosiphon (Ag.), 239, 241, 245
— lomentarium (Ag.), 246
Scytosiphonaceæ, 245
Seaweeds, 184, 190, 191, 235, 237
Sebacina (Tul.), 389
Secondary capitulum, 178, 163
— embryo-sac, 14, 52, 3
— growth in thickness, 49, 116, 125
— markings, 420, 353
— prothallium, 39
Secotium (Kze.), 395
— erythrocephalum (Tul.), 396
Secreting system, 139
Seftenbergia (Cord.), 122, 94
— ophidermatica (Stur), 94
Seirospora Griffithsiana (Harv.), 204
Seirospore, 196, 204, 180
Selaginella (Spring), 19, 20, 38, 39, 47, 20–27
— caulescens (Spr.), 20
— denticulata (Lk.), 24
— inæqualifolia (Spr.), 22, 23, 25–27
— Martensii (Spr.), 20, 21
Selaginellaceæ, 12, 13, 18, 19, 38, 115, 130, 20–33, 87–91
Selaginelleæ, 38, 39, 20–27
Selenastrum (Reinsch), 186, 303
Seta, 134, 144, 146, 110, 112
Sewage-fungus, 454
Sheath, 50, 428, 430, 434, 437
Shepherd's Purse, 326
Shield, 177, 163
Side-view (of diatoms), 421, 354
Sieve-hypha, 244
Sieve-plate, 240, 244, 217
Sieve-tube, 18, 58, 240, 244, 217
Sigillaria, 117, 118, 89, 91
Sigillariostrobus, 117, 118
Silica, 102, 107, 419, 420
Siphon, 192
Siphoneæ, 186, 280, 281, 290, 308, 410, 248, 249
Siphonocladaceæ, 186, 190, 279, 280, 281, 288, 304, 411, 256–258
Siphonocladus (Schr.), 289
Sirogonium (Ktz.), 264, 265, 267
Sirosiphon (Ktz.), 439, 441, 449
Sirosiphoneæ, 437, 438, 439, 441
Solenites (L. & H.), 119
Sorastreæ, 186, 291, 302, 414, 418, 264, 265
Sorastrum (Ktz.), 186, 302, 264
— spinulosum (Näg.), 264
Sordaria (Ces. and De Not.), 354, 359, 360, 370
Sorede, 319, 282
Sorophore, 37, 19
Sorosporium (Rud.), 350, 352
— Saponariæ (Rud.), 351

SOR

Sorus, 11, 20, 24, 72, 79, 94, 122, 196, 237, 244, 251, 347, 19, 58, 59, 71, 218
Spatoglossum (Ktz.), 254
Special bundle-sheath, 109
'Sperm,' 7, 8
'Sperm-cell,' 8
'Spermatia,' 7, 360
'Spermatozoid,' 8
Spermocarp, 180, 221, 165, 167, 168, 199
'Spermogone,' 7, 360
Spermothamnion (Aresch.), 209, 176
— hermaphroditum (Näg.), 176
Sphacelaria (Lyngb.), 249, 223, 224
— cirrhosa (Ag.), 223, 224
Sphacelariaceæ, 237, 239, 241, 249, 223, 224
Sphacele, 249, 223
Sphacelia (Lév.), 376, 309
Sphacelotheca (de By.), 352
— Hydropiperis (de By.), 350
Sphærobolus (Tode), 399
Sphærocarpus (Mich.), 160, 166, 147
— terrestris (Sm.), 147
Sphærococcaceæ, 208, 178, 187
Sphærococcus (Stackh.), 208
Sphærogonium (Rostaf.), 444
Sphæroplea (Ag.), 227, 203, 204
— annulina (Ag.), 226, 203, 204
Sphæropleaceæ, 188, 220, 226, 203, 204
'Sphærosp re,' 195
Sphærozyga (Ag.), 430, 433
Sphagnaceæ, 132, 136, 139, 142, 144, 145, 151 125–131
Sphagnum (L.), 138, 156, 172, 125–131
— acutifolium (Ehrh.), 125–127, 129–131
— cymbifolium (Dill), 128
— squarrosum (Pers.), 131
Sphenoglossum (Emm.), 114
Sphenophylleæ, 129, 130, 101
Sphenophyllum (Brongn.), 129, 130, 101
Sphenopteris crenata (L. & H.), 120, 92
Spherocrystal, 93
Sphyridium (Flot.), 361
Spinellus fusiger (Van Tiegh.), 338
Spiral bands, 152
Spirillum (Ehrb.), 450, 451, 382
Spirochæta (Ehrb.), 451
Spirogyra (Lk.), 4, 264, 265, 266, 267, 330, 342, 236, 238
— bellis (Hass.), 238
— crassa (Ktz.), 264
— porticalis (Vauch.), 236
Spirulina (Lk.), 429, 441, 370
— tenuissima (Ktz.), 370
Splachnidium (Grev.), 235, 236
Splachnum (B. & S.), 149
— ampullaceum (L.), 116
Splenic fever, 452, 454
Spongiocarpeæ, 209
Spongocladia (Aresch.), 276, 290
Spongodieæ, 289
Spontaneous generation, 450
movement, 415, 427, 431, 442
Sporange, 6 (see also under Vasc. Crypt., Musc., Algæ, Fungi, Mycet., and Prot.), 2, 37, 40, 41, 55, 56, 58–60, 63, 68, 71, 72, 75, 76, 83, 93, 110, 111, 114–118, 122, 124, 131, 133, 134, 137, 145, 158, 293, 334
— dehiscence of, 79, 94, 135, 155, 159, 55
Sporangiole, 339
Sporangiophore, 328, 337, 339, 287
Sporangiospore, 6
Sporangites (Daws.), 115
Spore, 5 (see also under Vasc. Crypt., Musc., Algæ, Fungi, Mycet., and Prot.), 37, 84, 85, 158, 181, 245, 249, 272, 275, 303, 317, 318, 332, 335, 361, 362, 379–381

STR

Spore-sac, 147, 112
Sporid, 6, 314, 350, 351, 352, 362, 373, 385, 386 390, 299, 314
Sporiferous filaments, 192
Sporocarp, 11, 353 (see also under Vasc. Crypt. Algæ, and Sporocarpeæ), 5–7, 12, 18, 19 302–305
— origin of, 359
Sporocarpeæ, 353, 300–331
Sporocarpon (Williams.), 114
Sporochnaceæ, 245
Sporochnus (Ag.), 241, 246, 219
— pedunculatus (Ag.), 219
Sporodinia (Lk.), 338, 339
— grandis (Lk.), 337
Sporogenous tissue, 20, 46, 33
Sporogone, 134, 136, 144, 159, 163, 107, 110, 131, 142, 143, 145, 148, 156, 158
Sporophore, 309, 325, 327, 328, 336, 337, 340, 343, 363, 365, 367, 374, 388, 389, 404, 272, 275, 288, 293, 296, 303, 304, 327–329, 330
Sporophore (compound), 391, 392, 393, 395, 396, 397, 398, 273, 319, 320
Sporophyll, 46, 51, 72, 99, 110, 116, 32, 72, 75, 76, 83, 88
Sporophyte, 10, 17, 132, 135, 159
Sporophytic budding, 69
Sprouting, 307, 314, 339, 352, 380, 381, 389, 268, 313
Spyridia (Harv.), 193, 209
Spyridiaceæ, 209
Squamariaceæ, 189, 190, 191, 202, 210, 188
Stag's-horn-moss, 61
Staurastrum (Mey.), 269, 270, 239, 240
— Arachne (Ralfs), 239
— teliferum (Ralfs), 240
Stauroneis (Ehrb.), 426
Staurospermum (Ktz.), 260, 263, 234
— capucinum (Ktz.), 263
— gracillimum (Hass.), 234
Stemmatopteris (Cord.), 124, 95
— insignis (Cord.), 124
Stemonitis fusca (Roth), 334
Stephanosphæra (Cohn), 186, 299, 301
Stephensia (Tul.), 358
Stereocaulon ramulosum (Ach.), 279
Sterigma, 86, 340, 367, 370, 372, 376, 384, 386, 389, 394, 275, 304
Stichid, 196, 174
Stictosphæria (Tul.), 373
Stigeoclonium (Ktz.), 276, 284, 417
Stigma, 133, 143
Stigmaria, 118, 90, 91
— ficoides (Brongn.), 118, 90
Stigmatic cell, 17, 27, 133, 143, 158, 15, 109
Stigmatomyces (Karst.), 378
— Baeri (Peyr.), 312
— Muscæ (Karst.), 312
Stigonema (Ag.), 439, 441, 449, 366
— compactum (Kirchn.), 440
— minutum (Hass.), 366
Stigonemeæ, 437, 438, 445, 366
Stipe, 230, 239, 241, 242, 286, 391, 319, 320
Stipule, 92, 174
Stœchospermum (Ktz.), 254
Stolon, 78, 133, 139
Stomate, 19, 72, 78, 93, 107, 144, 169, 23, 82, 153–155
Stomium, 79, 55
Stoneworts, 181
Strand-mycele, 309
Streblonema (Derb.), 239
Stroma, 350, 352, 355, 356, 370, 375, 311
Struthiopteris (L.), 66
— germanica (L.), 66, 69, 77
Struvea (Sond.), 289

STY

Stylospore, 339, 362
Stypocaulon (Ktz.), 249
Subhymenial layer, 355, 368, 372, 393, 394, 305
Submerged leaf, 28, 7, 8
Subsidiary cell, 107
Sulphur, 454, 381, 382
Surirella (Turp.), 426, 357
— splendida (Ktz.), 357
Suspensor, 41, 56, 337, 338, 21, 293–296
Suture, 421
Swarm-cell, 198, 218, 239, 292, 196, 259, 263
Swarming motion, 446
Swarm-spore, 312, 401, 402, 405, 252, 332
Sykidion (Wright), 414
Symbiosis, 318
Symploca (Ktz.), 441, 444, 372, 373
— hydnoides (Ktz.), 372
— violacea (Hauck), 373
Synalissa symphorea (Nyl.), 279
Synange, 94, 122, 71, 94
Syncephalis (Van Tiegh.), 340
Synchytrieæ, 344, 345, 347
Synchytrium Taraxaci (de By.), 347
Synechococcus (Näg.), 447, 449
Synedra (Ehrb.), 426, 357
— Arcus (Ktz.), 357
Syngeneticæ, 188, 237, 256, 232
Syzygites (Ehrb.), 337

TABELLARIA (Ehrb.), 426
Tangle, 244
Tannin-cell 34, 76, 92
Taonia (Ag.), 254
Tapetal cells, 20, 25, 36, 60, 80, 33
Tapete, 20, 80
Targionia (Mich.), 171
Targonieæ, 171
Tayloria (Hook.), 149
Teeth of Hydneæ, 393
— — peristome, 135, 145, 111
— — sheath, 102, 105, 124, 83
Teleutospore, 6, 385, 386, 390, 391, 274, 314, 315
Terfezia (Tul.), 358
Terminal nodule (diatoms), 353
Tetmemorus (Ralfs), 268
Tetrachytrium triceps (Sorok.), 347
Tetrapedia (Reinsch), 447
Tetraphis (Hedw.), 149
— pellucida (Hedw.), 140, 114
Tetraspora (Lk.), 219, 418, 448
— gelatinosa (Desv.), 418
Tetrasporange, 195, 217, 254, 170, 172–174, 177, 182, 183, 186, 190, 230
Tetraspore, 6, 185, 195, 217, 254, 195, 231
Thalassiophyllum (Post.), 244
Thalloid Hepaticæ, 135, 156, 160, 135, 136, 143–159
Thallophytes, 2, 3, 4, 135, 156, 184
Thallus, 132, 184, 191, 306, 430, 432, 153, 188, 199, 212, 227, 230, 307, 358
— compound, 306
Thamnidium (Lk.), 339
Theca, 134
Thecaphora hyalina (Fingerh.), 351
— Lathyri (Kuhn), 351
Thelephoreæ, 391
Thorea (Bory), 211
Thrush-fungus, 38
Tilletia (Tul.), 350, 351
— caries (Tul.), 315, 299
Tilopterideæ, 249,
Tilopteris (Ktz.), 249

UST

Tmesipteris (Bernh.), 61, 63, 41
— tannensis (Bernh.), 41
Todea (Willd.), 89, 90
— superba (Col.), 90
Tolypella (A. Br.), 182
Tolyposporium (Woron.), 351
Tolypothrix (Ktz.), 429, 437, 438, 439, 440, 441
— amphibia (Zopf), 440
Tortula (Hedw.), 149
Trabecule, 41, 52, 147, 23, 32, 33, 112
Tracheide, 18, 76
Trama, 394, 395, 396, 397, 321
Traquairia (Carruth.), 114
Tree-ferns, 71, 75, 78, 85, 57
Tremella (Dill.), 389
— mesenterica (Retz.), 316
Tremellineæ, 343, 388, 389
Tremelloid Uredineæ, 386
Trentepohlia (Mart.), 185, 280, 284, 247
— Bleischii (Rabenh.), 247
Triceratium (Ehrb.), 420, 357
— Favus (Ehrb.), 357
Trichocoma (Jungh.), 319
Trichogyne, 188, 199, 219, 220, 360, 369, 370, 371, 372, 373, 378, 379, 385, 175, 176, 179, 192, 306, 307
Trichomanes (L.), 70, 71, 86, 87, 88, 61
— alatum (Sw.), 86
— pyxidiferum (L.), 86, 61
Trichome, 428, 359
Trichophilus (Web.), 280
Trichophore, 199, 176, 179
'Trichosporange,' 237
Tripoli, 424
Trochopteris (Gardn.), 91
Truffle-family, 357
Truffles, 358
Tuber, 56, 112, 38, 86
— (Mich.), 357, 358
— rufum (Pico), 301
Tuberaceæ, 308, 357
Tubereæ, 358
Tubular organ, 161
Tubuli of Polyporeæ, 393
Tubulinæ, 401
Tuburcinia (Berk.), 350, 351, 352
— Trientalis (Berk.), 352
Tulostoma (Pers.), 399

UDOTEA (Lmx.), 289
Udoteaceæ, 289, 258
Ulodendron (Sternb.), 116
Ulothrix (Ktz.), 185, 277, 278, 279, 246
— implexa (Ktz.), 246
— zonata (Ktz.), 277
Ulotrichaceæ, 187, 273, 277, 246
Ulva (L.), 189, 217, 304, 196
Ulvaceæ, 189, 190, 195, 196, 198, 217, 218, 219, 279, 418, 196–198
Umbelliferæ, 327, 348, 364
Unicellular plants, 184, 281, 284, 286, 288, 427
Unilocular zoosporange, 187, 237, 212, 219, 220, 223
Uredineæ, 4, 312, 315, 383, 391, 274, 314, 315
— (tremelloid), 386, 390
Uredo (Pers.), 383, 385
Uredospore, 6, 385, 386, 274, 314, 315
Urn, 134
Urococcus (Hass.), 418, 350
— insignis (Hass), 350
Urocystis (Rabenh.), 350, 351, 352
Urospora (Aresch.), 273, 276
Usnea barbata (Fr.), 282
Ustilagineæ, 4, 315, 349, 299

UST

Ustilago (Pers.), 351
— carbo (Tul.), 351
— destruens (Tul.), 351
— Hydropiperis (Schm.), 350
— longissima (Tul.), 351
Ustulina (Tul.), 373

Vaccinium Vitis-idæa, 388
Vagine, 134, 135, 144, 146 163, 131
Vallecular canal, 105
Valonia (Gin.), 289, 411, 257
— macrophysa (Ktz.), 257
Valoniaceæ, 28, 257
Valve, 420, 353–355
Valve-view, 421, 353, 354
Vampyrella (Cienk.), 405
Vascular bundle-sheath, 18, 76, 39
— Cryptogams, 1, 2, 10, 2, 4-101
— cylinder, 58, 39
Vaucheria (DC.), 186, 281, 283, 284, 248, 249
— dichotoma (Lyngb.), 249
— sessilis (Vauch.), 248
Veil, 50
Velum partiale, 393
— universale, 393, 397, 318
Venter, 16, 39, 68, 133, 143, 159, 109, 158
Ventral canal-cell, 16, 27, 40, 69, 143, 159, 45
Verbascum, 364
Vesicle, 23, 17, 44
Vessel, 76
Vibrio (Cohn), 440, 451
Vidalia (Lmx.), 194, 209
Vine, 327
— mildew, 364
Violet-stone, 280
Vittaria (Sm.), 67, 85
Volkmannia (Sternb.), 127
Volva, 393, 318
Volvocineæ, 291, 292, 259
Volvox (L.), 4, 292, 295, 259
— globator (L.), 292, 259
Vorticella, 199

Water-beetles, 378
Water-net, 296
Weissia (Hedw.), 149
'Wendungszellen,' 179, 166
Whip-shaped filaments, 178, 163

ZYG

Woodsia (R. Br.), 85, 58
Woodwardia (Sm.), 85, 58
— radicans (Sm.), 78
Woronina (Cornu), 345, 347
'Woronin's hypha,' 360, 373, 385
Wrangelia (Ag.), 209
Wrangeliaceæ, 209, 176, 177

Xanthidium (Ehrb.), 269, 239
— cristatum (Bréb.), 239
Xenococcus (Rostaf.), 444
Xylaria (Hill), 360, 373, 385
Xylarieæ, 373

Yeast, 380, 268, 313,
Yellow cells of Radiolarians, 318

Zamia, 1
Zanardinia (Nardo), 239, 251, 252, 228, 229
— collaris (Crouan), 228, 229
Zippea (Cord.), 124
Zonal view (diatoms), 421
Zonaria (Harv.), 254
Zonate (tetraspores), 195
Zoogamete, 185 (see also under Algæ and Prot.), 196, 222, 246, 247, 252, 260, 263
Zooglœa, 433, 444, 451
Zoosphere, 252, 295, 229
Zoosporange, 220 (see also under Algæ, Fungi and Prot.), 201, 252–255, 258, 259, 272, 277, 287–289, 291, 297
Zoosporangiophore, 326, 275, 287
Zoospore, 6 (see also under Algæ, Fungi, and Prot.), 199, 201, 232, 242, 259, 277, 289, 291, 292, 297
Zygnema (Ktz.), 258, 259, 264, 265, 267, 237
- pectinatum (Ag.), 237
Zygnemaceæ, 187, 258, 259, 264, 236 233
Zygochytrium (Sorok.), 341
Zygodon (H. & T.), 149
Zygogonium (Ktz.), 267
Zygomycetes, 4, 335, 293–299
Zygosperm, 218 (see also under Algæ and Fungi), 222, 233–238, 241, 252, 263, 293–296
'Zygosporeæ,' 3
Zygosporites (Williams.), 114

PRINTED BY
SPOTTISWOODE AND CO., NEW-STREET SQUARE
LONDON

STANDARD SCIENTIFIC WORKS.

WORKS BY JOHN TYNDALL, F.R.S.

FRAGMENTS of SCIENCE. 2 vols. crown 8vo. 16*s.*

HEAT a MODE of MOTION. Crown 8vo. 12*s.*

SOUND. With 204 Woodcuts. Crown 8vo. 10*s.* 6*d.*

ESSAYS on the FLOATING MATTER of the AIR in RELATION to PUTREFACTION and INFECTION. With 24 Woodcuts. Cr. 8vo. 7*s.* 6*d.*

RESEARCHES on DIAMAGNETISM and MAGNE-CRYSTALLIC ACTION, including the Question of Diamagnetic Polarity. Cr. 8vo. 12*s.*

LECTURES on LIGHT, delivered in America in 1872 and 1873. With Portrait, Plate, and Diagrams. Crown 8vo. 5*s.*

LESSONS in ELECTRICITY at the Royal Institution, 1875–6. With 58 Woodcuts. Crown 8vo. 2*s.* 6*d.*

NOTES of a COURSE of SEVEN LECTURES on ELECTRICAL PHENOMENA and THEORIES, delivered at the Royal Institution. Crown 8vo. 1*s.* sewed; 1*s.* 6*d.* cloth.

NOTES of a COURSE of NINE LECTURES on LIGHT, delivered at the Royal Institution. Crown 8vo. 1*s.* sewed; 1*s.* 6*d.* cloth.

FARADAY as a DISCOVERER. Fcp. 8vo. 3*s.* 6*d.*

WATTS' DICTIONARY of CHEMISTRY. Revised and Entirely Rewritten by H. Forster Morley, M.A. D.Sc. Fellow of, and lately Assistant-Professor of Chemistry in, University College, London, and M. M. Pattison Muir, M.A. F.R.S.E. Fellow and Prælector in Chemistry, of Gonville and Caius College, Cambridge; assisted by eminent Contributors. To be published in Four Volumes, 8vo. Volume I. (*Abies—Chemical Change*) price 42*s.* Now ready.

The ELEMENTS of CHEMISTRY, Theoretical and Practical. By W. Allen Miller, M.D. LL.D. Re-edited, with Additions, by H. Macleod, F.C.S. 3 vols. 8vo.

Part I. Chemical Physics, 16*s.* | Part II. Inorganic Chemistry, 24*s.*

Part III. Organic Chemistry, 31*s.* 6*d.*

A SHORT TEXT-BOOK of INORGANIC CHEMISTRY. By Dr. Hermann Kolbe, Professor of Chemistry in the University of Leipzig. Translated and Edited by T. S. Humpidge, Ph.D. B.Sc. (Lond.) With a Coloured Table of Spectra and 66 Wood Engravings. Crown 8vo. 7*s.* 6*d.*

MODERN THEORIES of CHEMISTRY. By Professor Lothair Meyer. Translated from the Fifth Edition of the German by P. Phillipa Bedson, D.Sc. (Lond.) B.Sc. (Vict.) F.C.S. Professor of Chemistry, Durham College of Science; and W. Carleton Williams, B.Sc. (Vict.) F.C.S. Professor of Chemistry, Firth College, Sheffield. 8vo. 18*s.*

The FUNDAMENTAL PRINCIPLES of CHEMISTRY PRACTICALLY TAUGHT, by a New Method. By Robert Galloway, M.R.I.A. F.C.S. With 71 Woodcuts and numerous Exercises and Answers. Crown 8vo. 6*s.* 6*d.*

London: LONGMANS, GREEN, & CO.

SELECT METHODS in CHEMICAL ANALYSIS (chiefly Inorganic). By WILLIAM CROOKES, F.R.S. V.P.C.S. With 37 Illustrations. 8vo. 24*s.*

SPECTRUM ANALYSIS in its APPLICATION to TERRESTRIAL SUBSTANCES, and the Physical Constitution of the Heavenly Bodies. By the late Dr. H. SCHELLEN. Translated by JANE and CAROLINE LASSELL. Edited, with Notes, by Capt. W. DE W. ABNEY, R.E. Second Edition. With 14 Plates (including Ångström's and Cornu's Maps) and 291 Woodcuts. 8vo. £1. 11*s.* 6*d.*

A TREATISE on MAGNETISM, General and Terrestrial. By H. LLOYD, D.D. D.C.L. 8vo. 10*s.* 6*d.*

HANDBOOK of PRACTICAL TELEGRAPHY. By R. S. CULLEY, M.Inst.C.E. Plates and Woodcuts. 8vo. 16*s.*

POPULAR LECTURES on SCIENTIFIC SUBJECTS. By Professor HELMHOLTZ. Translated and Edited by EDMUND ATKINSON. Ph.D. F.C.S. With a Preface by Professor TYNDALL, F.R.S. and 68 Woodcuts. 2 vols. crown 8vo. 15*s.*; or separately, 7*s.* 6*d.* each.

On the SENSATIONS of TONE as a PHYSIOLOGICAL BASIS FOR THE THEORY OF MUSIC. By Professor HELMHOLTZ. Translated by A. J. ELLIS, F.R.S. Second English Edition. Royal 8vo. 21*s.*

The ROTIFERA or 'WHEEL ANIMALCULES.' By C. T. HUDSON, LL.D. and P. H. GOSSE, F.R.S. 6 Parts 4to. price 10*s.* 6*d.* each. Complete in Two Volumes, 4to. price £3. 10*s.*

A COURSE of LECTURES on ELECTRICITY, &c. delivered before the Society of Arts. By GEORGE FORBES, M.A. F.R.S. (L. & E.) Crown 8vo. 5*s.*

The TESTING of MATERIALS of CONSTRUCTION; a Text-book for the Engineering Laboratory, and a Collection of the Results of Experiment. By WILLIAM CAWTHORNE UNWIN, F.R.S. M.Inst.C.E. Professor of Engineering at the Central Institution of the City and Guilds of London Institute. With 5 Plates and 141 Illustrations and Diagrams. 8vo. 21*s.*

OLD and NEW ASTRONOMY. By RICHARD A. PROCTOR.

*** This work will be completed in twelve monthly parts and a supplementary section. In each there will be sixty-four pages, imperial octavo, many cuts, and two plates, or one large folding plate. The price of each part will be 2*s.* 6*d.*; that of the supplementary section, containing tables, index, and preface, 1*s.*

CELESTIAL OBJECTS for COMMON TELESCOPES. By the Rev. T. W. WEBB, M.A. Map, Plate, Woodcuts. Crown 8vo. 9*s.*

ASTRONOMY for AMATEURS: a Practical Manual of Telescopic Research adapted to Moderate Instruments. Edited by J. A. WESTWOOD OLIVER, with the assistance of Messrs. MAUNDER, GRUBB, GORE, DENNING, FRANKS, ELGER, BURNHAM, CAPRON, BACKHOUSE, and others. With several Illustrations. Crown 8vo. 7*s.* 6*d.*

WEATHER CHARTS and STORM WARNINGS. By ROBERT H. SCOTT, M.A. F.R.S. Secretary to the Meteorological Council. With numerous Illustrations. Third Edition, Revised and Enlarged. Crown 8vo. 6*s.*

Catalogue of Books

PUBLISHED BY

MESSRS. LONGMANS, GREEN, & CO.

Abbey and Overton.—*THE ENGLISH CHURCH IN THE EIGHTEENTH CENTURY.* By CHARLES J. ABBEY and JOHN H. OVERTON. Cr. 8vo. 7*s.* 6*d.*

Abbott.—*WORKS BY T. K. ABBOTT.*

THE ELEMENTS OF LOGIC. 12mo. 3*s.*

ELEMENTARY THEORY OF THE TIDES: the Fundamental Theories demonstrated without Mathematics. Crown 8vo. 2*s.*

Acton. — *MODERN COOKERY FOR PRIVATE FAMILIES.* By ELIZA ACTON. With 150 Woodcuts. Fcp. 8vo. 4*s.* 6*d.*

Adams.—*PUBLIC DEBTS:* an Essay on the Science of Finance. By HENRY C. ADAMS, Ph.D. 8vo. 12*s.* 6*d.*

A. K. H. B.—*THE ESSAYS AND CONTRIBUTIONS OF A. K. H. B.* Cr. 8vo.

Autumn Holidays of a Country Parson, 3*s.* 6*d.*
Changed Aspects of Unchanged Truths, 3*s.* 6*d.*
Commonplace Philosopher, 3*s.* 6*d.*
Counsel and Comfort from a City Pulpit, 3*s.* 6*d.*
Critical Essays of a Country Parson, 3*s.* 6*d.*
Graver Thoughts of a Country Parson. Three Series, 3*s.* 6*d.* each.
Landscapes, Churches, and Moralities, 3*s.* 6*d.*
Leisure Hours in Town, 3*s.* 6*d.*
Lessons of Middle Age, 3*s.* 6*d.*
Our Little Life. Two Series, 3*s.* 6*d.* each.
Our Homely Comedy and Tragedy, 3*s.* 6*d.*
Present Day Thoughts, 3*s.* 6*d.*
Recreations of a Country Parson. Three Series, 3*s.* 6*d.* each.
Seaside Musings, 3*s.* 6*d.*
Sunday Afternoons in the Parish Church of a Scottish University City, 3*s.* 6*d.*
'To Meet the Day' through the Christian Year: being a Text of Scripture, with an Original Meditation and a Short Selection in Verse for Every Day, 4*s.* 6*d.*

Allen. — *FORCE AND ENERGY:* a Theory of Dynamics. By GRANT ALLEN. 8vo. 7*s.* 6*d.*

Amos.—*A PRIMER OF THE ENGLISH CONSTITUTION AND GOVERNMENT.* By SHELDON AMOS. Crown 8vo. 6*s.*

Anstey.—*THE BLACK POODLE,* and other Stories. By F. ANSTEY, Author of 'Vice Versâ.' Cr. 8vo. 2*s.* bds.; 2*s.* 6*d.* cl.

Archer.—*MASKS OR FACES?* A Study in the Psychology of Acting. By WILLIAM ARCHER. Crown 8vo. 6*s.* 6*d.*

Aristotle.—*THE WORKS OF.*

THE POLITICS, G. Bekker's Greek Text of Books I. III. IV. (VII.) with an English Translation by W. E. BOLLAND, M.A.; and short Introductory Essays by A. LANG, M.A. Crown 8vo. 7*s.* 6*d.*

THE POLITICS: Introductory Essays. By ANDREW LANG. (From Bolland and Lang's 'Politics.') Crown 8vo. 2*s.* 6*d.*

THE ETHICS; Greek Text, illustrated with Essays and Notes. By Sir ALEXANDER GRANT, Bart. M.A. LL.D. 2 vols. 8vo. 32*s.*

THE NICOMACHEAN ETHICS, Newly Translated into English. By ROBERT WILLIAMS, Barrister-at-Law. Crown 8vo. 7*s.* 6*d.*

Armstrong.—*WORKS BY GEORGE FRANCIS ARMSTRONG, M.A.*

POEMS: Lyrical and Dramatic. Fcp. 8vo. 6*s.*

KING SAUL. (The Tragedy of Israel, Part I.) Fcp. 8vo. 5*s.*

KING DAVID. (The Tragedy of Israel, Part II.) Fcp. 8vo. 6*s.*

KING SOLOMON. (The Tragedy of Israel, Part III.) Fcp. 8vo. 6*s.*

UGONE: A Tragedy. Fcp. 8vo. 6*s.*

A GARLAND FROM GREECE; Poems. Fcp. 8vo. 9*s.*

STORIES OF WICKLOW; Poems. Fcp. 8vo. 9*s.*

VICTORIA REGINA ET IMPERATRIX: a Jubilee Song from Ireland, 1887. 4to. 2*s.* 6*d.* cloth gilt.

MEPHISTOPHELES IN BROADCLOTH: a Satire. Fcp. 8vo. 4*s.*

THE LIFE AND LETTERS OF EDMUND J. ARMSTRONG. Fcp. 8vo. 7*s.* 6*d.*

Armstrong.—*WORKS BY EDMUND J. ARMSTRONG.*

POETICAL WORKS. Fcp. 8vo. 5*s.*

ESSAYS AND SKETCHES. Fcp. 8vo. 5*s.*

Arnold. — *WORKS BY THOMAS ARNOLD, D.D. Late Head-master of Rugby School.*

INTRODUCTORY LECTURES ON MODERN HISTORY, delivered in 1841 and 1842. 8vo. 7*s.* 6*d.*

SERMONS PREACHED MOSTLY IN THE CHAPEL OF RUGBY SCHOOL. 6 vols. crown 8vo. 30*s.* or separately, 5*s.* each.

MISCELLANEOUS WORKS. 8vo. 7*s.* 6*d.*

Arnold.—*A MANUAL OF ENGLISH LITERATURE*, Historical and Critical. By THOMAS ARNOLD, M.A. Cr. 8vo. 7*s.* 6*d.*

Arnott.—*THE ELEMENTS OF PHYSICS OR NATURAL PHILOSOPHY.* By NEIL ARNOTT, M.D. Edited by A. BAIN, LL.D. and A. S. TAYLOR, M.D. F.R.S. Woodcuts. Crown 8vo. 12*s.* 6*d.*

Ashby. — *NOTES ON PHYSIOLOGY FOR THE USE OF STUDENTS PREPARING FOR EXAMINATION.* With 120 Woodcuts. By HENRY ASHBY, M.D. Fcp. 8vo. 5*s.*

Ashby and Wright.—*THE DISEASES OF CHILDREN, MEDICAL AND SURGICAL.* By HENRY ASHBY, M.D. and GEO. A. WRIGHT, B.A. F.R.C.S. 8vo. 21*s.*

Atelier (The) du Lys; or, an Art Student in the Reign of Terror. By the Author of 'Mademoiselle Mori.' Crown 8vo. 2*s.* 6*d.*

Bacon.—*THE WORKS AND LIFE OF.*

COMPLETE WORKS. Edited by R. L. ELLIS, J. SPEDDING, and D. D. HEATH. 7 vols. 8vo. £3. 13*s.* 6*d.*

LETTERS AND LIFE, INCLUDING ALL HIS OCCASIONAL WORKS. Edited by J. SPEDDING. 7 vols. 8vo. £4. 4*s.*

THE ESSAYS; with Annotations. By RICHARD WHATELY, D.D., 8vo. 10*s.* 6*d.*

THE ESSAYS; with Introduction, Notes, and Index. By E. A. ABBOTT, D.D. 2 vols. fcp. 8vo. price 6*s.* Text and Index only, without Introduction and Notes, in 1 vol. fcp. 8vo. 2*s.* 6*d.*

Bagehot.—*WORKS BY WALTER BAGEHOT.*

BIOGRAPHICAL STUDIES. 8vo. 12*s.*

ECONOMIC STUDIES. 8vo. 10*s.* 6*d.*

LITERARY STUDIES. 2 vols. 8vo. 28*s.*

THE POSTULATES OF ENGLISH POLITICAL ECONOMY. Crown 8vo. 2*s.* 6*d.*

A PRACTICAL PLAN FOR ASSIMILATING THE ENGLISH AND AMERICAN MONEY AS A STEP TOWARDS A UNIVERSAL MONEY. Crown 8vo. 2*s.* 6*d.*

The BADMINTON LIBRARY, edited by the DUKE OF BEAUFORT, K.G., assisted by ALFRED E. T. WATSON

Hunting. By the DUKE OF BEAUFORT, K.G. and MOWBRAY MORRIS. With 53 Illustrations by J. Sturgess, J. Charlton, and Agnes M. Biddulph. Crown 8vo. 10*s.* 6*d.*

Fishing. By H. CHOLMONDELEY-PENNELL.

Vol. I. Salmon, Trout, and Grayling. With 150 Illustrations. Cr. 8vo. 10*s.* 6*d.*

Vol. II. Pike and other Coarse Fish. With 58 Illustrations. Cr. 8vo. 10*s.* 6*d.*

Racing and Steeplechasing. By the EARL OF SUFFOLK AND BERKSHIRE, W. G. CRAVEN, &c. With 56 Illustrations by J. Sturgess. Cr. 8vo. 10*s.* 6*d.*

Shooting. By Lord WALSINGHAM and Sir RALPH PAYNE-GALLWEY, Bart. With 21 full-page Illustrations and 149 Woodcuts by A. J. Stuart-Wortley, C. Whymper, J. G. Millais, &c.

Vol. I. Field and Covert. Cr. 8vo. 10*s.* 6*d.*

Vol. II. Moor and Marsh. Cr. 8vo. 10*s.* 6*d.*

Cycling. By VISCOUNT BURY, K.C.M.G. and G. LACY HILLIER. With 19 Plates and 61 Woodcuts by Viscount Bury and Joseph Pennell. Cr. 8vo. 10*s.* 6*d.*

Athletics and Football. By MONTAGUE SHEARMAN. With 6 full-page Illustrations and 45 Woodcuts by Stanley Berkeley, and from Photographs by G. Mitchell. Cr. 8vo. 10*s.* 6*d.*

Boating. By W. B. WOODGATE. With 10 full-page Illustrations and 39 Woodcuts in the Text. Cr. 8vo. 10*s.* 6*d.*

Cricket. By A. G. STEEL and the Hon. R. H. LYTTELTON. With 11 full-page Illustrations and 52 Woodcuts in the Text, by Lucien Davis. Cr. 8vo. 10*s.* 6*d.*

Driving. By the DUKE OF BEAUFORT. With Illustrations by J. Sturgess and G. D. Giles. Crown 8vo. 10*s.* 6*d.*

Fencing, Boxing, and Wrestling. By WALTER H. POLLOCK, F. C. GROVE, WALTER ARMSTRONG, E. B. MICHELL, and M. PRÉVOST. Crown 8vo. 10*s.* 6*d.*

Golf. By HORACE HUTCHINSON, the Right Hon. A. J. BALFOUR, M.P. Sir WILLIAM SIMPSON, Bart. and other Writers. [*In the press.*

Riding. By W. R. WEIR, the EARL OF SUFFOLK AND BERKSHIRE, the DUKE OF BEAUFORT, and A. E. T. WATSON. [*In the press.*

Tennis, Lawn Tennis, Racquets, and Fives. [*In preparation.*

Yachting. By Lord BRASSEY, Lord DUNRAVEN, and other Writers. [*In preparation.*

Bagwell. — *IRELAND UNDER THE TUDORS*, with a Succinct Account of the Earlier History. By RICHARD BAGWELL, M.A. Vols. I. and II. From the first invasion of the Northmen to the year 1578. 2 vols. 8vo. 32*s.*

Bain. — *WORKS BY ALEXANDER BAIN, LL.D.*

MENTAL AND MORAL SCIENCE. Crown 8vo. 10*s.* 6*d.*

SENSES AND THE INTELLECT. 8vo. 15*s.*

EMOTIONS AND THE WILL. 8vo. 15*s.*

LOGIC, DEDUCTIVE AND INDUCTIVE. PART I. *Deduction*, 4*s.* PART II. *Induction*, 6*s.* 6*d.*

PRACTICAL ESSAYS. Cr. 8vo. 2*s.*

Baker.—*BY THE WESTERN SEA:* a Summer Idyll. By JAMES BAKER, Author of 'John Westacott' &c. Crown 8vo. 6*s.*

Baker. — '*WAR WITH CRIME*': being a Selection of Reprinted Papers on Crime, Prison Discipline, &c. By T. BARWICK LL. BAKER. 8vo. 12*s.* 6*d.*

Baker.—*WORKS BY SIR S. W. BAKER.*

EIGHT YEARS IN CEYLON. Crown 8vo. Woodcuts. 5*s.*

THE RIFLE AND THE HOUND IN CEYLON. Crown 8vo. Woodcuts. 5*s.*

Bale.—*A HANDBOOK FOR STEAM USERS;* being Notes on Steam Engine and Boiler Management and Steam Boiler Explosions. By M. POWIS BALE, M.I.M.E. A.M.I.C.E. Fcp. 8vo. 2*s.* 6*d.*

Ball.—*WORKS BY THE RT. HON. J. T. BALL, LL.D. D.C.L.*

THE REFORMED CHURCH OF IRELAND (1537–1886). 8vo. 7*s.* 6*d.*

HISTORICAL REVIEW OF THE LEGISLATIVE SYSTEMS OPERATIVE IN IRELAND, from the Invasion of Henry the Second to the Union (1172–1800). 8vo. 6*s.*

Barker.—*A SHORT MANUAL OF SURGICAL OPERATIONS.* By A. E. J. BARKER, F.R.C.S. With 61 Woodcuts. Crown 8vo. 12*s.* 6*d.*

Barnett.—*PRACTICABLE SOCIALISM:* Essays on Social Reform. By the Rev. S. A. BARNETT, M.A. and Mrs. BARNETT. Crown 8vo. 2*s.* 6*d.*

Becker.—*WORKS BY PROFESSOR BECKER, translated from the German by the Rev. F. METCALFE.*

GALLUS; or, Roman Scenes in the Time of Augustus. Post 8vo. 7*s.* 6*d.*

CHARICLES; or, Illustrations of the Private Life of the Ancient Greeks. Post 8vo. 7*s.* 6*d.*

Beaconsfield.—*WORKS BY THE EARL OF BEACONSFIELD, K.G.*

NOVELS AND TALES. The Hughenden Edition. With 2 Portraits and 11 Vignettes. 11 vols. Crown 8vo. 42*s.*

Endymion.	Henrietta Temple.
Lothair.	Contarini Fleming, &c
Coningsby.	Alroy, Ixion, &c.
Tancred.	Sybil.	The Young Duke, &c.
Venetia.	Vivian Grey.

NOVELS AND TALES. Cheap Edition, complete in 11 vols. Crown 8vo. 1*s.* each, boards; 1*s.* 6*d.* each, cloth.

Bennett and Murray.—*A HANDBOOK OF CRYPTOGAMIC BOTANY.* By A. W. BENNETT, M.A. B.Sc. F.L.S. and GEORGE R. MILNE MURRAY, F.L.S. With 378 Illustrations. 8vo. 16*s.*

Bentley.—*A TEXT-BOOK OF ORGANIC MATERIA MEDICA.* By ROBT. BENTLEY, M.R.C.S.Eng. F.L.S. With 62 Illustrations. Crown 8vo. 7*s.* 6*d.*

Book (The) of Wedding Days. Arranged on the Plan of a Birthday Book. With 96 Illustrated Borders, Frontispiece, and Title-page by WALTER CRANE; and Quotations for each Day. Compiled and Arranged by K. E. J. REID, MAY ROSS, and MABEL BAMFIELD. 4to. 21*s.*

Boultbee.—*A COMMENTARY ON THE 39 ARTICLES* of the Church of England. By the Rev. T. P. BOULTBEE. Cr. 8vo. 6*s.*

Bourne.—*WORKS BY JOHN BOURNE.*

CATECHISM OF THE STEAM ENGINE in its various Applications in the Arts, to which is now added a chapter on Air and Gas Engines, and another devoted to Useful Rules, Tables, and Memoranda. Illustrated by 212 Woodcuts. Crown 8vo. 7*s.* 6*d.*

HANDBOOK OF THE STEAM ENGINE; a Key to the Author's Catechism of the Steam Engine. With 67 Woodcuts. Fcp. 8vo. 9*s.*

RECENT IMPROVEMENTS IN THE STEAM ENGINE. With 124 Woodcuts. Fcp. 8vo. 6*s.*

Bowen.—*A SELECTION FROM THE DESPATCHES AND LETTERS OF THE RIGHT HON. SIR GEORGE F. BOWEN, G.C.M.G. D.C.L. LL.D.* Edited by STANLEY LANE-POOLE, Author of the 'Life of Lord Stratford de Redcliffe.' 2 vols. 8vo.

Bowen. — *HARROW SONGS AND OTHER VERSES.* By EDWARD E. BOWEN. Fcp. 8vo. 2*s* 6*d.*

Brassey. — *WORKS BY LADY BRASSEY.*

A VOYAGE IN THE 'SUNBEAM,' OUR HOME ON THE OCEAN FOR ELEVEN MONTHS.
Library Edition. With 8 Maps and Charts, and 118 Illustrations, 8vo. 21*s.*
Cabinet Edition. With Map and 66 Illustrations, crown 8vo. 7*s.* 6*d.*
School Edition. With 37 Illustrations, fcp. 2*s.* cloth, or 3*s.* white parchment with gilt edges.
Popular Edition. With 60 Illustrations, 4to. 6*d.* sewed, 1*s.* cloth.

SUNSHINE AND STORM IN THE EAST.
Library Edition. With 2 Maps and 114 Illustrations, 8vo. 21*s.*
Cabinet Edition. With 2 Maps and 114 Illustrations, crown 8vo. 7*s.* 6*d.*
Popular Edition. With 103 Illustrations, 4to. 6*d.* sewed, 1*s.* cloth.

IN THE TRADES, THE TROPICS, AND THE 'ROARING FORTIES.'
Cabinet Edition. With Map and 220 Illustrations, crown 8vo. 7*s.* 6*d.*
Popular Edition. With 183 Illustrations, 4to. 6*d.* sewed, 1*s.* cloth.

THE LAST VOYAGE TO INDIA AND AUSTRALIA IN THE 'SUNBEAM.' With Charts and Maps, and 40 Illustrations in Monotone (20 full-page), and nearly 200 Illustrations in the Text from Drawings by R. T. PRITCHETT. 8vo. 21*s.*

THREE VOYAGES IN THE 'SUNBEAM.' Popular Edition. With 346 Illustrations, 4to. 2*s.* 6*d.*

Bray.—*THE PHILOSOPHY OF NECESSITY*; or, Law in Mind as in Matter. By CHARLES BRAY. With Portrait. Crown 8vo. 5*s.*

Brinkley's Astronomy. Re-edited and Revised by J. W. STUBBS, D.D. and F. BRUNNOW, Ph.D. With 49 Diagrams. Crown 8vo. 6*s.*

Browne.—*AN EXPOSITION OF THE 39 ARTICLES*, Historical and Doctrinal. By E. H. BROWNE, D.D., Bishop of Winchester. 8vo. 16*s.*

Bryant.—*EDUCATIONAL ENDS*; or, the Ideal of Personal Development. By SOPHIE BRYANT, D.Sc.Lond. Crown 8vo. 6*s.*

Buckle.—*HISTORY OF CIVILISATION IN ENGLAND AND FRANCE, SPAIN AND SCOTLAND.* By HENRY THOMAS BUCKLE. 3 vols. crown 8vo. 24*s.*

Buckton.—*WORKS BY MRS. C. M. BUCKTON.*

FOOD AND HOME COOKERY. With 11 Woodcuts. Crown 8vo. 2*s.* 6*d.*

HEALTH IN THE HOUSE. With 41 Woodcuts and Diagrams. Crown 8vo. 2*s.*

Bull.—*WORKS BY THOMAS BULL.*

HINTS TO MOTHERS ON THE MANAGEMENT OF THEIR HEALTH during the Period of Pregnancy and in the Lying-in Room. Fcp. 8vo. 1*s.* 6*d.*

THE MATERNAL MANAGEMENT OF CHILDREN IN HEALTH AND DISEASE. Fcp. 8vo. 1*s.* 6*d.*

Bullinger.—*A CRITICAL LEXICON AND CONCORDANCE TO THE ENGLISH AND GREEK NEW TESTAMENT.* By the Rev. E. W. BULLINGER, D.D. Royal 8vo. 15*s.*

Burdett.—*PRINCE, PRINCESS, AND PEOPLE:* an Account of the Social Progress and Development of our own Times, as illustrated by the Public Life and Work of their Royal Highnesses the Prince and Princess of Wales, 1863–1889. By HENRY C. BURDETT. 8vo. 21*s.*

Carlyle. — *THOMAS AND JANE WELSH CARLYLE.*

THOMAS CARLYLE, a History of his Life. By J. A. FROUDE, M.A. Vols. I. and II. 1795–1835, 8vo. 32*s.* Vols. III. and IV. 1834–1881, 8vo. 32*s.*

LETTERS AND MEMORIALS OF JANE WELSH CARLYLE. Prepared for publication by THOMAS CARLYLE, and edited by J. A. FROUDE, M.A. 3 vols. 8vo. 36*s.*

Case.—*PHYSICAL REALISM*; being an Analytical Philosophy from the Physical Objects of Science to the Physical Data of Sense. By THOMAS CASE, M.A. Fellow and Senior Tutor C.C.C. 8vo. 15*s.*

Cecil. — *NOTES OF MY JOURNEY ROUND THE WORLD.* By EVELYN CECIL. With 15 Plates. 1 vol. 8vo. 12*s.* 6*d.*

Chisholm. — *HANDBOOK OF COMMERCIAL GEOGRAPHY.* By G. G. CHISHOLM, M.A., B.Sc. With 29 Maps. 8vo. 16*s.*

Churchill. — *SPEECHES BY LORD RANDOLPH CHURCHILL, M.P.* Selected and Edited, with an Introduction and Notes, by LOUIS J. JENNINGS, M.P. 2 vols. 8vo. 24*s.*

Clavers, the Despot's Champion: a Scots Biography. By A SOUTHERN. Crown 8vo. 7*s.* 6*d.*

Clerk.—*THE GAS ENGINE.* By DUGALD CLERK. With 101 Illustrations and Diagrams. Crown 8vo. 7*s.* 6*d.*

Clodd.—*THE STORY OF CREATION:* a Plain Account of Evolution. By EDWARD CLODD. With 77 Illustrations. Crown 8vo. 6*s.*

Clutterbuck. — *THE SKIPPER IN ARCTIC SEAS.* By W. J. CLUTTERBUCK, one of the Authors of 'Three in Norway.' With numerous Illustrations. Crown 8vo.

Coats.—*A MANUAL OF PATHOLOGY.* By JOSEPH COATS, M.D. Pathologist to the Western Infirmary and the Sick Children's Hospital, Glasgow. With 364 Illustrations. 8vo. 31*s.* 6*d.*

Colenso.—*THE PENTATEUCH AND BOOK OF JOSHUA CRITICALLY EXAMINED.* By J. W. COLENSO, D.D. late Bishop of Natal. Crown 8vo. 6*s.*

Comyn.—*ATHERSTONE PRIORY:* a Tale. By L. N. COMYN. Cr. 8vo. 2*s.* 6*d.*

Conder. — *A HANDBOOK TO THE BIBLE,* or Guide to the Study of the Holy Scriptures derived from Ancient Monuments and Modern Exploration. By F. R. CONDER, and Lieut. C. R. CONDER, R.E. Post 8vo. 7*s.* 6*d.*

Conington. — *WORKS BY JOHN CONINGTON, M.A.*

THE ÆNEID OF VIRGIL. Translated into English Verse. Crown 8vo. 6*s.*

THE POEMS OF VIRGIL. Translated into English Prose. Crown 8vo. 6*s.*

Conybeare & Howson. — *THE LIFE AND EPISTLES OF ST. PAUL.* By the Rev. W. J. CONYBEARE, M.A. and the Very Rev. J. S. HOWSON, D.D.
Library Edition. 2 vols. 8vo. 21*s.*
Student's Edition. 1 vol. crown 8vo. 6*s.*

Cooke. — *TABLETS OF ANATOMY.* By THOMAS COOKE, F.R.C.S. Eng. B.A. B.Sc. M.D. Paris. A selection of the Tablets believed to be most useful to Students generally. Post 4to. 7*s.* 6*d.*

Coolidge.—*SWISS TRAVEL AND SWISS GUIDE-BOOKS.* By W. A. B. COOLIDGE, Fellow of Magdalen College, Oxford, and Editor of the *Alpine Journal.* Crown 8vo. 10*s.* 6*d.*

Cox.—*A GENERAL HISTORY OF GREECE:* from the Earliest Period to the Death of Alexander the Great; with a Sketch of the History to the Present Time. By the Rev. Sir G. W. COX, Bart., M.A. With 11 Maps and Plans. Crown 8vo. 7*s.* 6*d.*

Crawford.—*REMINISCENCES OF FOREIGN TRAVEL.* By ROBERT CRAWFORD, M.A. Author of 'Across the Pampas and the Andes.' Crown 8vo. 5*s.*

Creighton. — *HISTORY OF THE PAPACY DURING THE REFORMATION.* By the Rev. M. CREIGHTON, M.A. 8vo. Vols. I. and II. 1378–1464, 32*s.*; Vols. III. and IV. 1464–1518, 24*s.*

Crookes. — *SELECT METHODS IN CHEMICAL ANALYSIS* (chiefly Inorganic). By WILLIAM CROOKES, F.R.S. V.P.C.S. With 37 Illustrations. 8vo. 24*s.*

Crump. — *WORKS BY ARTHUR CRUMP:*

A SHORT ENQUIRY INTO THE FORMATION OF POLITICAL OPINION, from the Reign of the Great Families to the Advent of Democracy. 8vo. 7*s.* 6*d.*

AN INVESTIGATION INTO THE CAUSES OF THE GREAT FALL IN PRICES WHICH TOOK PLACE COINCIDENTLY WITH THE DEMONETISATION OF SILVER BY GERMANY. 8vo. 6*s.*

Culley.—*HANDBOOK OF PRACTICAL TELEGRAPHY.* By R. S. CULLEY, M. Inst. C.E. Plates and Woodcuts. 8vo. 16*s.*

Curzon.—*RUSSIA IN CENTRAL ASIA IN 1888.* By the Hon. GEORGE CURZON, M.P. Fellow of All Souls' College, Oxford. With Illustrations, Maps, Appendices, and an Index. 1 vol. 8vo.

Davidson.—*AN INTRODUCTION TO THE STUDY OF THE NEW TESTAMENT,* Critical, Exegetical, and Theological. By the Rev. S. DAVIDSON, D.D. LL.D. Revised Edition. 2 vols. 8vo. 30*s.*

Davidson.—*WORKS BY WILLIAM L. DAVIDSON, M.A.*

THE LOGIC OF DEFINITION EXPLAINED AND APPLIED. Crown 8vo. 6*s.*

LEADING AND IMPORTANT ENGLISH WORDS EXPLAINED AND EXEMPLIFIED. Fcp. 8vo. 3*s.* 6*d.*

De Redcliffe.—*THE LIFE OF THE RIGHT HON. STRATFORD CANNING: VISCOUNT STRATFORD DE REDCLIFFE.* By STANLEY LANE-POOLE. With 3 Portraits. 2 vols. 8vo. 36*s.*

De Salis. — *WORKS BY MRS. DE SALIS.*

SAVOURIES À LA MODE. Fcp. 8vo. 1*s.* boards.

ENTRÉES À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

SOUPS AND DRESSED FISH À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

OYSTERS À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

SWEETS AND SUPPER DISHES À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

DRESSED VEGETABLES À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

DRESSED GAME AND POULTRY À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

PUDDINGS AND PASTRY À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

CAKES AND CONFECTIONS À LA MODE. Fcp. 8vo. 1*s.* 6*d.* boards.

De Tocqueville. — *DEMOCRACY IN AMERICA.* By ALEXIS DE TOCQUEVILLE. Translated by HENRY REEVE, C.B. 2 vols. crown 8vo. 16*s.*

Deland. — *WORKS BY MRS. DELAND.*

JOHN WARD, PREACHER: a Story. Crown 8vo. Cabinet Edition, 6*s.*; Popular Edition, 2*s.* boards, 2*s.* 6*d.* cloth.

THE OLD GARDEN, and other Verses. Fcp. 8vo. 5*s.*

FLORIDA DAYS. With 12 Full-page Plates (2 Etched and 4 in Colours), and about 50 Illustrations in the Text, by LOUIS K. HARLOW. 8vo.

Dickinson. — *WORKS BY W. HOWSHIP DICKINSON, M.D. CANTAB.*

ON RENAL AND URINARY AFFECTIONS. With 12 Plates and 122 Woodcuts. 3 vols. 8vo. £3. 4*s.* 6*d.*

THE TONGUE AS AN INDICATOR OF DISEASE: being the Lumbeian Lectures delivered at the Royal College of Physicians in March 1888. 8vo. 7*s.* 6*d.*

Dixon. — *RURAL BIRD LIFE;* Essays on Ornithology, with Instructions for Preserving Objects relating to that Science. By CHARLES DIXON. With 45 Woodcuts. Crown 8vo. 5*s.*

Dowell. — *A HISTORY OF TAXATION AND TAXES IN ENGLAND FROM THE EARLIEST TIMES TO THE YEAR* 1885. By STEPHEN DOWELL, Assistant Solicitor of Inland Revenue. Second Edition, Revised and Altered. (4 vols. 8vo.) Vols. I. and II. The History of Taxation, 21*s.* Vols. III. and IV. The History of Taxes, 21*s.*

Doyle. — *THE OFFICIAL BARONAGE OF ENGLAND.* By JAMES E. DOYLE. Showing the Succession, Dignities, and Offices of every Peer from 1066 to 1885. Vols. I. to III. With 1,600 Portraits, Shields of Arms, Autographs, &c. 3 vols. 4to. £5. 5*s.*

Doyle. — *WORKS BY J. A. DOYLE,* Fellow of All Souls College, Oxford.

THE ENGLISH IN AMERICA: VIRGINIA, MARYLAND, AND THE CAROLINAS. 8vo. 18*s.*

THE ENGLISH IN AMERICA: THE PURITAN COLONIES. 2 vols. 8vo. 36*s.*

Doyle. — *MICAH CLARKE:* his Statement; with some Account of his Journey from Havant to Taunton with Decimus Saxon in the Summer of 1685. Also of the Adventures that befell them during the Western Rebellion, and of their intercourse with James Duke of Monmouth, Lord Grey, and other Persons of Quality. By A. CONAN DOYLE. Crown 8vo. 6*s.*

Dublin University Press Series (The): a Series of Works undertaken by the Provost and Senior Fellows of Trinity College, Dublin.

Abbott's (T. K.) Codex Rescriptus Dublinensis of St. Matthew. 4to. 21*s.*

——— Evangeliorum Versio Antehieronymiana ex Codice Usseriano (Dublinensi). 2 vols. crown 8vo. 21*s.*

Allman's (G. J.) Greek Geometry from Thales to Euclid. 8vo. 10*s.* 6*d.*

Burnside (W. S.) and Panton's (A. W.) Theory of Equations. 8vo. 12*s.* 6*d.*

Casey's (John) Sequel to Euclid's Elements. Crown 8vo. 3*s.* 6*d.*

——— Analytical Geometry of the Conic Sections. Crown 8vo. 7*s.* 6*d.*

Davies's (J. F.) Eumenides of Æschylus. With Metrical English Translation. 8vo. 7*s.*

Dublin Translations into Greek and Latin Verse. Edited by R. Y. Tyrrell. 8vo. 12*s.* 6*d.*

Graves's (R. P.) Life of Sir William Hamilton. 3 vols. 15*s.* each.

Griffin (R. W.) on Parabola, Ellipse, and Hyperbola. Crown 8vo. 6*s.*

Hobart's (W. K.) Medical Language of St. Luke. 8vo. 16*s.*

Leslie's (T. E. Cliffe) Essays in Political Economy. 8vo. 10*s.* 6*d.*

Macalister's (A.) Zoology and Morphology of Vertebrata. 8vo. 10*s.* 6*d.*

[*Continued on next page.*

Dublin University Press Series (The)—*continued.*

MacCullagh's (James) Mathematical and other Tracts. 8vo. 15*s.*

Maguire's (T.) Parmenides of Plato, Text with Introduction, Analysis, &c. 8vo. 7*s.* 6*d.*

Monck's (W. H. S.) Introduction to Logic. Crown 8vo. 5*s.*

Purser's (J. M.) Manual of Histology. Fcp. 8vo. 5*s.*

Roberts's (R. A.) Examples in the Analytic Geometry of Plane Curves. Fcp. 8vo. 5*s.*

Southey's (R.) Correspondence with Caroline Bowles. Edited by E. Dowden. 8vo. 14*s.*

Thornhill's (W. J.) The Æneid of Virgil, freely translated into English Blank Verse. Crown 8vo. 7*s.* 6*d.*

Tyrrell's (R. Y.) Cicero's Correspondence. Vols. I. and II. 8vo. each 12*s.*

——— The Acharnians of Aristophanes, translated into English Verse. Crown 8vo. 2*s.* 6*d.*

Webb's (T. E.) Goethe's Faust, Translation and Notes. 8vo. 12*s.* 6*d.*

——— The Veil of Isis: a Series of Essays on Idealism. 8vo. 10*s.* 6*d.*

Wilkins's (G.) The Growth of the Homeric Poems. 8vo. 6*s.*

Edersheim.—*WORKS BY THE REV. ALFRED EDERSHEIM, D.D.*

THE LIFE AND TIMES OF JESUS THE MESSIAH. 2 vols. 8vo. 24*s.*

PROPHECY AND HISTORY IN RELATION TO THE MESSIAH: the Warburton Lectures, 1880–1884. 8vo. 12*s.*

Ellicott. — *WORKS BY C. J. ELLICOTT, D.D.* Bishop of Gloucester and Bristol.

A CRITICAL AND GRAMMATICAL COMMENTARY ON ST. PAUL'S EPISTLES. 8vo.

I. CORINTHIANS. 16*s.*
GALATIANS. 8*s.* 6*d.*
EPHESIANS. 8*s.* 6*d.*
PASTORAL EPISTLES. 10*s.* 6*d.*
PHILIPPIANS, COLOSSIANS, and PHILEMON. 10*s.* 6*d.*
THESSALONIANS. 7*s.* 6*d.*

HISTORICAL LECTURES ON THE LIFE OF OUR LORD JESUS CHRIST. 8vo. 12*s.*

Ellis.—*MEMOIR OF WILLIAM ELLIS, AND AN ACCOUNT OF HIS CONDUCT-TEACHING.* By ETHEL E. ELLIS. With a Portrait. 8vo. 6*s.*

English Worthies. Fcp. 8vo. 1*s.* each, sewed; 1*s.* 6*d.* each, cloth.

DARWIN. By GRANT ALLEN.

MARLBOROUGH. By G. SAINTSBURY.

SHAFTESBURY (The First Earl). By H. D. TRAILL.

ADMIRAL BLAKE. By DAVID HANNAY.

RALEIGH. By EDMUND GOSSE.

STEELE. By AUSTIN DOBSON.

BEN JONSON. By J. A. SYMONDS.

CANNING. By FRANK H. HILL.

CLAVERHOUSE. By MOWBRAY MORRIS.

Erichsen.—*WORKS BY JOHN ERIC ERICHSEN, F.R.S.*

THE SCIENCE AND ART OF SURGERY: Being a Treatise on Surgical Injuries, Diseases, and Operations. With 1,025 Illustrations. 2 vols. 8vo. 48*s.*

ON CONCUSSION OF THE SPINE, NERVOUS SHOCKS, and other Obscure Injuries of the Nervous System. Cr. 8vo. 10*s.* 6*d.*

Ewald. — *WORKS BY PROFESSOR HEINRICH EWALD,* of Göttingen.

THE ANTIQUITIES OF ISRAEL. Translated from the German by H. S. SOLLY, M.A. 8vo. 12*s.* 6*d.*

THE HISTORY OF ISRAEL. Translated from the German. 8 vols. 8vo. Vols. I. and II. 24*s.* Vols. III. and IV. 21*s.* Vol. V. 18*s.* Vol. VI. 16*s.* Vol. VII. 21*s.* Vol. VIII. with Index to the Complete Work. 18*s.*

Fairbairn.—*WORKS BY SIR W. FAIRBAIRN, BART. C.E.*

A TREATISE ON MILLS AND MILLWORK, with 18 Plates and 333 Woodcuts. 1 vol. 8vo. 25*s.*

USEFUL INFORMATION FOR ENGINEERS. With many Plates and Woodcuts. 3 vols. crown 8vo. 31*s.* 6*d.*

Farrar. — *LANGUAGE AND LANGUAGES.* A Revised Edition of *Chapters on Language and Families of Speech.* By F. W. FARRAR, D.D. Crown 8vo. 6*s.*

Firth.—*OUR KIN ACROSS THE SEA.* By J. C. FIRTH, of Auckland, New Zealand. With a Preface by Mr. FROUDE. Fcp. 8vo. 6*s.*

Fitzwygram. — *HORSES AND STABLES.* By Major-General Sir F. FITZWYGRAM, Bart. With 19 pages of Illustrations. 8vo. 5*s.*

Fletcher.—*CHARACTERS IN 'MACBETH.'* Extracted from 'Studies of Shakespeare.' By GEORGE FLETCHER, 1847. Crown 8vo. 2*s.* 6*d.*

Forbes.—*A COURSE OF LECTURES ON ELECTRICITY*, delivered before the Society of Arts. By GEORGE FORBES. With 17 Illustrations. Crown 8vo. 5*s.*

Ford.—*THE THEORY AND PRACTICE OF ARCHERY.* By the late HORACE FORD. New Edition, thoroughly Revised and Re-written by W. BUTT, M.A. With a Preface by C. J. LONGMAN, M.A. F.S.A. 8vo. 14*s.*

Fox.—*THE EARLY HISTORY OF CHARLES JAMES FOX.* By the Right Hon. Sir G. O. TREVELYAN, Bart.
Library Edition, 8vo. 18*s.*
Cabinet Edition, cr. 8vo. 6*s.*

Francis.—*A BOOK ON ANGLING*; or, Treatise on the Art of Fishing in every branch; including full Illustrated List of Salmon Flies. By FRANCIS FRANCIS. Post 8vo. Portrait and Plates, 15*s.*

Freeman.—*THE HISTORICAL GEOGRAPHY OF EUROPE.* By E. A. FREEMAN. With 65 Maps. 2 vols. 8vo. 31*s.* 6*d.*

Froude.—*WORKS BY JAMES A. FROUDE.*

THE HISTORY OF ENGLAND, from the Fall of Wolsey to the Defeat of the Spanish Armada.
Cabinet Edition, 12 vols. cr. 8vo. £3. 12*s.*
Popular Edition, 12 vols. cr. 8vo. £2. 2*s.*

SHORT STUDIES ON GREAT SUBJECTS. 4 vols. crown 8vo. 24*s.*

CÆSAR: a Sketch. Crown 8vo. 6*s.*

THE ENGLISH IN IRELAND IN THE EIGHTEENTH CENTURY. 3 vols. crown 8vo. 18*s.*

OCEANA; OR, *ENGLAND AND HER COLONIES.* With 9 Illustrations. Crown 8vo. 2*s.* boards, 2*s.* 6*d.* cloth.

THE ENGLISH IN THE WEST INDIES; OR, *THE BOW OF ULYSSES.* With 9 Illustrations. Crown 8vo. 2*s.* boards, 2*s.* 6*d.* cloth.

THE TWO CHIEFS OF DUNBOY; or, an Irish Romance of the Last Century. Crown 8vo. 6*s.*

THOMAS CARLYLE, a History of his Life, 1795 to 1835. 2 vols. 8vo. 32*s.* 1834 to 1881. 2 vols. 8vo. 32*s.*

Gairdner and Coats.—*ON THE DISEASES CLASSIFIED BY THE REGISTRAR-GENERAL AS TABES MESENTERICA.* By W. T. GAIRDNER, M.D. LL.D. *ON THE PATHOLOGY OF PHTHISIS PULMONALIS.* By JOSEPH COATS, M.D. With 28 Illustrations. 8vo. 12*s.* 6*d.*

Galloway.—*THE FUNDAMENTAL PRINCIPLES OF CHEMISTRY PRACTICALLY TAUGHT BY A NEW METHOD.* By ROBERT GALLOWAY, M.R.I.A. Cr. 8vo. 6*s.* 6*d.*

Ganot.—*WORKS BY PROFESSOR GANOT.* Translated by E. ATKINSON, Ph.D. F.C.S.

ELEMENTARY TREATISE ON PHYSICS. With 5 Coloured Plates and 923 Woodcuts. Crown 8vo. 15*s.*

NATURAL PHILOSOPHY FOR GENERAL READERS AND YOUNG PERSONS. With 2 Plates, 518 Woodcuts, and an Appendix of Questions. Cr. 8vo. 7*s.* 6*d.*

Gardiner.—*WORKS BY SAMUEL RAWSON GARDINER, LL.D.*

HISTORY OF ENGLAND, from the Accession of James I. to the Outbreak of the Civil War, 1603–1642. 10 vols. crown 8vo. price 6*s.* each.

A HISTORY OF THE GREAT CIVIL WAR, 1642–1649. (3 vols.) Vol. I. 1642–1644. With 24 Maps. 8vo. 21*s.* Vol. II. 1644–1647. With 21 Maps. 8vo. 24*s.*

Garrod.—*WORKS BY SIR ALFRED BARING GARROD, M.D. F.R.S.*

A TREATISE ON GOUT AND RHEUMATIC GOUT (RHEUMATOID ARTHRITIS). With 6 Plates, comprising 21 Figures (14 Coloured), and 27 Illustrations engraved on Wood. 8vo. 21*s.*

THE ESSENTIALS OF MATERIA MEDICA AND THERAPEUTICS. New Edition, revised and adapted to the New Edition of the British Pharmacopœia, by NESTOR TIRARD, M.D. Cr. 8vo. 12*s.* 6*d.*

Gerard.—*ORTHODOX*: a Novel. By DOROTHEA GERARD. Crown 8vo. 6*s.*

Gibbs.—*ENGLAND AND SOUTH AFRICA.* By EDMUND J. GIBBS. 8vo.

Gibson—*A TEXT-BOOK OF ELEMENTARY BIOLOGY.* By R. J. HARVEY GIBSON, M.A. With 192 Illustrations. Fcp. 8vo. 6*s.*

Godolphin.—*THE LIFE OF THE EARL OF GODOLPHIN*, Lord High Treasurer 1702–1710. By the Hon. HUGH ELLIOT, M.P. 8vo. 15*s.*

Goethe.—*FAUST.* A New Translation, chiefly in Blank Verse; with Introduction and Notes. By JAMES ADEY BIRDS. Crown 8vo. 6*s.*

FAUST. The Second Part. A New Translation in Verse. By JAMES ADEY BIRDS. Crown 8vo. 6*s.*

Gray. — *ANATOMY, DESCRIPTIVE AND SURGICAL.* By HENRY GRAY, F.R.S. With 569 Woodcut Illustrations, a large number of which are coloured. Re-edited by T. PICKERING PICK. Royal 8vo. 36*s.*

Green.—*THE WORKS OF THOMAS HILL GREEN*, late Whyte's Professor of Moral Philosophy, Oxford. Edited by R. L. NETTLESHIP, Fellow of Balliol College (3 vols.) Vols. I. and II.—Philosophical Works. 8vo. 16*s.* each. Vol. III.—Miscellanies. With Index to the three Volumes and Memoir. 8vo. 21*s.*

THE WITNESS OF GOD, AND FAITH: Two Lay Sermons. By T. H. GREEN. Fcp. 8vo. 2*s.*

Greville. — *A JOURNAL OF THE REIGNS OF KING GEORGE IV. KING WILLIAM IV. AND QUEEN VICTORIA.* By the late C. C. F. GREVILLE, Esq. Edited by H. REEVE, C.B. Cabinet Edition. 8 vols. Crown 8vo. 6*s.* each.

Gwilt.—*AN ENCYCLOPÆDIA OF ARCHITECTURE.* By JOSEPH GWILT, F.S.A. Illustrated with more than 1,700 Engravings on Wood. Revised by WYATT PAPWORTH. 8vo. 52*s.* 6*d.*

Haggard.—*WORKS BY H. RIDER HAGGARD.*

SHE. With 32 Illustrations by M. GREIFFENHAGEN and C. H. M. KERR. Crown 8vo. 3*s.* 6*d.*

ALLAN QUATERMAIN. With 31 Illustrations by C. H. M. KERR. Crown 8vo. 3*s.* 6*d.*

MAIWA'S REVENGE; OR, THE WAR OF THE LITTLE HAND. Crown 8vo. 2*s.* boards; 2*s.* 6*d.* cloth.

COLONEL QUARITCH, V.C. A Novel. Crown 8vo. 6*s.*

CLEOPATRA: being an Account of the Fall and Vengeance of Harmachis, the Royal Egyptian, as set forth by his own hand. With 29 Full-page Illustrations by M. Greiffenhagen and R. Caton Woodville. Crown 8vo. 6*s.*

Hargreaves.—*LITERARY WORKERS;* or, Pilgrims to the Temple of Honour. By JOHN GEORGE HARGREAVES, M.A. 4to. 7*s.* 6*d.*

Harte.—*NOVELS BY BRET HARTE.*

IN THE CARQUINEZ WOODS. Fcp. 8vo. 1*s.* boards; 1*s.* 6*d.* cloth.

ON THE FRONTIER. 16mo. 1*s.*

BY SHORE AND SEDGE. 16mo. 1*s.*

Hartwig.—*WORKS BY DR. HARTWIG.*

THE SEA AND ITS LIVING WONDERS. With 12 Plates and 303 Woodcuts. 8vo. 10*s.* 6*d.*

THE TROPICAL WORLD. With 8 Plates, and 172 Woodcuts. 8vo. 10*s.* 6*d.*

THE POLAR WORLD. With 3 Maps, 8 Plates, and 85 Woodcuts. 8vo. 10*s.* 6*d.*

THE SUBTERRANEAN WORLD. With 3 Maps and 80 Woodcuts. 8vo. 10*s.* 6*d.*

THE AERIAL WORLD. With Map, 8 Plates, and 60 Woodcuts. 8vo. 10*s.* 6*d.*

The following books are extracted from the foregoing works by Dr. HARTWIG:—

HEROES OF THE ARCTIC REGIONS. With 19 Illustrations. Crown 8vo. 2*s.* cloth extra, gilt edges.

WONDERS OF THE TROPICAL FORESTS. With 40 Illustrations. Crown 8vo. 2*s.* cloth extra, gilt edges.

WORKERS UNDER THE GROUND; or, Mines and Mining. With 29 Illustrations. Crown 8vo. 2*s.* cloth extra, gilt edges.

MARVELS OVER OUR HEADS. With 29 Illustrations. Crown 8vo. 2*s.* cloth extra, gilt edges.

MARVELS UNDER OUR FEET. With 22 Illustrations. Crown 8vo. 2*s.* cloth extra, gilt edges.

DWELLERS IN THE ARCTIC REGIONS. With 29 Illustrations. Crown 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

WINGED LIFE IN THE TROPICS. With 55 Illustrations. Crown 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

VOLCANOES AND EARTHQUAKES. With 30 Illustrations. Crown 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

WILD ANIMALS OF THE TROPICS. With 66 Illustrations. Crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.

SEA MONSTERS AND SEA BIRDS. With 75 Illustrations. Crown 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

DENIZENS OF THE DEEP. With 117 Illustrations. Crown 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

Hassall.—*THE INHALATION TREATMENT OF DISEASES OF THE ORGANS OF RESPIRATION,* including Consumption. By ARTHUR HILL HASSALL, M.D. With 19 Illustrations of Apparatus. Cr. 8vo. 12*s.* 6*d.*

Havelock. — *MEMOIRS OF SIR HENRY HAVELOCK, K.C.B.* By JOHN CLARK MARSHMAN. Crown 8vo. 3*s.* 6*d.*

Hearn.—*THE GOVERNMENT OF ENGLAND;* its Structure and its Development. By WILLIAM EDWARD HEARN, Q.C. 8vo. 16*s.*

Helmholtz.—*WORKS BY PROFESSOR HELMHOLTZ.*

ON THE SENSATIONS OF TONE AS A PHYSIOLOGICAL BASIS FOR THE THEORY OF MUSIC. Royal 8vo. 28*s.*

POPULAR LECTURES ON SCIENTIFIC SUBJECTS. With 68 Woodcuts. 2 vols. Crown 8vo. 15*s.* or separately, 7*s.* 6*d.* each.

Herschel.—*OUTLINES OF ASTRONOMY.* By Sir J. F. W. HERSCHEL, Bart. M.A. With Plates and Diagrams. Square crown 8vo. 12*s.*

Hester's Venture: a Novel. By the Author of 'The Atelier du Lys.' Crown 8vo. 2*s.* 6*d.*

Hewitt. — *THE DIAGNOSIS AND TREATMENT OF DISEASES OF WOMEN, INCLUDING THE DIAGNOSIS OF PREGNANCY.* By GRAILY HEWITT, M.D. With 211 Engravings. 8vo. 24*s.*

Higginson. — *THE AFTERNOON LANDSCAPE:* Poems and Translations. By THOMAS WENTWORTH HIGGINSON, Colonel U.S. Army. Fcp. 8vo. 5*s.*

Historic Towns. Edited by E. A. FREEMAN, D.C.L. and Rev. WILLIAM HUNT, M.A. With Maps and Plans. Crown 8vo. 3*s.* 6*d.* each.

LONDON. By W. E. LOFTIE.
EXETER. By E. A. FREEMAN.
BRISTOL. By W. HUNT.
OXFORD. By C. W. BOASE.
COLCHESTER. By E. L. CUTTS.
CINQUE PORTS. By MONTAGU BURROWS.
CARLISLE. By MANDELL CREIGHTON.

Holmes.—*A SYSTEM OF SURGERY,* Theoretical and Practical, in Treatises by various Authors. Edited by TIMOTHY HOLMES, M.A. and J. W. HULKE, F.R.S. 3 vols. royal 8vo. £4. 4*s.*

Hopkins.—*CHRIST THE CONSOLER;* a Book of Comfort for the Sick. By ELLICE HOPKINS. Fcp. 8vo. 2*s.* 6*d.*

Howitt.—*VISITS TO REMARKABLE PLACES,* Old Halls, Battle-Fields, Scenes illustrative of Striking Passages in English History and Poetry. By WILLIAM HOWITT. With 80 Illustrations engraved on Wood. Crown 8vo. 5*s.*

Hudson & Gosse.—*THE ROTIFERA OR 'WHEEL-ANIMALCULES.'* By C. T. HUDSON, LL.D. and P. H. GOSSE, F.R.S. With 30 Coloured Plates. In 6 Parts. 4to. 10*s.* 6*d.* each. Complete in 2 vols. 4to. £3. 10*s.* With Supplement, £4. 4*s.* Supplement separately, 12*s.* 6*d.*

Hullah.—*WORKS BY JOHN HULLAH.*

COURSE OF LECTURES ON THE HISTORY OF MODERN MUSIC. 8vo. 8*s.* 6*d.*

COURSE OF LECTURES ON THE TRANSITION PERIOD OF MUSICAL HISTORY. 8vo. 10*s.* 6*d.*

Hume.—*THE PHILOSOPHICAL WORKS OF DAVID HUME.* Edited by T. H. GREEN and T. H. GROSE. 4 vols. 8vo. 56*s.* Or separately, Essays, 2 vols. 28*s.* Treatise of Human Nature. 2 vols. 28*s.*

Hurlbert.—*FRANCE AND HER REPUBLIC:* a Record of Things Seen and Heard in the 'Centennial Year,' 1889. By WM. HENRY HURLBERT, Author of 'Ireland under Coercion.'

Hutchinson. —*WORKS BY HORACE G. HUTCHINSON.*

CRICKETING SAWS AND STORIES. Consisting of scraps of advice to duffers (chiefly on points of cricketing culture) by one of them—of saws not sharp enough to hurt anybody, and of stories picked up between the innings. With rectilinear Illustrations by the Author. 16mo. 1*s.*

THE RECORD OF A HUMAN SOUL. Fcp. 8vo. 3*s.* 6*d.*

Huth.—*THE MARRIAGE OF NEAR KIN,* considered with respect to the Law of Nations, the Result of Experience, and the Teachings of Biology. By ALFRED H. HUTH. Royal 8vo. 21*s.*

In the Olden Time: a Tale of the Peasant War in Germany. By Author of 'Mademoiselle Mori.' Cr. 8vo. 2*s.* 6*d.*

Ingelow.—*WORKS BY JEAN INGELOW.*

POETICAL WORKS. Vols. I. and II. Fcp. 8vo. 12*s.* Vol. III. Fcp. 8vo. 5*s.*

LYRICAL AND OTHER POEMS. Selected from the Writings of JEAN INGELOW. Fcp. 8vo. 2*s.* 6*d.* cloth plain; 3*s.* cloth gilt.

Irving.—*PHYSICAL AND CHEMICAL STUDIES IN ROCK-METAMORPHISM,* based on the Thesis written for the D.Sc. Degree in the University of London, 1888. By the Rev. A. IRVING, D.Sc. Lond. 8vo. 5*s.*

James.—*THE LONG WHITE MOUNTAIN;* or, a Journey in Manchuria, with an Account of the History, Administration, and Religion of that Province. By H. E. JAMES, of Her Majesty's Bombay Civil Service. With Illustrations and a Map. 1 vol. 8vo. 24*s.*

Jameson.—*WORKS BY MRS. JAMESON.*

LEGENDS OF THE SAINTS AND MARTYRS. With 19 Etchings and 187 Woodcuts. 2 vols. 31*s.* 6*d.*

LEGENDS OF THE MADONNA, the Virgin Mary as represented in Sacred and Legendary Art. With 27 Etchings and 165 Woodcuts. 1 vol. 21*s.*

LEGENDS OF THE MONASTIC ORDERS. With 11 Etchings and 88 Woodcuts. 1 vol. 21*s.*

HISTORY OF THE SAVIOUR, His Types and Precursors. Completed by Lady EASTLAKE. With 13 Etchings and 281 Woodcuts. 2 vols. 42*s.*

Jefferies.—*FIELD AND HEDGEROW*: last Essays of RICHARD JEFFERIES. Crown 8vo. 6*s.*

Jenkin. — *PAPERS, LITERARY, SCIENTIFIC, &c.* By the late FLEEMING JENKIN, F.R.S.S. L. & E. Edited by SIDNEY COLVIN, M.A. and J. A. EWING, F.R.S. With Memoir by ROBERT LOUIS STEVENSON. 2 vols. 8vo. 32*s.*

Jessop.—*JUDGE LYNCH:* a Tale of the California Vineyards. By GEORGE H. JESSOP. Crown 8vo. 6*s.*

Johnson.—*THE PATENTEE'S MANUAL;* a Treatise on the Law and Practice of Letters Patent. By J. JOHNSON and J. H. JOHNSON. 8vo. 10*s.* 6*d.*

Johnston.—*A GENERAL DICTIONARY OF GEOGRAPHY*, Descriptive, Physical, Statistical, and Historical; a complete Gazetteer of the World. By KEITH JOHNSTON. Medium 8vo. 42*s.*

Johnstone.—*A SHORT INTRODUCTION TO THE STUDY OF LOGIC.* By LAURENCE JOHNSTONE. Crown 8vo. 2*s.* 6*d.*

Jordan. — *WORKS BY WILLIAM LEIGHTON JORDAN, F.R.G.S.*

THE OCEAN: a Treatise on Ocean Currents and Tides and their Causes. 8vo. 21*s.*

THE NEW PRINCIPLES OF NATURAL PHILOSOPHY. With 13 plates. 8vo. 21*s.*

THE WINDS: an Essay in Illustration of the New Principles of Natural Philosophy. Crown 8vo. 2*s.*

THE STANDARD OF VALUE. 8vo. 6*s.*

Jukes.—*WORKS BY ANDREW JUKES.*

THE NEW MAN AND THE ETERNAL LIFE. Crown 8vo. 6*s.*

THE TYPES OF GENESIS. Crown 8vo. 7*s.* 6*d.*

THE SECOND DEATH AND THE RESTITUTION OF ALL THINGS. Crown 8vo. 3*s.* 6*d.*

THE MYSTERY OF THE KINGDOM. Crown 8vo. 2*s.* 6*d.*

THE NAMES OF GOD IN HOLY SCRIPTURE: a Revelation of His Nature and Relationships. Crown 8vo. 4*s.* 6*d.*

Justinian.—*THE INSTITUTES OF JUSTINIAN;* Latin Text, chiefly that of Huschke, with English Introduction, Translation, Notes, and Summary. By THOMAS C. SANDARS, M.A. 8vo. 18*s.*

Kalisch. — *WORKS BY M. M. KALISCH, M.A.*

BIBLE STUDIES. Part I. The Prophecies of Balaam. 8vo. 10*s.* 6*d.* Part II. The Book of Jonah. 8vo. 10*s.* 6*d.*

COMMENTARY ON THE OLD TESTAMENT; with a New Translation. Vol. I. Genesis, 8vo. 18*s.* or adapted for the General Reader, 12*s.* Vol. II. Exodus, 15*s.* or adapted for the General Reader, 12*s.* Vol. III. Leviticus, Part I. 15*s.* or adapted for the General Reader, 8*s.* Vol. IV. Leviticus, Part II. 15*s.* or adapted for the General Reader, 8*s.*

HEBREW GRAMMAR. With Exercises. Part I. 8vo. 12*s.* 6*d.* Key, 5*s.* Part II. 12*s.* 6*d.*

Kant.—*WORKS BY EMMANUEL KANT.*

CRITIQUE OF PRACTICAL REASON. Translated by T. K. Abbott. 8vo. 12*s.* 6*d.*

INTRODUCTION TO LOGIC, AND HIS ESSAY ON THE MISTAKEN SUBTILTY OF THE FOUR FIGURES. Translated by T. K. Abbott. With Notes by S. T. Coleridge. 8vo. 6*s.*

Kendall.—*WORKS BY MAY KENDALL.*

FROM A GARRET. Crown 8vo. 6*s.*

DREAMS TO SELL; Poems. Fcp. 8vo. 6*s.*

'SUCH IS LIFE': a Novel. Crown 8vo. 6*s.*

Killick.—*HANDBOOK TO MILL'S SYSTEM OF LOGIC.* By the Rev. A. H. KILLICK, M.A. Crown 8vo. 3*s.* 6*d.*

Kirkup.—*AN INQUIRY INTO SOCIALISM.* By THOMAS KIRKUP, Author of the Article on 'Socialism' in the 'Encyclopædia Britannica.' Crown 8vo. 5*s.*

Kolbe.—*A SHORT TEXT-BOOK OF INORGANIC CHEMISTRY.* By Dr. HERMANN KOLBE. Translated from the German by T. S. HUMPIDGE, Ph.D. With a Coloured Table of Spectra and 66 Illustrations. Crown 8vo. 7*s.* 6*d.*

Ladd.—*ELEMENTS OF PHYSIOLOGICAL PSYCHOLOGY:* a Treatise of the Activities and Nature of the Mind from the Physical and Experimental Point of View. By GEORGE T. LADD. 8vo. 21*s.*

Lang.—*WORKS BY ANDREW LANG.*

MYTH, RITUAL, AND RELIGION. 2 vols. crown 8vo. 21*s.*

CUSTOM AND MYTH; Studies of Early Usage and Belief. With 15 Illustrations. Crown 8vo. 7*s.* 6*d.*

LETTERS TO DEAD AUTHORS. Fcp. 8vo. 6*s.* 6*d.*

BOOKS AND BOOKMEN. With 2 Coloured Plates and 17 Illustrations. Cr. 8vo. 6*s.* 6*d.*

GRASS OF PARNASSUS. A Volume of Selected Verses. Fcp. 8vo. 6*s.*

LETTERS ON LITERATURE. Crown 8vo. 6*s.* 6*d.*

BALLADS OF BOOKS. Edited by ANDREW LANG. Fcp. 8vo. 6*s.*

THE BLUE FAIRY BOOK. Edited by ANDREW LANG. With numerous Illustrations by H. J. Ford and G. P. Jacomb Hood. Crown 8vo. 6*s.*

Laughton.—*STUDIES IN NAVAL HISTORY;* Biographies. By J. K. LAUGHTON, M.A. 8vo. 10*s.* 6*d.*

Lavigerie.—*CARDINAL LAVIGERIE AND SLAVERY IN AFRICA.* 1 vol. 8vo.

Lecky.—*WORKS BY W. E. H. LECKY.*

HISTORY OF ENGLAND IN THE EIGHTEENTH CENTURY. 8vo. Vols. I. & II. 1700–1760. 36*s.* Vols. III. & IV. 1760–1784. 36*s.* Vols. V. & VI. 1784–1793. 36*s.*

THE HISTORY OF EUROPEAN MORALS FROM AUGUSTUS TO CHARLEMAGNE. 2 vols. crown 8vo. 16*s.*

HISTORY OF THE RISE AND INFLUENCE OF THE SPIRIT OF RATIONALISM IN EUROPE. 2 vols. crown 8vo. 16*s.*

Lees and Clutterbuck.—*B. C.* 1887, *A RAMBLE IN BRITISH COLUMBIA.* By J. A. LEES and W. J. CLUTTERBUCK, Authors of 'Three in Norway.' With Map and 75 Illustrations from Sketches and Photographs by the Authors. Crown 8vo. 6*s.*

Lewes.—*THE HISTORY OF PHILOSOPHY,* from Thales to Comte. By GEORGE HENRY LEWES. 2 vols. 8vo. 32*s.*

Light through the Crannies.—Parables and Teachings from the other Side. First Series. Crown 8vo. 1*s.* sewed; 1*s.* 6*d.* cloth.

Lindt.—*PICTURESQUE NEW GUINEA.* By J. W. LINDT, F.R.G.S. With 50 Full-page Photographic Illustrations. 4to. 42*s.*

Liveing.—*WORKS BY ROBERT LIVEING, M.A. and M.D. Cantab.*

HANDBOOK ON DISEASES OF THE SKIN. Fcp 8vo. 5*s.*

NOTES ON THE TREATMENT OF SKIN DISEASES. 18mo. 3*s.*

Lloyd.—*A TREATISE ON MAGNETISM,* General and Terrestrial. By H. LLOYD, D.D. D.C.L. 8vo. 10*s.* 6*d.*

Lloyd.—*THE SCIENCE OF AGRICULTURE.* By F. J. LLOYD. 8vo. 12*s.*

Longman.—*HISTORY OF THE LIFE AND TIMES OF EDWARD III.* By WILLIAM LONGMAN. 2 vols. 8vo. 28*s.*

Longman.—*WORKS BY FREDERICK W. LONGMAN, Balliol College, Oxon.*

CHESS OPENINGS. Fcp. 8vo. 2*s.* 6*d.*

FREDERICK THE GREAT AND THE SEVEN YEARS' WAR. Fcp. 8vo. 2*s.* 6*d.*

A NEW POCKET DICTIONARY OF THE GERMAN AND ENGLISH LANGUAGES. Square 18mo. 2*s.* 6*d.*

Longman's Magazine. Published Monthly. Price Sixpence. Vols. 1–13, 8vo. price 5*s.* each.

Longmans' New Atlas. Political and Physical. For the Use of Schools and Private Persons. Consisting of 40 Quarto and 16 Octavo Maps and Diagrams, besides Insets and 16 Quarto Plates of Views, &c. Engraved and Lithographed by EDWARD STANFORD. Edited by GEO. G. CHISHOLM, M.A. B.Sc. Imp. 4to. or imp. 8vo. 12*s.* 6*d.*

Longmore.—*WORKS BY SURGEON-GENERAL SIR T. LONGMORE.*

GUNSHOT INJURIES; their History, Characteristic Features, Complications, and General Treatment. With 58 Illustrations. 8vo. 31*s.* 6*d.*

THE ILLUSTRATED OPTICAL MANUAL; or, Handbook of Instructions for the Guidance of Surgeons in Testing Quality and Range of Vision, and in Distinguishing and dealing with Optical Defects in General. With 74 Drawings and Diagrams. 8vo. 14*s.*

Loudon.—*WORKS BY J. C. LOUDON, F.L.S.*

ENCYCLOPÆDIA OF GARDENING. With 1,000 Woodcuts. 8vo. 21*s.*

ENCYCLOPÆDIA OF AGRICULTURE; the Laying-out, Improvement, and Management of Landed Property. With 1,100 Woodcuts. 8vo. 21*s.*

ENCYCLOPÆDIA OF PLANTS; the Specific Character, &c. of all Plants found in Great Britain. With 12,000 Woodcuts. 8vo. 42*s.*

Lubbock.—*THE ORIGIN OF CIVILIZATION AND THE PRIMITIVE CONDITION OF MAN.* By Sir J. LUBBOCK, Bart. M.P. F.R.S. With Illustrations. 8vo. 18*s.*

Lyall.—*THE AUTOBIOGRAPHY OF A SLANDER.* By EDNA LYALL, Author of 'Donovan,' 'We Two,' &c. Fcp. 8vo. 1*s.* sewed.

Lyra Germanica; Hymns Translated from the German by Miss C. WINKWORTH. Fcp. 8vo. 5*s.*

Macaulay.—*WORKS AND LIFE OF LORD MACAULAY.*

HISTORY OF ENGLAND FROM THE ACCESSION OF JAMES THE SECOND:
Popular Edition, 2 vols. crown 8vo. 5*s.*
Student's Edition, 2 vols. crown 8vo. 12*s.*
People's Edition, 4 vols. crown 8vo. 16*s.*
Cabinet Edition, 8 vols. post 8vo. 48*s.*
Library Edition, 5 vols. 8vo. £4.

CRITICAL AND HISTORICAL ESSAYS, with LAYS of ANCIENT ROME, in 1 volume:
Popular Edition, crown 8vo. 2*s.* 6*d.*
Authorised Edition, crown 8vo. 2*s.* 6*d.* or 3*s.* 6*d.* gilt edges.

CRITICAL AND HISTORICAL ESSAYS:
Student's Edition, 1 vol. crown 8vo. 6*s.*
People's Edition, 2 vols. crown 8vo. 8*s.*
Cabinet Edition, 4 vols. post 8vo. 24*s.*
Library Edition, 3 vols. 8vo. 36*s.*

[*Continued above.*

Macaulay—*WORKS AND LIFE OF LORD MACAULAY—continued.*

ESSAYS which may be had separately price 6*d.* each sewed, 1*s.* each cloth:
Addison and Walpole.
Frederick the Great.
Croker's Boswell's Johnson.
Hallam's Constitutional History.
Warren Hastings. (3*d.* sewed, 6*d.* cloth.)
The Earl of Chatham (Two Essays).
Ranke and Gladstone.
Milton and Machiavelli.
Lord Bacon.
Lord Clive.
Lord Byron, and The Comic Dramatists of the Restoration.

The Essay on Warren Hastings annotated by S. HALES, 1*s.* 6*d.*
The Essay on Lord Clive annotated by H. COURTHOPE BOWEN, M.A. 2*s.* 6*d.*

SPEECHES:
People's Edition, crown 8vo. 3*s.* 6*d.*

MISCELLANEOUS WRITINGS:
People's Edition, 1 vol. crown 8vo. 4*s.* 6*d.*
Library Edition, 2 vols. 8vo. 21*s.*

LAYS OF ANCIENT ROME, &c.
Illustrated by G. Scharf, fcp. 4to. 10*s.* 6*d.*
——— Bijou Edition, 18mo. 2*s.* 6*d.* gilt top.
——— Popular Edition, fcp. 4to. 6*d.* sewed, 1*s.* cloth.
Illustrated by J. R. Weguelin, crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.
Cabinet Edition, post 8vo. 3*s.* 6*d.*
Annotated Edition, fcp. 8vo. 1*s.* sewed 1*s.* 6*d.* cloth, or 2*s.* 6*d.* cloth extra, gilt edges.

MISCELLANEOUS WRITINGS AND SPEECHES:
Popular edition, 1 vol. crown 8vo. 2*s.* 6*d.*
Student's Edition, in 1 vol. crown 8vo. 6*s.*
Cabinet Edition, including Indian Penal Code, Lays of Ancient Rome, and Miscellaneous Poems, 4 vols. post 8vo. 24*s.*

SELECTIONS FROM THE WRITINGS OF LORD MACAULAY. Edited, with Occasional Notes, by the Right Hon. Sir G. O. TREVELYAN, Bart. Crown 8vo. 6*s.*

COMPLETE WORKS OF LORD MACAULAY.
Library Edition, 8 vols. 8vo. £5. 5*s.*
Cabinet Edition, 16 vols. post 8vo. £4. 16*s.*

THE LIFE AND LETTERS OF LORD MACAULAY. By the Right Hon. Sir G. O. TREVELYAN, Bart.
Popular Edition, 1 vol. crown 8vo. 2*s.* 6*d.*
Student's Edition, 1 vol. crown 8vo. 6*s.*
Cabinet Edition, 2 vols. post 8vo. 12*s.*
Library Edition, 2 vols. 8vo. 36*s.*

Macdonald.—*WORKS BY GEORGE MACDONALD, LL.D.*

UNSPOKEN SERMONS. First and Second Series. Crown 8vo. 3*s.* 6*d.* each. Third Series. Crown 8vo. 7*s.* 6*d.*

THE MIRACLES OF OUR LORD. Crown 8vo. 3*s.* 6*d.*

A BOOK OF STRIFE, IN THE FORM OF THE DIARY OF AN OLD SOUL: Poems. 12mo. 6*s.*

Macfarren.—*WORKS BY SIR G. A. MACFARREN.*

LECTURES ON HARMONY, delivered at the Royal Institution. 8vo. 12*s.*

ADDRESSES AND LECTURES, delivered at the Royal Academy of Music, &c. Crown 8vo. 6*s.* 6*d.*

Macleod.—*WORKS BY HENRY D. MACLEOD, M.A.*

THE ELEMENTS OF BANKING. Crown 8vo. 5*s.*

THE THEORY AND PRACTICE OF BANKING. Vol. I. 8vo. 12*s.* Vol. II. 14*s.*

THE THEORY OF CREDIT. 2 vols. 8vo. Vol. I. 7*s.* 6*d.*

[*Vol. II. in the press.*

McCulloch. — *THE DICTIONARY OF COMMERCE AND COMMERCIAL NAVIGATION* of the late J. R. MCCULLOCH, of H.M. Stationery Office. Latest Edition, containing the most recent Statistical Information by A. J. WILSON. 1 vol. medium 8vo. with 11 Maps and 30 Charts, price 63*s.* cloth, or 70*s.* strongly half-bound in russia.

McDougall.—*MEMOIRS OF FRANCIS THOMAS MCDOUGALL, D.C.L. F.R.C.S.* sometime Bishop of Labuan and Sarawak, and of Harriette his Wife. By her Brother, CHARLES JOHN BUNYON. 8vo.

Mademoiselle Mori: a Tale of Modern Rome. By the Author of 'The Atelier du Lys.' Crown 8vo. 2*s.* 6*d.*

Mahaffy.—*A HISTORY OF CLASSICAL GREEK LITERATURE.* By the Rev. J. P. MAHAFFY, M.A. Crown 8vo. Vol. I. Poets, 7*s.* 6*d.* Vol. II. Prose Writers, 7*s.* 6*d.*

Malmesbury. — *MEMOIRS OF AN EX-MINISTER:* an Autobiography. By the Earl of MALMESBURY, G.C.B. Crown 8vo. 7*s.* 6*d.*

Manning.—*THE TEMPORAL MISSION OF THE HOLY GHOST;* or, Reason and Revelation. By H. E. MANNING, D.D. Cardinal-Archbishop. Crown 8vo. 8*s.* 6*d.*

Manuals of Catholic Philosophy (*Stonyhurst Series*):

LOGIC. By RICHARD F. CLARKE, S.J. Crown 8vo. 5*s.*

FIRST PRINCIPLES OF KNOWLEDGE. By JOHN RICKABY, S.J. Crown 8vo. 5*s.*

MORAL PHILOSOPHY (ETHICS AND NATURAL LAW). By JOSEPH RICKABY, S.J. Crown 8vo. 5*s.*

NATURAL THEOLOGY. By BERNARD BOEDDER, S.J. Crown 8vo. 6*s.* 6*d.* [*Nearly ready.*

PSYCHOLOGY. By MICHAEL MAHER, S.J. Crown 8vo. 6*s.* 6*d.* [*Preparing.*

GENERAL METAPHYSICS. By JOHN RICKABY, S.J. Crown 8vo. 5*s.* [*Preparing.*

Martin.—*NAVIGATION AND NAUTICAL ASTRONOMY.* Compiled by Staff-Commander W. R. MARTIN, R.N. Sanctioned for use in the Royal Navy by the Lords Commissioners of the Admiralty. Royal 8vo. 18*s.*

Martineau—*WORKS BY JAMES MARTINEAU, D.D.*

HOURS OF THOUGHT ON SACRED THINGS. Two Volumes of Sermons. 2 vols. crown 8vo. 7*s.* 6*d.* each.

ENDEAVOURS AFTER THE CHRISTIAN LIFE. Discourses. Crown 8vo. 7*s.* 6*d.*

Matthews.—*PEN AND INK.* Papers on Subjects of More or Less Importance. By BRANDER MATTHEWS. Crown 8vo. 5*s.*

Maunder's Treasuries.

BIOGRAPHICAL TREASURY. Edited by W. L. R. CATES. New Edition, with Supplement brought down to 1889, by Rev. JAS. WOOD. Fcp. 8vo. 6*s.*

TREASURY OF NATURAL HISTORY; or, Popular Dictionary of Zoology. Fcp. 8vo. with 900 Woodcuts, 6*s.*

TREASURY OF GEOGRAPHY, Physical, Historical, Descriptive, and Political. With 7 Maps and 16 Plates. Fcp. 8vo. 6*s.*

SCIENTIFIC AND LITERARY TREASURY. Fcp. 8vo. 6*s.*

HISTORICAL TREASURY: Outlines of Universal History, Separate Histories of all Nations. Revised by the Rev. Sir G. W. COX, Bart. M.A. Fcp. 8vo. 6*s.*

TREASURY OF KNOWLEDGE AND LIBRARY OF REFERENCE. Comprising an English Dictionary and Grammar, Universal Gazetteer, Classical Dictionary, Chronology, Law Dictionary, &c. Fcp. 8vo. 6*s.*

[*Continued on next page.*

Maunder's Treasuries—*continued.*

THE TREASURY OF BIBLE KNOWLEDGE. By the Rev. J. AYRE, M.A. With 5 Maps, 15 Plates, and 300 Woodcuts. Fcp. 8vo. 6*s.*

THE TREASURY OF BOTANY. Edited by J. LINDLEY, F.R.S. and T. MOORE, F.L.S. With 274 Woodcuts and 20 Steel Plates. 2 vols. fcp. 8vo. 12*s.*

Max Müller.—*WORKS BY F. MAX MÜLLER, M.A.*

BIOGRAPHICAL ESSAYS. Crown 8vo. 7*s.* 6*d.*

SELECTED ESSAYS ON LANGUAGE, MYTHOLOGY AND RELIGION. 2 vols. crown 8vo. 16*s.*

LECTURES ON THE SCIENCE OF LANGUAGE. 2 vols. crown 8vo. 16*s.*

THREE LECTURES ON THE SCIENCE OF LANGUAGE AND ITS PLACE IN GENERAL EDUCATION, delivered at the Oxford University Extension Meeting, 1889. Crown 8vo. 2*s.*

INDIA, WHAT CAN IT TEACH US? A Course of Lectures delivered before the University of Cambridge. 8vo. 12*s.* 6*d.*

HIBBERT LECTURES ON THE ORIGIN AND GROWTH OF RELIGION, as illustrated by the Religions of India. Crown 8vo. 7*s.* 6*d.*

INTRODUCTION TO THE SCIENCE OF RELIGION: Four Lectures delivered at the Royal Institution. Crown 8vo. 7*s.* 6*d.*

NATURAL RELIGION. The Gifford Lectures, delivered before the University of Glasgow in 1888. Crown 8vo. 10*s.* 6*d.*

THE SCIENCE OF THOUGHT. 8vo. 21*s.*

BIOGRAPHIES OF WORDS, AND THE HOME OF THE ARYAS. Crown 8vo. 7*s.* 6*d.*

A SANSKRIT GRAMMAR FOR BEGINNERS. New and Abridged Edition. By A. A. MACDONELL. Crown 8vo. 6*s.*

May.—*THE CONSTITUTIONAL HISTORY OF ENGLAND SINCE THE ACCESSION OF GEORGE III.* 1760–1870. By the Right Hon. Sir THOMAS ERSKINE MAY, K.C.B. 3 vols. crown 8vo. 18*s.*

Meath.—*WORKS BY THE EARL OF MEATH (Lord Brabazon).*

SOCIAL ARROWS: Reprinted Articles on various Social Subjects. Cr. 8vo. 5*s.*

PROSPERITY OR PAUPERISM? Physical, Industrial, and Technical Training. (Edited by the EARL OF MEATH). 8vo. 5*s.*

Melbourne.—*THE MELBOURNE PAPERS:* being a Selection from Documents in the possession of Earl Cowper, K.G. Edited by LLOYD C. SANDERS, B.A. With Preface by EARL COWPER. 1 vol. 8vo.

Melville.—*NOVELS BY G. J. WHYTE MELVILLE.* Crown 8vo. 1*s.* each, boards; 1*s.* 6*d.* each, cloth.

The Gladiators.	Holmby House.
The Interpreter.	Kate Coventry.
Good for Nothing.	Digby Grand.
The Queen's Maries.	General Bounce.

Mendelssohn.—*THE LETTERS OF FELIX MENDELSSOHN.* Translated by Lady WALLACE. 2 vols. crown 8vo. 10*s.*

Merivale.—*WORKS BY THE VERY REV. CHARLES MERIVALE, D.D. Dean of Ely.*

HISTORY OF THE ROMANS UNDER THE EMPIRE. 8 vols. post 8vo. 48*s.*

THE FALL OF THE ROMAN REPUBLIC: a Short History of the Last Century of the Commonwealth. 12mo. 7*s.* 6*d.*

GENERAL HISTORY OF ROME FROM B.C. 753 TO A.D. 476. Crown 8vo. 7*s.* 6*d.*

THE ROMAN TRIUMVIRATES. With Maps. Fcp. 8vo. 2*s.* 6*d.*

Meyer.—*MODERN THEORIES OF CHEMISTRY.* By Professor LOTHAR MEYER. Translated, from the Fifth Edition of the German, by P. PHILLIPS BEDSON, D.Sc. (Lond.) B.Sc. (Vict.) F.C.S.; and W. CARLETON WILLIAMS, B.Sc. (Vict.) F.C.S. 8vo. 18*s.*

Mill.—*ANALYSIS OF THE PHENOMENA OF THE HUMAN MIND.* By JAMES MILL. With Notes, Illustrative and Critical. 2 vols. 8vo. 28*s.*

Mill.—*WORKS BY JOHN STUART MILL.*

PRINCIPLES OF POLITICAL ECONOMY. Library Edition, 2 vols. 8vo. 30*s.* People's Edition, 1 vol. crown 8vo. 5*s.*

A SYSTEM OF LOGIC, Ratiocinative and Inductive. Crown 8vo. 5*s.*

ON LIBERTY. Crown 8vo. 1*s.* 4*d.*

ON REPRESENTATIVE GOVERNMENT. Crown 8vo. 2*s.*

UTILITARIANISM. 8vo. 5*s.*

EXAMINATION OF SIR WILLIAM HAMILTON'S PHILOSOPHY. 8vo. 16*s.*

NATURE, THE UTILITY OF RELIGION, AND THEISM. Three Essays. 8vo. 5*s.*

Miller. — *WORKS BY W. ALLEN MILLER, M.D. LL.D.*

THE ELEMENTS OF CHEMISTRY, Theoretical and Practical. Re-edited, with Additions, by H. MACLEOD, F.C.S. 3 vols. 8vo.
Vol. I. CHEMICAL PHYSICS, 16*s.*
Vol. II. INORGANIC CHEMISTRY, 24*s.*
Vol. III. ORGANIC CHEMISTRY, 31*s.* 6*d.*

AN INTRODUCTION TO THE STUDY OF INORGANIC CHEMISTRY. With 71 Woodcuts. Fcp. 8vo. 3*s.* 6*d.*

Mitchell.—*A MANUAL OF PRACTICAL ASSAYING.* By JOHN MITCHELL, F.C.S. Revised, with the Recent Discoveries incorporated. By W. CROOKES, F.R.S. 8vo. Woodcuts, 31*s.* 6*d.*

Mitchell.—*DISSOLUTION AND EVOLUTION AND THE SCIENCE OF MEDICINE:* an Attempt to Co-ordinate the necessary Facts of Pathology and to Establish the First Principles of Treatment. By C. PITFIELD MITCHELL. 8vo. 16*s.*

Molesworth. — *MARRYING AND GIVING IN MARRIAGE:* a Novel. By Mrs. MOLESWORTH. Fcp. 8vo. 2*s.* 6*d.*

Mozley.—*THE WORD.* By the Rev. T. MOZLEY, Author of 'Reminiscences of Oriel College and the Oxford Movement.' Crown 8vo.

Mulhall.—*HISTORY OF PRICES SINCE THE YEAR* 1850. By MICHAEL G. MULHALL. Crown 8vo. 6*s.*

Murchison.—*WORKS BY CHARLES MURCHISON, M.D. LL.D. &c.*

A TREATISE ON THE CONTINUED FEVERS OF GREAT BRITAIN. Revised by W. CAYLEY, M.D. Physician to the Middlesex Hospital. 8vo. with numerous Illustrations, 25*s.*

CLINICAL LECTURES ON DISEASES OF THE LIVER, JAUNDICE, AND ABDOMINAL DROPSY. Revised by T. LAUDER BRUNTON, M.D. and Sir JOSEPH FAYRER, M.D. 8vo. with 43 Illustrations, 24*s.*

Murray.—*A DANGEROUS CATSPAW:* a Story. By DAVID CHRISTIE MURRAY and HENRY MURRAY. Crown 8vo. 6*s.*

Murray.—*GOBI OR SHAMO:* a Story of Three Songs. (*The Great Desert of Gobi or Shamo.*—Cornwell's Geography.) By G. G. A. MURRAY, New College, Oxford. 1 vol. crown 8vo.

Murray and Herman. — *WILD DARRIE:* a Story. By CHRISTIE MURRAY and HENRY HERMAN. Crown 8vo. 6*s.*

Nelson.—*LETTERS AND DESPATCHES OF HORATIO, VISCOUNT NELSON.* Selected and arranged by JOHN KNOX LAUGHTON, M.A. 8vo. 16*s.*

Nesbit.—*WORKS BY E. NESBIT.*

LAYS AND LEGENDS. Cr. 8vo. 5*s.*

LEAVES OF LIFE: Verses. Cr. 8vo. 5*s.*

Newman.—*ON THE DISEASES OF THE KIDNEY AMENABLE TO SURGICAL TREATMENT.* By DAVID NEWMAN, M.D. 8vo. 16*s.*

Newman.— *WORKS BY CARDINAL NEWMAN.*

APOLOGIA PRO VITÂ SUÂ. Crown 8vo. 6*s.*

THE IDEA OF A UNIVERSITY DEFINED AND ILLUSTRATED. Crown 8vo. 7*s.*

HISTORICAL SKETCHES. 3 vols. crown 8vo. 6*s.* each.

THE ARIANS OF THE FOURTH CENTURY. Crown 8vo. 6*s.*

SELECT TREATISES OF ST. ATHANASIUS IN CONTROVERSY WITH THE ARIANS. Freely Translated. 2 vols. crown 8vo. 15*s.*

DISCUSSIONS AND ARGUMENTS ON VARIOUS SUBJECTS. Crown 8vo. 6*s.*

AN ESSAY ON THE DEVELOPMENT OF CHRISTIAN DOCTRINE. Crown 8vo. 6*s.*

CERTAIN DIFFICULTIES FELT BY ANGLICANS IN CATHOLIC TEACHING CONSIDERED. Vol. 1, crown 8vo. 7*s.* 6*d.*; Vol. 2, crown 8vo. 5*s.* 6*d.*

THE VIA MEDIA OF THE ANGLICAN CHURCH, ILLUSTRATED IN LECTURES &c. 2 vols. crown 8vo. 6*s.* each.

ESSAYS, CRITICAL AND HISTORICAL. 2 vols. crown 8vo. 12*s.*

ESSAYS ON BIBLICAL AND ON ECCLESIASTICAL MIRACLES. Crown 8vo. 6*s.*

AN ESSAY IN AID OF A GRAMMAR OF ASSENT. 7*s.* 6*d.*

PRESENT POSITION OF CATHOLICS IN ENGLAND. Crown 8vo. 7*s.*

CALLISTA: a Tale of the Third Century. Crown 8vo. 6*s.*

THE DREAM OF GERONTIUS. 16mo. 6*d.* sewed, 1*s.* cloth.

VERSES ON VARIOUS OCCASIONS. Crown 8vo. 6*s.*

Newnham.—*WORKS BY THE REV. H. P. NEWNHAM.*

THE ALL-FATHER: Sermons preached in a Village Church. With Preface by EDNA LYALL. Cr. 8vo. 4*s.* 6*d.*

THY HEART WITH MY HEART: Four Letters on the Holy Communion. 18mo. 3*d.* sewed; 6*d.* cloth limp; 8*d.* cl.

Noble.—*HOURS WITH A THREE-INCH TELESCOPE.* By Captain W. NOBLE. With a Map of the Moon. Cr. 8vo. 4*s.* 6*d.*

Northcott.—*LATHES AND TURNING*, Simple, Mechanical, and Ornamental. By W. H. NORTHCOTT. With 338 Illustrations. 8vo. 18*s.*

Oliphant.—*NOVELS BY MRS. OLIPHANT.*

MADAM. Cr. 8vo. 1*s.* bds.; 1*s.* 6*d.* cl.

IN TRUST.—Cr. 8vo. 1*s.* bds.; 1*s.* 6*d.* cl.

LADY CAR: the Sequel of a Life. Crown 8vo. 6*s.*

Oliver.—*ASTRONOMY FOR AMATEURS:* a Practical Manual of Telescopic Research adapted to Moderate Instruments. Edited by J. A. WESTWOOD OLIVER. With several Illustrations. Crown 8vo. 7*s.* 6*d.*

Owen.—*A MANUAL OF ANATOMY FOR SENIOR STUDENTS.* By EDMUND OWEN, M.B. F.R.C.S. Surgeon to St. Mary's Hospital, London, and co-Lecturer on Surgery, late Lecturer on Anatomy in its Medical School. With numerous Illustrations. Crown 8vo.

Owen.—*THE COMPARATIVE ANATOMY AND PHYSIOLOGY OF THE VERTEBRATE ANIMALS.* By Sir RICHARD OWEN, K.C.B. &c. With 1,472 Woodcuts. 3 vols. 8vo. £3. 13*s.* 6*d.*

Paget.—*WORKS BY SIR JAMES PAGET, BART. F.R.S. D.C.L. &c.*

CLINICAL LECTURES AND ESSAYS. Edited by F. HOWARD MARSH. 8vo. 15*s.*

LECTURES ON SURGICAL PATHOLOGY. Re-edited by the AUTHOR and W. TURNER, M.B. 8vo. with 131 Woodcuts, 21*s.*

Pasteur.—*LOUIS PASTEUR*, his Life and Labours. By his SON-IN-LAW. Translated from the French by Lady CLAUD HAMILTON. Crown 8vo. 7*s.* 6*d.*

Payen.—*INDUSTRIAL CHEMISTRY;* a Manual for Manufacturers and for Colleges or Technical Schools; a Translation of PAYEN'S 'Précis de Chimie Industrielle.' Edited by B. H. PAUL. With 698 Woodcuts. Medium 8vo. 42*s.*

Payn.—*NOVELS BY JAMES PAYN.*

THE LUCK OF THE DARRELLS. Crown 8vo. 1*s.* boards; 1*s.* 6*d.* cloth.

THICKER THAN WATER. Crown 8vo. 1*s.* boards; 1*s.* 6*d.* cloth.

Pears.—*THE FALL OF CONSTANTINOPLE:* being the Story of the Fourth Crusade. By EDWIN PEARS. 8vo. 16*s.*

Pennell.—*OUR SENTIMENTAL JOURNEY THROUGH FRANCE AND ITALY.* By JOSEPH and ELIZABETH ROBINS PENNELL. With a Map and 120 Illustrations by Joseph Pennell. Crown 8vo. 6*s.* cloth or vegetable vellum.

Perring.—*HARD KNOTS IN SHAKESPEARE.* By Sir PHILIP PERRING, Bart. 8vo. 7*s.* 6*d.*

Pole.—*THE THEORY OF THE MODERN SCIENTIFIC GAME OF WHIST.* By W. POLE, F.R.S. Fcp. 8vo. 2*s.* 6*d.*

Pollock.—*A NINE MEN'S MORRICE:* Stories Collected and Re-collected. By WALTER HERRIES POLLOCK. Crown 8vo. 6*s.*

Porter.—*THE HISTORY OF THE CORPS OF ROYAL ENGINEERS.* By Major-General WHITWORTH PORTER, R.E. 2 vols. 8vo. 36*s.*

Prendergast.—*IRELAND*, from the Restoration to the Revolution, 1660–1690. By JOHN P. PRENDERGAST. 8vo. 5*s.*

Proctor.—*WORKS BY R. A. PROCTOR.*

OLD AND NEW ASTRONOMY. 12 Parts, 2*s.* 6*d.* each. Supplementary Section, 1*s.* Complete in 1 vol. 4to. 36*s.*

[*In course of publication.*

THE ORBS AROUND US; a Series of Essays on the Moon and Planets, Meteors and Comets. With Chart and Diagrams, crown 8vo. 5*s.*

OTHER WORLDS THAN OURS; The Plurality of Worlds Studied under the Light of Recent Scientific Researches. With 14 Illustrations, crown 8vo. 5*s.*

THE MOON; her Motions, Aspects, Scenery, and Physical Condition. With Plates, Charts, Woodcuts, &c. Cr. 8vo. 5*s.*

UNIVERSE OF STARS; Presenting Researches into and New Views respecting the Constitution of the Heavens. With 22 Charts and 22 Diagrams, 8vo. 10*s.* 6*d.*

LARGER STAR ATLAS for the Library, in 12 Circular Maps, with Introduction and 2 Index Pages. Folio, 15*s.* or Maps only, 12*s.* 6*d.*

THE STUDENT'S ATLAS. In Twelve Circular Maps on a Uniform Projection and one Scale, with Two Index Maps. With a letterpress Introduction illustrated by several cuts. 8vo. 5*s.*

NEW STAR ATLAS for the Library, the School, and the Observatory, in 12 Circular Maps. Crown 8vo. 5*s.*

[*Continued on next page.*

Proctor.—*WORKS BY R. A. PROCTOR, Continued.*

LIGHT SCIENCE FOR LEISURE HOURS; Familiar Essays on Scientific Subjects. 3 vols. crown 8vo. 5*s.* each.

CHANCE AND LUCK; a Discussion of the Laws of Luck, Coincidences, Wagers, Lotteries, and the Fallacies of Gambling &c. Crown 8vo. 2*s.* boards; 2*s.* 6*d.* cloth.

STUDIES OF VENUS-TRANSITS; an Investigation of the Circumstances of the Transits of Venus in 1874 and 1882. With 7 Diagrams and 10 Plates. 8vo. 5*s.*

HOW TO PLAY WHIST: WITH THE LAWS AND ETIQUETTE OF WHIST. Crown 8vo. 3*s.* 6*d.*

HOME WHIST: an Easy Guide to Correct Play. 16mo. 1*s.*

THE POETRY OF ASTRONOMY. A Series of Familiar Essays. Crown 8vo. 5*s.*

THE STARS IN THEIR SEASONS An Easy Guide to a Knowledge of the Star Groups, in 12 Large Maps. Imperial 8vo. 5*s.*

STAR PRIMER. Showing the Starry Sky Week by Week, in 24 Hourly Maps. Crown 4to. 2*s.* 6*d.*

THE SEASONS PICTURED IN 48 *SUN-VIEWS OF THE EARTH,* and 24 Zodiacal Maps, &c. Demy 4to. 5*s.*

STRENGTH AND HAPPINESS. Crown 8vo. 5*s.*

STRENGTH: How to get Strong and keep Strong, with Chapters on Rowing and Swimming, Fat, Age, and the Waist. With 9 Illustrations. Crown 8vo. 2*s.*

ROUGH WAYS MADE SMOOTH. Familiar Essays on Scientific Subjects. Crown 8vo. 5*s.*

OUR PLACE AMONG INFINITIES. A Series of Essays contrasting our Little Abode in Space and Time with the Infinities Around us. Crown 8vo. 5*s.*

THE EXPANSE OF HEAVEN. Essays on the Wonders of the Firmament. Crown 8vo. 5*s.*

THE GREAT PYRAMID, OBSERVATORY, TOMB, AND TEMPLE. With Illustrations. Crown 8vo. 5*s.*

PLEASANT WAYS IN SCIENCE. Crown 8vo. 5*s.*

MYTHS AND MARVELS OF ASTRONOMY. Crown 8vo. 5*s.*

NATURE STUDIES. By GRANT ALLEN, A. WILSON, T. FOSTER, E. CLODD, and R. A. PROCTOR. Crown 8vo. 5*s.*

LEISURE READINGS. By E. CLODD, A. WILSON, T. FOSTER, A. C. RANYARD, and R. A. PROCTOR. Crown 8vo. 5*s.*

Prothero.—*THE PIONEERS AND PROGRESS OF ENGLISH FARMING.* By ROWLAND E. PROTHERO. Crown 8vo. 5*s.*

Pryce.—*THE ANCIENT BRITISH CHURCH:* an Historical Essay. By JOHN PRYCE, M.A. Canon of Bangor. Crown 8vo. 6*s.*

Quain's Elements of Anatomy. The Ninth Edition. Re-edited by ALLEN THOMSON, M.D. LL.D. F.R.S.S. L. & E. EDWARD ALBERT SCHÄFER, F.R.S. and GEORGE DANCER THANE. With upwards of 1,000 Illustrations engraved on Wood, of which many are Coloured. 2 vols. 8vo. 18*s.* each.

Quain.—*A DICTIONARY OF MEDICINE.* By Various Writers. Edited by R. QUAIN, M.D. F.R.S. &c. With 138 Woodcuts. Medium 8vo. 31*s.* 6*d.* cloth, or 40*s.* half-russia; to be had also in 2 vols. 34*s.* cloth.

Rawlinson.—*THE HISTORY OF PHŒNICIA.* By GEORGE RAWLINSON, M.A. Canon of Canterbury, &c. With numerous Illustrations. 8vo. 24*s.*

Reader.—*WORKS BY EMILY E. READER.*

ECHOES OF THOUGHT: a Medley of Verse. Fcp. 8vo. 5*s.* cloth, gilt top.

THE GHOST OF BRANKINSHAW and other Tales. With 9 Full-page Illustrations. Fcp. 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

VOICES FROM FLOWER-LAND, in Original Couplets. A Birthday-Book and Language of Flowers. 16mo. 1*s.* 6*d.* limp cloth; 2*s.* 6*d.* roan, gilt edges, or in vegetable vellum, gilt top.

FAIRY PRINCE FOLLOW-MY-LEAD; or, the *MAGIC BRACELET.* Illustrated by WM. READER. Crown 8vo. 2*s.* 6*d.* gilt edges; or 3*s.* 6*d.* vegetable vellum, gilt edges.

Reeve.—*COOKERY AND HOUSEKEEPING.* By Mrs. HENRY REEVE. With 8 Coloured Plates and 37 Woodcuts. Crown 8vo. 5*s.*

Rendle and Norman.—*THE INNS OF OLD SOUTHWARK,* and their Associations. By WILLIAM RENDLE, F.R.C.S. Author of 'Old Southwark and its People,' and PHILIP NORMAN, F.S.A. With numerous Illustrations. Royal 8vo. 28*s.*

Reply (A) to Dr. Lightfoot's Essays. By the Author of 'Supernatural Religion.' 1 vol. 8vo. 6*s.*

Rich.—*A DICTIONARY OF ROMAN AND GREEK ANTIQUITIES.* With 2,000 Woodcuts. By A. RICH, B.A. Cr. 8vo. 7*s.* 6*d.*

Richardson.—*WORKS BY BENJAMIN WARD RICHARDSON, M.D.*

THE HEALTH OF NATIONS: a Review of the Works—Economical, Educational, Sanitary, and Administrative—of EDWIN CHADWICK, C.B. With a Biographical Dissertation by BENJAMIN WARD RICHARDSON, M.D. F.R.S. 2 vols. 8vo. 28*s.*

THE COMMONHEALTH: a Series of Essays on Health and Felicity for Every-Day Readers. Crown 8vo. 6*s.*

THE SON OF A STAR: a Romance of the Second Century. Crown 8vo. 6*s.*

Riley.—*ATHOS;* or, the Mountain of the Monks. By ATHELSTAN RILEY, M.A. F.R.G.S. With Map and 29 Illustrations. 8vo. 21*s.*

Riley. — *OLD-FASHIONED ROSES.* Verses and Sonnets. By J. W. RILEY. Fcp. 8vo. 5*s.*

Rivers.—*WORKS BY THOMAS RIVERS.*

THE ORCHARD-HOUSE. With 25 Woodcuts. Crown 8vo. 5*s.*

THE MINIATURE FRUIT GARDEN; or, the Culture of Pyramidal and Bush Fruit Trees, with Instructions for Root Pruning. With 32 Illustrations. Fcp. 8vo. 4*s.*

Roberts.—*GREEK THE LANGUAGE OF CHRIST AND HIS APOSTLES.* By ALEXANDER ROBERTS, D.D. 8vo. 18*s.*

Robinson. — *THE NEW ARCADIA,* and other Poems. By A. MARY F. ROBINSON. Crown 8vo. 6*s.*

Roget.—*THESAURUS OF ENGLISH WORDS AND PHRASES,* Classified and Arranged so as to facilitate the Expression of Ideas and assist in Literary Composition. By PETER M. ROGET. Crown 8vo. 10*s.* 6*d.*

Ronalds. — *THE FLY-FISHER'S ENTOMOLOGY.* By ALFRED RONALDS. With 20 Coloured Plates. 8vo. 14*s.*

Russell.—*A LIFE OF LORD JOHN RUSSELL (EARL RUSSELL, K.G.).* By SPENCER WALPOLE, Author of 'A History of England from 1815.' With 2 Portraits. 2 vols. 8vo. 36*s.*

Schäfer. — *THE ESSENTIALS OF HISTOLOGY, DESCRIPTIVE AND PRACTICAL.* For the use of Students. By E. A. SCHÄFER, F.R.S. With 281 Illustrations. 8vo. 6*s.* or Interleaved with Drawing Paper, 8*s.* 6*d.*

Schellen. — *SPECTRUM ANALYSIS IN ITS APPLICATION TO TERRESTRIAL SUBSTANCES,* and the Physical Constitution of the Heavenly Bodies. By Dr. H. SCHELLEN. Translated by JANE and CAROLINE LASSELL. Edited by Capt. W. DE W. ABNEY. With 14 Plates (including Angström's and Cornu's Maps) and 291 Woodcuts. 8vo. 31*s.* 6*d.*

Scott.—*WEATHER CHARTS AND STORM WARNINGS.* By ROBERT H. SCOTT, M.A. F.R.S. With numerous Illustrations. Crown 8vo. 6*s.*

Seebohm.—*WORKS BY FREDERIC SEEBOHM.*

THE OXFORD REFORMERS — JOHN COLET, ERASMUS, AND THOMAS MORE; a History of their Fellow-Work. 8vo. 14*s.*

THE ENGLISH VILLAGE COMMUNITY Examined in its Relations to the Manorial and Tribal Systems, &c. 13 Maps and Plates. 8vo. 16*s.*

THE ERA OF THE PROTESTANT REVOLUTION. With Map. Fcp. 8vo. 2*s.* 6*d.*

Sennett. — *THE MARINE STEAM-ENGINE;* a Treatise for the use of Engineering Students and Officers of the Royal Navy. By RICHARD SENNETT, Engineer-in-Chief of the Royal Navy. With 244 Illustrations. 8vo. 21*s.*

Sewell. — *STORIES AND TALES.* By ELIZABETH M. SEWELL. Crown 8vo. 1*s.* 6*d.* each, cloth plain; 2*s.* 6*d.* each, cloth extra, gilt edges:—

Amy Herbert.	Margaret Percival.
The Earl's Daughter.	Laneton Parsonage.
The Experience of Life.	Ursula.
A Glimpse of the World.	Gertrude.
Cleve Hall.	Ivors.
Katharine Ashton.	

Shakespeare. — *BOWDLER'S FAMILY SHAKESPEARE.* Genuine Edition, in 1 vol. medium 8vo. large type, with 36 Woodcuts, 14*s.* or in 6 vols. fcp. 8vo. 21*s.*

OUTLINES OF THE LIFE OF SHAKESPEARE. By J. O. HALLIWELL-PHILLIPPS, F.R.S. 2 vols. Royal 8vo. £1. 1*s.*

SHAKESPEARE'S TRUE LIFE. By JAMES WALTER. With 400 Illustrations. 4to. 15*s.*

Short.—*SKETCH OF THE HISTORY OF THE CHURCH OF ENGLAND TO THE REVOLUTION OF 1688.* By T. V. SHORT, D.D. Crown 8vo. 7*s.* 6*d.*

Smith, H. F.—*THE HANDBOOK FOR MIDWIVES.* By HENRY FLY SMITH, M.B. Oxon. M.R.C.S. With 41 Woodcuts. Crown 8vo. 5*s.*

Smith, J. H.—*THE WHITE UMBRELLA IN MEXICO.* With numerous Illustrations. By J. HOPKINSON SMITH. Fcp. 8vo. 6*s.* 6*d.*

Smith, R. Bosworth. — *CARTHAGE AND THE CARTHAGINIANS.* By R. BOSWORTH SMITH, M.A. Maps, Plans, &c. Crown 8vo. 10*s.* 6*d.*

Smith, R. H.—*GRAPHICS;* or, The Art of Calculation by Drawing Lines, applied to Mathematics, Theoretical Mechanics, and Engineering, including the Kinetics and Dynamics of Machinery, &c. By ROBERT H. SMITH.

PART I. Text, with separate Atlas of Plates, 8vo. 15*s.*

Smith, T.—*A MANUAL OF OPERATIVE SURGERY ON THE DEAD BODY.* By THOMAS SMITH, Surgeon to St. Bartholomew's Hospital. A New Edition, re-edited by W. J. WALSHAM. With 46 Illustrations. 8vo. 12*s.*

Southey.—*THE POETICAL WORKS OF ROBERT SOUTHEY,* with the Author's last Corrections and Additions. Medium 8vo. with Portrait, 14*s.*

Stanley. — *A FAMILIAR HISTORY OF BIRDS.* By E. STANLEY, D.D. Revised and enlarged, with 160 Woodcuts. Crown 8vo. 6*s.*

Steel.—*WORKS BY J. H. STEEL, M.R.C.V.S.*

A TREATISE ON THE DISEASES OF THE DOG; being a Manual of Canine Pathology. Especially adapted for the Use of Veterinary Practitioners and Students. With 88 Illustrations. 8vo. 10*s.* 6*d.*

A TREATISE ON THE DISEASES OF THE OX; being a Manual of Bovine Pathology specially adapted for the use of Veterinary Practitioners and Students. With 2 Plates and 117 Woodcuts. 8vo. 15*s.*

A TREATISE ON DISEASES OF THE SHEEP: being a Manual of Ovine Pathology for the use of Veterinary Practitioners and Students. Illustrated. 8vo.

Stephen. — *ESSAYS IN ECCLESIASTICAL BIOGRAPHY.* By the Right Hon. Sir J. STEPHEN, LL.D. Crown 8vo. 7*s.* 6*d.*

Stevenson.—*WORKS BY ROBERT LOUIS STEVENSON.*

A CHILD'S GARDEN OF VERSES. Small fcp. 8vo. 5*s.*

THE DYNAMITER. Fcp. 8vo. 1*s.* swd. 1*s.* 6*d.* cloth.

STRANGE CASE OF DR. JEKYLL AND MR. HYDE. Fcp. 8vo. 1*s.* sewed; 1*s.* 6*d.* cloth.

Stevenson and Osbourne.—*THE WRONG BOX.* By ROBERT LOUIS STEVENSON and LLOYD OSBOURNE. Crown 8vo. 5*s.*

Stock. — *DEDUCTIVE LOGIC.* By ST. GEORGE STOCK. Fcp. 8vo. 3*s.* 6*d.*

Stockton. — *THE GREAT WAR SYNDICATE.* By FRANK R. STOCKTON, Author of 'Rudder Grange.' Fcp. 8vo. 1*s.* sewed.

'Stonehenge.' — *THE DOG IN HEALTH AND DISEASE.* By 'STONEHENGE.' With 84 Wood Engravings. Square crown 8vo. 7*s.* 6*d.*

Stoney. — *THE THEORY OF THE STRESSES ON GIRDERS AND SIMILAR STRUCTURES.* With Practical Observations on the Strength and other Properties of Materials. By BINDON B. STONEY, LL.D. F.R.S. M.I.C.E. With 5 Plates and 143 Illustrations. Royal 8vo. 36*s.*

Sully.—*WORKS BY JAMES SULLY.*

OUTLINES OF PSYCHOLOGY, with Special Reference to the Theory of Education. 8vo. 12*s.* 6*d.*

THE TEACHER'S HANDBOOK OF PSYCHOLOGY, on the Basis of 'Outlines of Psychology.' Crown 8vo. 6*s.* 6*d.*

Sumner.—*THE BESOM MAKER, AND OTHER COUNTRY FOLK SONGS.* Collected and Illustrated by HEYWOOD SUMNER. With Music. 4to. 2*s.* 6*d.* boards.

Supernatural Religion; an Inquiry into the Reality of Divine Revelation. Complete Edition, thoroughly revised. 3 vols. 8vo. 36*s.*

Swinburne. — *PICTURE LOGIC;* an Attempt to Popularise the Science of Reasoning. By A. J. SWINBURNE, B.A. Post 8vo. 5*s.*

Tangena Tree (The): a True Story from Madagascar. By AGNES MARION. Crown 8vo. 1*s.*

Thompson.—*WORKS BY D. GREENLEAF THOMPSON.*

THE PROBLEM OF EVIL: an Introduction to the Practical Sciences. 8vo. 10*s.* 6*d.*

A SYSTEM OF PSYCHOLOGY. 2 vols. 8vo. 36*s.*

THE RELIGIOUS SENTIMENTS OF THE HUMAN MIND. 8vo. 7*s.* 6*d.*

SOCIAL PROGRESS: an Essay. 8vo. 7*s.* 6*d.*

Three in Norway. By Two of THEM. With a Map and 59 Illustrations from Sketches by the Authors. Cr. 8vo. 2*s.* boards; 2*s.* 6*d.* cloth.

Times and Days: being Essays in Romance and History. Fcp. 8vo. 5*s.*

Todd.—*ON PARLIAMENTARY GOVERNMENT IN ENGLAND:* its Origin, Development, and Practical Operation. By ALPHEUS TODD, LL.D. C.M.G. Second Edition. In Two Volumes—VOL. I. 8vo. 24*s.* Vol. II. 8vo. 30*s.*

Tomson.—*THE BIRD BRIDE:* a Volume of Ballads and Sonnets. By GRAHAM R. TOMSON. Fcp. 8vo. 6*s.*

Trevelyan.—*WORKS BY THE RIGHT HON. SIR G. O. TREVELYAN, BART.*

THE LIFE AND LETTERS OF LORD MACAULAY.

POPULAR EDITION, 1 vol. crown 8vo. 2*s.* 6*d.*

STUDENT'S EDITION, 1 vol. cr. 8vo. 6*s.*

CABINET EDITION, 2 vols. cr. 8vo. 12*s.*

LIBRARY EDITION, 2 vols. 8vo. 36*s.*

THE EARLY HISTORY OF CHARLES JAMES FOX. Library Edition, 8vo. 18*s.* Cabinet Edition, crown 8vo. 6*s.*

Trollope.—*NOVELS BY ANTHONY TROLLOPE.*

THE WARDEN. Crown 8vo. 1*s.* boards; 1*s.* 6*d.* cloth.

BARCHESTER TOWERS. Crown 8vo. 1*s.* boards; 1*s.* 6*d.* cloth.

Tuttle.—*HISTORY OF PRUSSIA UNDER FREDERIC THE GREAT,* 1740–1756. By HERBERT TUTTLE. With 2 Maps. 2 vols. crown 8vo. 18*s.*

Twells.—*COLLOQUIES ON PREACHING.* By the Rev. H. TWELLS, M.A. Crown 8vo. 5*s.*

Tyndall.—*WORKS BY JOHN TYNDALL.*

FRAGMENTS OF SCIENCE. 2 vols. crown 8vo. 16*s.*

HEAT A MODE OF MOTION. Cr. 8vo. 12*s.*

[*Continued above.*

Tyndall.—*WORKS BY JOHN TYNDALL, Continued.*

SOUND. With 204 Woodcuts. Crown 8vo. 10*s.* 6*d.*

RESEARCHES ON DIAMAGNETISM AND MAGNE-CRYSTALLIC ACTION. With 8 Plates and numerous Illustrations. Crown 8vo. 12*s.*

ESSAYS ON THE FLOATING-MATTER OF THE AIR in relation to Putrefaction and Infection. With 24 Woodcuts. Crown 8vo. 7*s.* 6*d.*

LECTURES ON LIGHT, delivered in America in 1872 and 1873. With 57 Diagrams. Crown 8vo. 5*s.*

LESSONS IN ELECTRICITY AT THE ROYAL INSTITUTION, 1875–76. With 58 Woodcuts. Crown 8vo. 2*s.* 6*d.*

NOTES OF A COURSE OF SEVEN LECTURES ON ELECTRICAL PHENOMENA AND THEORIES, delivered at the Royal Institution. Crown 8vo. 1*s.* sewed, 1*s.* 6*d.* cloth.

NOTES OF A COURSE OF NINE LECTURES ON LIGHT, delivered at the Royal Institution. Crown 8vo. 1*s.* sewed, 1*s.* 6*d.* cloth.

FARADAY AS A DISCOVERER. Fcp. 8vo. 3*s.* 6*d.*

Unwin.—*THE TESTING OF MATERIALS OF CONSTRUCTION:* a Text-Book for the Engineering Laboratory. By W. CAWTHORNE UNWIN, F.R.S. With 5 Plates and 141 Woodcuts. 8vo. 21*s.*

Vignoles.—*THE LIFE OF C. B. VIGNOLES, F.R.S. SOLDIER AND CIVIL ENGINEER.* Compiled from Original Diaries, Letters, and Documents by his Son, OLINTHUS J. VIGNOLES, M.A. With several Original Illustrations and Portraits. 8vo. 16*s.*

Ville.—*ON ARTIFICIAL MANURES,* their Chemical Selection and Scientific Application to Agriculture. By GEORGES VILLE. Translated and edited by W. CROOKES. With 31 Plates. 8vo. 21*s.*

Virgil.—*PUBLI VERGILI MARONIS BUCOLICA, GEORGICA, ÆNEIS;* the Works of VIRGIL, Latin Text, with English Commentary and Index. By B. H. KENNEDY, D.D. Cr. 8vo. 10*s.* 6*d.*

THE ÆNEID OF VIRGIL. Translated into English Verse. By JOHN CONINGTON, M.A. Crown 8vo. 6*s.*

THE POEMS OF VIRGIL. Translated into English Prose. By JOHN CONINGTON, M.A. Crown 8vo. 6*s.*

Walker. — *THE CORRECT CARD;* or, How to Play at Whist; a Whist Catechism. By Major A. CAMPBELL-WALKER, F.R.G.S. Fcp. 8vo. 2*s.* 6*d.*

Walpole.—*HISTORY OF ENGLAND FROM THE CONCLUSION OF THE GREAT WAR IN* 1815. By SPENCER WALPOLE. 5 vols. 8vo. Vols. I. and II. 1815–1832, 36*s.*; Vol. III. 1832–1841, 18*s.*; Vols. IV. and V. 1841–1858, 36*s.*

Waters. — *PARISH REGISTERS IN ENGLAND:* their History and Contents. By ROBERT E. CHESTER WATERS, B.A. 8vo. 5*s.*

Watts' *DICTIONARY OF CHEMISTRY.* Revised and entirely Re-written by H. FORSTER MORLEY, M.A. D.Sc.; and M. M. PATTISON MUIR, M.A. F.R.S.E. Assisted by Eminent Contributors. To be published in 4 vols. 8vo. Vols. I. & II. A—Indigo. 42*s.* each.

Webb.—*CELESTIAL OBJECTS FOR COMMON TELESCOPES.* By the Rev. T. W. WEBB. Crown 8vo. 9*s.*

Wellington.—*LIFE OF THE DUKE OF WELLINGTON.* By the Rev. G. R. GLEIG, M.A. Crown 8vo. 6*s.*

Wendt. — *PAPERS ON MARITIME LEGISLATION,* with a Translation of the German Mercantile Laws relating to Maritime Commerce. By ERNEST EMIL WENDT, D.C.L. Royal 8vo. £1. 11*s.* 6*d.*

West.—*WORKS BY CHARLES WEST, M.D. &c.*

LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD. 8vo. 18*s.*

THE MOTHER'S MANUAL OF CHILDREN'S DISEASES. Crown 8vo. 2*s.* 6*d.*

Whately. — *WORKS BY E. JANE WHATELY.*

ENGLISH SYNONYMS. Edited by R. WHATELY, D.D. Fcp. 8vo. 3*s.*

LIFE AND CORRESPONDENCE OF RICHARD WHATELY, D.D., late Archbishop of Dublin. With Portrait. Crown 8vo. 10*s.* 6*d.*

Whately.—*WORKS BY R. WHATELY, D.D.*

ELEMENTS OF LOGIC. Cr. 8vo. 4*s.* 6*d.*

ELEMENTS OF RHETORIC. Crown 8vo. 4*s.* 6*d.*

LESSONS ON REASONING. Fcp. 8vo. 1*s.* 6*d.*

BACON'S ESSAYS, with Annotations. 8vo. 10*s.* 6*d.*

Wilcocks.—*THE SEA FISHERMAN.* Comprising the Chief Methods of Hook and Line Fishing in the British and other Seas, and Remarks on Nets, Boats, and Boating. By J. C. WILCOCKS. Profusely Illustrated. Crown 8vo. 6*s.*

Wilks.—*LECTURES ON PATHOLOGICAL ANATOMY.* By SAMUEL WILKS, M.D. F.R.S. and the late WALTER MOXON, M.D. F.R.C.P. Third Edition, thoroughly Revised. By SAMUEL WILKS, M.D. LL.D. F.R.S. 8vo. 18*s.*

Williams.—*PULMONARY CONSUMPTION;* its Etiology, Pathology, and Treatment. By C. J. B. WILLIAMS, M.D. and CHARLES THEODORE WILLIAMS, M.A. M.D. Oxon. With 4 Coloured Plates and 10 Woodcuts. 8vo. 16*s.*

Williams. — *MANUAL OF TELEGRAPHY.* By W. WILLIAMS, Superintendent of Indian Government Telegraphs. Illustrated by 93 Engravings. 8vo. 10*s.* 6*d.*

Williams. — *THE LAND OF MY FATHERS:* a Story of Life in Wales. By T. MARCHANT WILLIAMS. Cr. 8vo. 2*s.* 6*d.*

Willich. — *POPULAR TABLES* for giving Information for ascertaining the value of Lifehold, Leasehold, and Church Property, the Public Funds, &c. By CHARLES M. WILLICH. Edited by H. BENCE JONES. Crown 8vo. 10*s.* 6*d.*

Willoughby. — *EAST AFRICA AND ITS BIG GAME.* By Capt. Sir JOHN WILLOUGHBY, Bart. With Illustrations. 8vo.

Wilson.—*A MANUAL OF HEALTH-SCIENCE.* Adapted for Use in Schools. By ANDREW WILSON, F.R.S.E. F.L.S. &c. With 74 Illustrations. Crown 8vo. 2*s.* 6*d.*

Witt.—*WORKS BY PROF. WITT.* Translated by FRANCES YOUNGHUSBAND.

THE TROJAN WAR. Crown 8vo. 2*s.*

MYTHS OF HELLAS; or, Greek Tales. Crown 8vo. 3*s.* 6*d.*

THE WANDERINGS OF ULYSSES. Crown 8vo. 3*s.* 6*d.*

Wood.—*Works by Rev. J. G. Wood.*

Homes Without Hands; a Description of the Habitations of Animals, classed according to the Principle of Construction. With 140 Illustrations. 8vo. 10*s.* 6*d.*

Insects at Home; a Popular Account of British Insects, their Structure, Habits, and Transformations. With 700 Illustrations. 8vo. 10*s.* 6*d.*

Insects Abroad; a Popular Account of Foreign Insects, their Structure, Habits, and Transformations. With 600 Illustrations. 8vo. 10*s.* 6*d.*

Bible Animals; a Description of every Living Creature mentioned in the Scriptures. With 112 Illustrations. 8vo. 10*s.* 6*d.*

Strange Dwellings; a Description of the Habitations of Animals, abridged from 'Homes without Hands.' With 60 Illustrations. Crown 8vo. 5*s.*

Illustrated Stable Maxims. To be hung in Stables for the use of Grooms, Stablemen, and others who are in charge of Horses. On Sheet, 4*s.*

Out of Doors; a Selection of Original Articles on Practical Natural History. With 11 Illustrations. Crown 8vo. 5*s.*

Petland Revisited. With 33 Illustrations. Crown 8vo. 7*s.* 6*d.*

The following books are extracted from the foregoing works by the Rev. J. G. Wood:

Social Habitations and Parasitic Nests. With 18 Illustrations. Crown 8vo. 2*s.* cloth extra, gilt edges.

The Branch Builders. With 28 Illustrations. Crown 8vo. 2*s.* 6*d.* cloth extra, gilt edges.

Wild Animals of the Bible. With 29 Illustrations. Crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.

Domestic Animals of the Bible. With 23 Illustrations. Crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.

Bird-Life of the Bible. With 32 Illustrations. Crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.

Wonderful Nests. With 30 Illustrations. Crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.

Homes Under the Ground. With 28 Illustrations. Crown 8vo. 3*s.* 6*d.* cloth extra, gilt edges.

Wright.—*Hip Disease in Childhood,* with Special Reference to its Treatment by Excision. By G. A. Wright, B.A. M.B.Oxon. F.R.C.S.Eng. With 48 Original Woodcuts. 8vo. 10*s.* 6*d.*

Wylie.—*History of England under Henry the Fourth.* By James Hamilton Wylie, M.A. one of Her Majesty's Inspectors of Schools. (2 vols.) Vol. I, crown 8vo. 10*s.* 6*d.*

Youatt.—*Works by William Youatt.*

The Horse. Revised and enlarged by W. Watson, M.R.C.V.S. 8vo. Woodcuts, 7*s.* 6*d.*

The Dog. Revised and enlarged. 8vo. Woodcuts. 6*s.*

Younghusband.—*Works by Frances Younghusband.*

The Story of Our Lord, told in Simple Language for Children. With 25 Illustrations on Wood from Pictures by the Old Masters, and numerous Ornamental Borders, Initial Letters, &c. from Longmans' Illustrated New Testament. Crown 8vo. 2*s.* 6*d.*

The Story of Genesis. Crown 8vo. 2*s.* 6*d.*

Zeller.—*Works by Dr. E. Zeller.*

History of Eclecticism in Greek Philosophy. Translated by Sarah F. Alleyne. Crown 8vo. 10*s.* 6*d.*

The Stoics, Epicureans, and Sceptics. Translated by the Rev. O. J. Reichel, M.A. Crown 8vo. 15*s.*

Socrates and the Socratic Schools. Translated by the Rev. O. J. Reichel, M.A. Crown 8vo. 10*s.* 6*d.*

Plato and the Older Academy. Translated by Sarah F. Alleyne and Alfred Goodwin, B.A. Crown 8vo. 18*s.*

The Pre-Socratic Schools: a History of Greek Philosophy from the Earliest Period to the time of Socrates. Translated by Sarah F. Alleyne. 2 vols. crown 8vo. 30*s.*

Outlines of the History of Greek Philosophy. Translated by Sarah F. Alleyne and Evelyn Abbott. Crown 8vo. 10*s.* 6*d.*

EPOCHS OF ANCIENT HISTORY.

Edited by the Rev. Sir G. W. Cox, Bart. M.A. and by C. SANKEY, M.A. 10 volumes, fcp. 8vo. with Maps, price 2*s.* 6*d.* each.

THE GRACCHI, MARIUS, AND SULLA. By A. H. BEESLY, M.A. With 2 Maps.

THE EARLY ROMAN EMPIRE. From the Assassination of Julius Cæsar to the Assassination of Domitian. By the Rev. W. WOLFE CAPES, M.A. With 2 Maps.

THE ROMAN EMPIRE OF THE SECOND CENTURY, or the Age of the Antonines. By the Rev. W. WOLFE CAPES, M.A. With 2 Maps.

THE ATHENIAN EMPIRE FROM THE FLIGHT of Xerxes to the Fall of Athens. By the Rev. Sir G. W. COX, Bart. M.A. With 5 Maps.

THE RISE OF THE MACEDONIAN EMPIRE. By ARTHUR M. CURTEIS, M.A. With 8 Maps.

THE GREEKS AND THE PERSIANS. By the Rev. Sir G. W. Cox, Bart. M.A. With 4 Maps.

ROME TO ITS CAPTURE BY THE GAULS. By WILHELM IHNE. With a Map.

THE ROMAN TRIUMVIRATES. By the Very Rev. CHARLES MERIVALE, D.D. Dean of Ely. With a Map.

THE SPARTAN AND THEBAN SUPREMACIES. By CHARLES SANKEY, M.A. With 5 Maps.

ROME AND CARTHAGE, THE PUNIC WARS. By R. BOSWORTH SMITH, M.A. With 9 Maps and Plans.

EPOCHS OF MODERN HISTORY.

Edited by C. COLBECK, M.A. 19 volumes, fcp. 8vo. with Maps, price 2*s.* 6*d.* each.

THE BEGINNING OF THE MIDDLE AGES. By the Very Rev. RICHARD WILLIAM CHURCH, M.A. &c. Dean of St. Paul's. With 3 Maps.

THE NORMANS IN EUROPE. By Rev. A. H. JOHNSON, M.A. With 3 Maps.

THE CRUSADES. By the Rev. Sir G. W. COX, Bart. M.A. With a Map.

THE EARLY PLANTAGENETS. By the Right Rev. W. STUBBS, D.D. Bishop of Oxford. With 2 Maps.

EDWARD THE THIRD. By the Rev. W. WARBURTON, M.A. With 3 Maps.

THE HOUSES OF LANCASTER AND YORK; with the Conquest and Loss of France. By JAMES GAIRDNER. With 5 Maps.

THE EARLY TUDORS. By the Rev. C. E. MOBERLY, M.A.

THE ERA OF THE PROTESTANT REVOLUTION. By F. SEEBOHM. With 4 Maps.

THE AGE OF ELIZABETH. By the Rev. M. CREIGHTON, M.A. LL.D. With 5 Maps.

THE FIRST TWO STUARTS AND THE PURITAN REVOLUTION, 1603–1660. By SAMUEL RAWSON GARDINER. With 4 Maps.

THE THIRTY YEARS' WAR, 1618–1648. By SAMUEL RAWSON GARDINER. With a Map.

THE ENGLISH RESTORATION AND LOUIS XIV. 1648–1678. By OSMUND AIRY.

THE FALL OF THE STUARTS; AND WESTERN EUROPE from 1678 *to* 1697. By the Rev. EDWARD HALE, M.A. With 11 Maps and Plans.

THE AGE OF ANNE. By E. E. MORRIS, M.A. With 7 Maps and Plans.

THE EARLY HANOVERIANS. By E. E. MORRIS, M.A. With 9 Maps and Plans.

FREDERICK THE GREAT AND THE SEVEN YEARS' WAR. By F. W. LONGMAN. With 2 Maps.

THE WAR OF AMERICAN INDEPENDENCE, 1775–1783. By J. M. LUDLOW. With 4 Maps.

THE FRENCH REVOULTION, 1789–1795. By Mrs. S. R. GARDINER. With 7 Maps.

THE EPOCH OF REFORM, 1830–1850. By JUSTIN MCCARTHY, M P.

EPOCHS OF CHURCH HISTORY.

Edited by the Rev. MANDELL CREIGHTON. 16 volumes, fcp. 8vo. price 2*s.* 6*d.* each.

THE ENGLISH CHURCH IN OTHER LANDS. By the Rev. H. W. TUCKER.

THE HISTORY OF THE REFORMATION IN England. By the Rev. GEORGE G. PERRY.

THE CHURCH OF THE EARLY FATHERS. By ALFRED PLUMMER, D.D.

THE EVANGELICAL REVIVAL IN THE Eighteenth Century. By the Rev. J. H. OVERTON.

A HISTORY OF THE UNIVERSITY OF OXFORD. By the Hon. G. C. BRODRICK, D.C.L.

A HISTORY OF THE UNIVERSITY OF CAMbridge. By J. BASS MULLINGER, M.A.

THE ENGLISH CHURCH IN THE MIDDLE Ages. By Rev. W. HUNT, M.A.

THE ARIAN CONTROVERSY. By H. M. GWATKIN, M.A.

THE COUNTER-REFORMATION. By A. W. WARD.

THE CHURCH AND THE ROMAN EMPIRE. By the Rev. A. CARR.

THE CHURCH AND THE PURITANS, 1570–1660. By HENRY OFFLEY WAKEMAN.

THE CHURCH AND THE EASTERN EMPIRE. By the Rev. H. F. TOZER.

HILDEBRAND AND HIS TIMES. By the Rev. W. R. W. STEPHENS.

THE POPES AND THE HOHENSTAUFEN. By UGO BALZANI.

THE GERMAN REFORMATION. By Prof. MANDELL CREIGHTON.

WYCLIFFE AND EARLY MOVEMENTS FOR Reform. By R. LANE POOLE.

Spottiswoode & Co. Printers, New-street Square, London.